
4: SOILS

A. Soil Survey Maps

The *soil* is the unconsolidated mineral material on the immediate surface of the earth which serves as the medium for growth of land plants. The characteristics of each soil type have developed over time (usually many thousands of years) under the influence of the parent material (the bedrock that has broken down into small fragments to form the soil), climate (including moisture and temperature regimes), macro- and microorganisms, and topography. Soil is a basic resource for food production, in addition to its essential role in collecting and purifying water before it enters the ground water (Soil Science Society of America, 2017). However, soil itself can be a pollutant as dust in the air or as sediment in water.



The US Department of Agriculture Natural Resources Conservation Service (USDA NRCS) is the science-based agency which provides technical assistance regarding the conservation and management of soil, water, and other natural resources to private land owners, local, state, and federal agencies, and policy-makers (USDA NRCS, February 10, 2011).

One of these technical services is the soil survey. A *soil survey* is an inventory of the country's soil resources to determine soil characteristics and capabilities and to help people understand soils and their uses. Soil surveys help to identify the best ways to protect soil and water quality through the use of conservation practices, and to identify which sites are suitable (and the degree of suitability) for various land uses (e.g. septic systems, roads, agriculture).

The objective of soil mapping is to separate the landscape into segments that have similar use and management requirements. Therefore, this data set is not designed for use as a primary regulatory or management tool, but may be used as a broad scale reference source. According to the Soil Survey Geographic Database (also known as SSURGO) information, field investigations and data collection were carried out in sufficient detail to name map units and to identify accurately and consistently areas of about 5 acres. As with other GIS data sets, enlargement of the maps to a scale greater than the accuracy of the data can cause misinterpretation of the data. Onsite sampling, testing, and detailed study of specific sites is essential for determining intensive uses, and for managing farms and wetlands (USDA NRCS, February 17, 2012).

Beginning in 2005, the NRCS made its soil surveys available online (USDA NRCS, February 17, 2012). This provides the means for keeping the information current and available to the public. Users specify a geographic "area of interest" (must be less than 10,000 acres) and then may view a wide variety of tables of soil properties and soil interpretations. For this report,

the entire SSURGO (Soil Survey Geographic Database) spatial data and tabular data for Cape May County were downloaded for use in the GIS (USDA NRCS, September 28, 2016)¹⁵.

B. Soil Series and Map Units

Soil characteristics vary from place to place in slope, depth, drainage, erodibility and other characteristics that affect management. A *soil series* is a basic unit of soil classification consisting of soils that are essentially alike, except that they may differ in surface texture, stoniness, slope or some other attribute. A *map unit* is the area delineated on a soil map, representing an area dominated by one major kind of soil, and is named according to the classification of the dominant soil or soils. However, soils are natural systems, with natural variability, and the range of some observed properties may extend beyond the limits defined for the class. In addition, small areas of contrasting soils may not be visible on the maps. The databases included with the soils data describe the characteristics of each soil map unit. The NRCS has included both estimated and measured data on the physical and chemical soil properties and soil interpretations for engineering, water management, recreation, agronomic, woodland, range and wildlife uses of the soil.

There are 17 soil map units found in the Borough of West Cape May (**Figure 4a**), 15 of which are soil types (80% of land cover in the borough). The remaining units indicate locations where soils have either been removed (PHG) or covered by impervious materials (UR) (collectively 20% of the borough). The map units are described below, and the total acreage for each unit is provided. **Table 4.1** also summarizes some important properties of each soil type.

Map Unit Descriptions and Acreage

Sources: US Department of Agriculture, Natural Resource Conservation Service (2002 and August 2008).

Appoquinimink-Transquaking-Mispillion complex, 0 to 1 percent slopes, very frequently flooded (AptAv - 29 acres)

Very poorly drained soil with a high water capacity and a moderate to high organic content in the surface layer. Moderately to strongly saline within 30 inches of surface. Tidal marshes, tidal flats and drainageways.

Berryland and Mullica soils, 0 to 2 percent slopes, occasionally flooded (BEXAS - 160 acres)

Very poorly drained soil with a moderate water capacity and moderate organic content in the surface layer. Freshwater flats, floodplains, drainageways and depressions.

Dennisville sandy loam, 0 to 2 percent slopes (DenA - 10 acres)

Well drained soil with a low water capacity and low organic content in the surface layer. Broad ridges, summits and side slopes.

Downer sandy loam, 0 to 2 percent slopes, Northern Tidewater Area (DoeAO - 9 acres)

Well drained soil with a moderate water capacity and low organic content in the surface layer. Broad ridges, summits and side slopes.

Fort Mott sand, 0 to 5 percent slopes (FobB - 5 acres)

Well drained soil with a low or moderate water capacity and low organic content in the surface layer. Ridges, terraces, summits and side slopes.

¹⁵ The maps in this report use the most recent data available (Cape May County, NJ; Tabular Data US Department of Agriculture, Natural Resources Conservation Service (NRCS). September 28, 2016. Survey Area Version: 12. Survey Area Version Date: 09/28/2016.

Galloway loamy sand, 0 to 5 percent slopes (GamB - 23 acres)

Moderately well drained soil with a low water capacity and low organic content in the surface layer.

Terraces, flats, side slopes and depressions.

Hammonton loamy sand, 0 to 5 percent slopes (HbmB - 92 acres)

Moderately well drained soil with a moderate water capacity and low organic content in the surface layer.

Terraces, flats, depressions, side slopes and foot slopes.

Hammonton sandy loam, 0 to 2 percent slopes (HboA - 21 acres)

Moderately well drained soil with a moderate water capacity and low organic content in the surface layer.

Terraces, flats, depressions, side slopes and foot slopes.

Ingleside loamy sand, 0 to 5 percent slopes (IngB - 50 acres)

Well drained soil with a moderate water capacity and low organic content in the surface layer.

Flats, terraces, summits and side slopes.

Ingleside sandy loam, 0 to 2 percent slopes (InnA - 113 acres)

Well drained soil with a moderate water capacity and low organic content in the surface layer.

Flats, terraces, summits and side slopes.

Pawcatuck-Transquaking complex, 0 to 1 percent slopes, very frequently flooded (PdwAv - 51 acres)

Very poorly drained soil with a very high water capacity and very high organic content in the surface layer.

Strongly saline within 30 inches of surface. Tidal marshes, tidal flats and drainageways.

Pits, sand and gravel (PHG - 37 acres)

Not a soil type. Pits, sand and gravel are open areas from which soil material has been removed by human activity.

They often have steep, unstable slope faces.

Psammaquents, sulfidic substratum, 0 to 3 percent slopes, frequently flooded (PstAt - 8 acres)

Fill material in tidal marshes, ranging from 2 to 4 feet in thickness, with fine materials in the upper part and silty organic materials in the lower part.

Psamments, wet substratum, 0 to 3 percent slopes, rarely flooded (PsvAr - 30 acres)

Fill material in floodplains, flats and swamps, ranging from 2 to 4 feet in thickness, with fine materials in the upper part and typically more loamy or mucky organic materials in the lower part.

Udorthents, refuse substratum, 0 to 8 percent slopes (UdrB - 5 acres)


Sanitary landfills on convex summits, side slopes or broad flats; typically a clay surface above a sandy or loamy material mixed with household or industrial refuse.

Urban land (UR - 110 acres)

Not a soil type. Urban land is land where most of the soil surface is covered by asphalt, concrete, buildings, or other impervious material.

Urban land-Psamments, wet substratum complex, 0 to 8 percent slopes, rarely flooded (USPSBR - less than an acre)

Urban land developed over floodplains, flats, swamps or barrier islands that have been altered or filled. Generally sandy material of variable characteristics that has been placed on top of wet soils.




Legend

Parcels	Map Unit Symbol	GamB	PdwAv
West Cape May	AptAv	HbmB	PstAt
Roads	BEXAS	HboA	PsvAr
Water	DenA	IngB	UR
Streams	DoeAO	InnA	USPSBR
	FobB	PHG	UdrB

Figure 4a: West Cape May Soil Map Units

There are 17 soil map units found in the borough, 15 of which are soil types. The remaining units indicate locations where soils have either been removed (PHG) or covered by impervious materials (UR).



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Table 4.1: Key Characteristics of Soil Types Found in West Cape May

Map unit symbol	Map unit name	Depth to root-restrictive layer (inches)	Depth to seasonal high water table (feet)	Flooding	Ponding	Potential frost action	Hydrologic group (defined in Table 4.2)	Natural drainage class	Hydric soil?	Farmland rating class	Septic disposal field rating class (NJ)	Septic limiting features (NJ)*
AptAv	Appoquinimink-Transquaking-Mispillion complex, 0 to 1 percent slopes, very frequently flooded	>60	0	very frequent	frequent	high	B/C/D	very poorly drained	Yes	Unique	very limited	DAZS, NPF, NPHS
BEXAS	Berryland and Mullica soils, 0 to 2 percent slopes, occasionally flooded	>60	0.0-0.5	occasional	occasional	high/moderate	A/D	very poorly drained	Yes	Unique	very limited	DAZS, NPF, NPHS
DenA	Dennisville sandy loam, 0 to 2 percent slopes	>60	3.5-4.5	none	none	moderate	A	well drained	No	Prime	somewhat limited	DAZS
DoeAO	Downer sandy loam, 0 to 2 percent slopes, Northern Tidewater Area	>60	> 6	none	none	moderate	A	well drained	No	Prime	not limited	-
FobB	Fort Mott sand, 0 to 5 percent slopes	>60	4.0->6	none	none	low	A	well drained	No	Statewide	not limited	-
GamB	Galloway loamy sand, 0 to 5 percent slopes	>60	1.0-1.5	none	none	low	A/D	somewhat poorly drained	No	Statewide	very limited	DAZS
HbmB	Hammonton loamy sand, 0 to 5 percent slopes	>60	1.5-3.5	none	none	moderate	B	moderately well drained	No	Statewide	somewhat limited	DAZS
HboA	Hammonton sandy loam, 0 to 2 percent slopes	>60	1.5-3.5	none	none	moderate	B	moderately well drained	No	Prime	somewhat limited	DAZS
IngB	Ingleside loamy sand, 0 to 5 percent slopes	>60	3.5-4.5	none	none	moderate	A	well drained	No	Prime	somewhat limited	DAZS
InnA	Ingleside sandy loam, 0 to 2 percent slopes	>60	3.5-4.5	none	none	moderate	A	well drained	No	Prime	somewhat limited	DAZS
PdwAv	Pawcatuck-Transquaking complex, 0 to 1 percent slopes, very frequently flooded	>60	0	very frequent	frequent	high	A/D	very poorly drained	Yes	Unique	very limited	DAZS, NPF, NPHS,
PHG	Pits, sand and gravel	areas where soil has been removed										
PstAt	Psammaquents, sulfidic substratum, 0 to 3 percent slopes, frequently flooded	>60	0	frequent	frequent	moderate	A/D	very poorly drained	Yes	Prime	very limited	DAZS, NPF, NPHS

Map unit symbol	Map unit name	Depth to root-restrictive layer (inches)	Depth to seasonal high water table (feet)	Flooding	Ponding	Potential frost action	Hydrologic group (defined in Table 4.2)	Natural drainage class	Hydric soil?	Farmland rating class	Septic disposal field rating class (NJ)	Septic limiting features (NJ)*
PsvAr	Psamments, wet substratum, 0 to 3 percent slopes, rarely flooded	>60	1.5-3.5	rare	none	low	A	moderately well drained	No	Prime	very limited	DAZS, NPF
UdrB	Udorthents, refuse substratum, 0 to 8 percent slopes	>60	> 6	none	none	high	B	well drained	No	Prime	not limited	-
UR	Urban land	areas where soil has been covered by impervious materials										
USPSBR	Urban land-Psamments, wet substratum complex, 0 to 8 percent slopes, rarely flooded	>60	1.0-2.0	rare	none	low	A/D	excessively drained	No	Prime	not rated to very limited	DAZS, NPF
* Septic System Limiting Factors: DAZS = Depth to apparent zone of saturation; NPF = Not Permitted - Flooding; NPHS = Not Permitted - Hydric Soil												
Sources: USDA NRCS, August 2008 and September 2016												

C. Soil Quality

Soil is arranged in horizontal layers called horizons. These horizons have technical designations largely useful for soil scientists to distinguish one soil series from another. The descriptions in the NRCS soil survey are done using soil in its native state where possible, so a soil profile which has been disturbed may not match the written description for the series. This is the way the degree of disturbance is assessed—by comparing the soil in its native condition to the profile observed at a specific site. For example, the upper horizon is often an *A horizon*, commonly known as “*topsoil*.” An *A horizon* typically exhibits increased organic matter, reduced clay percentage, a more granular structure of the soil aggregates, and a lower bulk density than the *B horizon* below it. If the *A horizon* is removed (a common practice in construction), this is evident to a trained observer and the soil would be described as having the *A horizon* missing. The material on the new surface does not automatically become an *A horizon* merely as a result of its position. It is possible over time for the newly exposed surface to acquire the characteristics of an *A horizon*, however this is not automatic and is highly management dependent. In technical writing, particularly in guidance documents intended for post-construction remediation, the use of the term “*topsoil*” should be used with caution if at all because there is no legal definition of *topsoil* and the materials available in commerce are highly variable in quality (Muldowney, 2011).

Soils vary naturally in their capacity to function. *Soil quality* is defined as the capacity of a specific kind of soil to function to sustain plant and animal productivity, maintain or enhance water and air quality, and support human health and habitation. *Inherent* or *intrinsic soil qualities* or characteristics of the soil are determined by factors of soil formation (climate, parent material, topography, time and biota). These are properties which cannot be altered by management except by actually replacing the present material with a different material altogether. An example of an inherent property is the percentage of sand in the soil's composition. Inherent soil quality is used to evaluate the suitability of soils for specific uses (buildings, roads, agriculture, septic systems, etc.). One measure of quality is soil particle size:

A loamy soil will have higher water holding capacity than a sandy soil, and therefore will have a higher inherent quality for storing water (USDA NRCS, November 2010).

Contrasting with intrinsic soil properties are management-dependent soil properties, also known as *dynamic soil qualities*. As the term suggests, these can be altered significantly (for better or for worse) by the management of a specific parcel of land, and changes can have significant consequences for overall environmental quality. Dynamic quality is determined by soil characteristics that are affected by human use and management practices, including physical, chemical and biological properties. Soil quality or health may be evaluated by either comparison to a reference condition that represents full capacity of a soil for a specific function, or to a baseline for the management-dependent soils properties (such as before and after a land use change) (USDA NRCS, November 2010).

Degradation of soil quality occurs in many forms. Significant issues are cutting and filling, compaction, excess salt content and organic matter content. *Cutting and filling* operations actually remove, bury, or invert existing horizons such that they no longer behave in a hydrologically coherent way, with precipitation and gases readily able to enter the soil surface and transmit to horizons lower in the profile. *Compaction*, the increase of bulk density as a result of compression from the surface, is another common form of soil degradation. Compaction can be avoided by not working soil at too high a moisture content. Even foot traffic on a near saturated soil can result in lasting damage which does not resolve itself naturally. A compacted soil can have runoff characteristics more similar to pavement than to the soil in good condition (Muldowney, 2011).

Excess salt content often results from deicing salts but sometimes from fertilizer preparations. It is especially common on roadside verges. The remedy is either prevention or washing the salt from the profile with excess water. Sodium salts are especially damaging because sodium causes the clays to disperse. Calcium chloride is relatively harmless to plants and soil (Muldowney, 2011; Wikipedia, May 30, 2013).

Organic matter content is another dynamic soil property. Rutgers New Jersey Agricultural Experiment Station (see **Internet Resources**) provides a chart for interpretation of organic matter percentages in New Jersey soils. Soils with a high organic content are better able to resist other forms of degradation than soils with depleted organic matter. The amount of organic matter in the upper horizons of a soil is a measure of carbon storage. Soil is the largest terrestrial reservoir of carbon and has the greatest potential for long term storage if degraded soils are managed in a way that builds up carbon. Silt loams, like the Preakness, are able to store more carbon in the form of organic matter than sandier soil. Keeping soil in good condition reduces runoff, produces cleaner runoff, requires less irrigation, grows more robust plantings, and sequesters more atmospheric carbon than a damaged soil (Muldowney, 2011).

D. Characteristics of West Cape May Soils

Soil properties contained in the NRCS soil survey and mapped in **Figures 4b through 4g** are *intrinsic* soil properties. These are properties which cannot be altered by management except by actually replacing the present material with a different material altogether.

Depth to Bedrock

According to NJDEP (1999), *bedrock* is defined as "any solid body of rock, with or without fractures, which is not underlain by soil or unconsolidated rock material." The *depth to bedrock* is the distance from the land surface to bedrock. Each soil map unit is characterized by a range of depths to bedrock that is typical for the majority of that soil type. Depth to bedrock is

an important factor when determining the suitability of land for building roads, foundations and septic systems. Bedrock outcrops (depth to bedrock equals zero inches) are absent in West Cape May. All of the soil units mapped in the Borough have a depth to the root-restrictive layer exceeding five feet (see **Table 4.1**).

Depth to Seasonal High Water Table (Figure 4b)

The *depth to seasonal high water table* (SHWT) is the distance between the ground surface and the top of the water surface in the saturated part of a water bearing zone. A SHWT of less than one foot severely constrains development, and a SHWT between 1 and 3 feet also provides obstacles to development. On-site investigation will often reveal that these areas are actually wetlands or floodplains. High water tables impact the effectiveness of septic systems, and the freeze/thaw cycles cause frost heaving, which damages structures and roads.

Four of the soil units in West Cape May fall into the first category, with SHWTs of less than a foot. Those include the Appoquinimink-Transquaking-Mispillion complex (AptAv), the Berryland and Mullica soils (BEXAS), the Pawcatuck-Transquaking complex (PdwAv), and the Psammaquents sulfidic substratum (PstAt). Together they comprise 41% of the soil cover in the borough. Another five units, which collectively comprise 24% of the borough's total soil cover, have SHWTs that range between 1.0-3.5 feet: Galloway loamy sand (GamB), Hammonton loamy sand and sandy loam (HbmB, HboA), Psammets wet substratum (PsvAr), and Urban land psammets - wet substratum complex (USPSBR). The remaining 35% of the soil cover in West Cape May has a minimum depth to seasonal high water table exceeding three feet (see **Table 4.1** and **Figure 4b**). Non-soil units were excluded from the percentage calculations in this section.

Hydrologic Soil Group (Figure 4c)

The *hydrologic soil grouping* describes a group of soils having similar runoff potential under similar storm and cover conditions (how much water would runoff compared to the rate that water would infiltrate into the ground). The definitions of the hydrologic soil groups are shown in **Table 4.2**.

Table 4.2: Hydrologic Soil Grouping

Class	Definition
A	High infiltration rates. Soils are deep, well to excessively drained sands and gravels.
B	Moderate infiltration rates. Deep and moderately deep, moderately well and well drained soils that have moderately coarse textures.
C	Slow infiltration rates. Soils with layers impeding downward movement of water, or soils that have moderately fine or fine textures.
D	Very slow infiltration rates. Soils are clayey, have a high water table, or are shallow to an impervious layer.
Source: USDA NRCS, 2008	

Six of the soil units in West Cape May fall into Class A, and three others into Class B (**Table 4.1**). The remaining six units have been assigned to multiple hydrologic soil groups. Dual ranks are indicative of soils that respond differently under varying hydrological conditions. In those cases, the first letter applies to the soil when it is in a drained condition (seasonal high water table at least two feet below the soil surface), and the last letter shows how the soil functions when the water table is higher (USDA - NRCS, 2007) (see **Figure 4c**).

Soil Drainage Class (Figure 4d)

Soil Drainage Class is a code identifying the natural drainage condition of the soil and refers to the frequency and duration of periods when the soil is free of saturation or partial saturation during soil formation. It does not refer to saturation due to recently altered drainage (manmade or natural). The categories are as follows: well drained, moderately well drained, excessively drained, somewhat excessively drained, poorly drained, and somewhat poorly drained. West Cape May has six soil types that are well drained, three that are moderately well drained, one that is somewhat poorly drained and three that are very poorly drained (see **Table 4.1**). The Urban land Psammets wet substratum complex, where wetlands have been filled in, is classified as excessively drained. Non-soil map units are also included in the figure in accordance with their runoff potential: Pit, sand and gravel (PHG) areas are very poorly drained and urban land (UR) is excessively drained. In total, 38% of the land in West Cape May is very poorly drained, 3% is poorly drained, 19% is moderately well drained, 25% is well drained and 15% is excessively drained (**Figure 4d**).


Hydric and Flooded Soils

Hydric soils are those soils that are wet long enough to periodically produce anaerobic conditions, thereby influencing the growth of plants. For delineation of hydric soils the ponding event must last longer than seven days. The four hydric soils found within West Cape May are the Appoquinimink-Transquaking-Mispiration complex (AptAv), the Berryland and Mullica soils (BEXAS), the Pawcatuck-Transquaking complex (PdwAv), and the Psammaquents sulfidic substratum (PstAt) (see **Table 4.1**). Collectively these map units comprise 33% of the borough. **Figure 6d** shows wetlands and hydric soils.

Annual flood frequency is a descriptive term used to describe the frequency of flooding that is likely to occur in a year. **Frequent** is > 50% chance of flooding in a given year; **occasional** is 5 to 50%; **rare** is 0 to 5% chance of flooding. In West Cape May, the hydric soils listed in the preceding paragraph are those most likely to experience flooding or ponding (see **Table 4.1**). **Figure 6c** shows floodplains, which encompass the frequently flooded soils.


Potential Frost Action (Figure 4e)

Potential Frost Action is an interpretational rating of the susceptibility of the soil to frost heaving. About half of the soil units mapped within West Cape May have moderate potential frost action, and those units account for 50% of the soil cover in the borough (**Table 4.1**). High potential for frost action is found in the Appoquinimink-Transquaking-Mispiration complex (AptAv), the Pawcatuck-Transquaking complex (PdwAv), and the Urdothents refuse substratum (UdrB), which together comprise 14% of the borough's soil cover. Berryland and Mullica soils (BEXAS), the most extensive map unit in West Cape May accounting for about a quarter of the borough's soil cover (26%), has a high/moderate potential for frost action. Conversely, the lowest potential for frost heave is found in the 10% of the borough's soil cover which includes Fort Mott sand (FobB), Galloway loamy sand (GamB), Psammets wet substratum (PsvAr) and Urban land Psammets wet substratum complex (USPSBR) (see **Figure 4e**). Non-soil units were excluded from the percentage calculations in this section.



**Figure 4b: West Cape May
Depth to High Water Table (minimum)**

The depth to a seasonal high water table is the distance between the ground surface and the top of the water surface in the saturated zone. A depth of less than one foot severely constrains development, and a depth between one and three feet also provides obstacles to development.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.






Figure 4c: West Cape May Hydrologic Group

Hydrologic soil groupings indicate the infiltration rates and runoff potential for various soil groups. Class A soils have the highest rates of infiltration, while Class D soils have the slowest.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.



Legend


- West Cape May
- Parcels
- Roads
- Water
- Streams

Drainage Class


- excessively drained
- moderately well drained
- well drained
- somewhat poorly drained
- very poorly drained

**Figure 4d: West Cape May
Drainage Class**

Soils in West Cape May range from well-drained to very poorly drained. Soils mapped as Excessively Drained indicate locations where wetlands have been filled in.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.



**Figure 4e: West Cape May
Potential Frost Action**


Potential Frost Action is an interpretational rating of the susceptibility of the soil to frost heaving. About half of the soil units mapped within West Cape May have a moderate ranking.

Legend

- West Cape May
- Parcels
- Roads
- Water
- Streams

Potential Frost Action

- High
- Moderate to High
- Moderate
- Low



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Farmland Suitability (Figure 4f)

New Jersey uses standard categories of soil quality developed by the USDA to rank the relative value of land for farming purposes, as described in **Table 4.3** (NJ SADC/CADB, 2003). Nine of the soil units in West Cape May are classified as Prime Farmland (40% of soil cover), and three others are ranked as being of Statewide Importance (20% of soil cover) (see **Table 4.1**). In the Unique category (40% of soil cover) are three of the four hydric soil types mapped in the borough: the Appoquinimink-Transquaking-Mispillion complex (AptAv), the Berryland and Mullica soils (BEXAS), and the Pawcatuck-Transquaking complex (PdwAv). None of the soils in West Cape May are classified as locally important (**Figure 4f**). Non-soil units were excluded from the percentage calculations in this section.


Table 4.3: Farmland Rating Classes.

Classification	Description
Prime	This land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber and oilseed crops and is also available for these uses. It has the soil quality, growing season, and moisture supply needed to economically produce sustained high yields of crops when treated and managed according to acceptable farming methods. Soils are not excessively erodible or saturated with water for a long period of time, and they either do not flood frequently or are protected from flooding.
Statewide	Soils of Statewide Importance are nearly Prime, and economically produce high yields of crops when treated and managed according to acceptable farming methods. Some may produce yields as high as Prime Farmland if conditions are favorable.
Local	Farmland of Local Importance includes those soils that are not prime or of statewide importance and are used for the production of high value food, fiber or horticultural crops.
Unique	These are soils with severe limitations which are used to produce special crops (e.g., cranberry bogs)
Source: USDA - NRCS, 2007.	

Septic Suitability (Figure 4g)

The NRCS SSURGO database provides an interpretation of limitations of each soil for *septic suitability*. The interpretation shown in **Figure 4g** is based on the N.J.A.C. 7:9A Standards for Individual Subsurface Sewage Disposal Systems, Subchapter 10 Disposal Fields. Factors which may affect the functioning of the system, and therefore limit septic suitability, are excessively coarse substratum or excessively coarse horizon (which allow effluent to percolate to ground water too rapidly); presence of water (including depth to perched zone of saturation, depth to apparent zone of saturation, flooding, and hydric soils); depth to restrictive layer (bedrock, restrictive horizon or restrictive substratum); and steep grades over 25%. N.J.A.C 7:9A prohibits septic systems in soils subject to flooding. In addition, septic disposal fields are prohibited in locations with the combination of slope greater than 10% and less than 50 feet upslope of any bedrock outcrop where signs of ground water seepage can be detected (NJDEP, 1999).

Three of the soil types in West Cape May do not have limited suitability for septic systems, and together they make up only three percent of the borough's soil cover. Those include the Downer sandy loam (DoeAO), the Fort Mott sand (FobB), and the Urdothents refuse substratum (UdrB). The remainder are classified as having either somewhat limited (47%) or very limited (50%) capacity for septic disposal (**Table 4.1**). Non-soil units were excluded from the percentage calculations in this section.



Legend


- West Cape May
- Parcels
- Roads
- Water
- Streams

Farmland Soils


- All areas are prime farmland
- Farmland of statewide importance
- Farmland of unique importance
- not a farmland soil

Figure 4f: West Cape May Farmland Soils

Soils ranked as *Prime* or of *Statewide Importance* are suitable for growing a wide variety of crops. Soils ranked as *Unique* are primarily wetlands, but may be suitable for certain specialty crops.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.



Legend


- West Cape May
- Parcels
- Roads
- Water
- Streams

Septic Disposal Field Rating Class (NJ)

- Not limited
- Not rated
- Not rated to Very limited
- Somewhat limited
- Very limited

Figure 4g: West Cape May Septic Disposal Field Rating Class (NJ)

The majority of soil types in West Cape May have either somewhat limited or very limited suitability for the development of septic fields. Suitability is typically limited by excessively coarse soils, the presence of water, shallow depth to bedrock, or steep gradients.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

References: Soils

Muldowney, Loren S. November 2011. Personal Communication. Soil Testing Laboratory, Department of Plant Biology and Pathology Rutgers School of Environmental and Biological Sciences Rutgers University.

NJDEP Division of Water Quality, Bureau of Nonpoint Source Pollution Control. August 15, 1999. N.J.A.C. 7:9A Standards for Individual Subsurface Sewage Disposal Systems, Subchapter 10. Disposal Fields. 130 pages. <http://www.state.nj.us/dep/dwq/pdf/njac79a.pdf>

NJ SADC/CADB. 2003. NJ State Agriculture Development Committee/ NJ County Agriculture Development Boards Farmland Preservation Program. Strategic Targeting Project Preliminary Report, March 2003. <http://www.nj.gov/agriculture/sadc/farmpreserve/resources/stp2003report.pdf>

Soil Science Society of America. 2017. Internet Glossary of Soil Science Terms. <https://www.soils.org/publications/soils-glossary>

US Department of Agriculture, Natural Resources Conservation Service (NRCS). September 28, 2016. Sewage Disposal Table for Cape May County, New Jersey. Technical Soil Services Handbook. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_053400. Accessed March 30, 2016.

US Department of Agriculture, Natural Resources Conservation Service (NRCS). September 28, 2016. USDA Soil Survey Geographic (SSURGO) database for Cape May County, New Jersey. Survey Area Version: 12 Survey Area Version Date: 09/28/2016. <http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx>

US Department of Agriculture, Natural Resources Conservation Service (NRCS). February 17, 2012. The Web Soil Survey Site. <http://websoilsurvey.nrcs.usda.gov/app/>

US Department of Agriculture, Natural Resources Conservation Service (NRCS). February 10, 2011. NRCS Soils Website. <http://soils.usda.gov/>

US Department of Agriculture, Natural Resources Conservation Service (NRCS). 2007. Part 630 Hydrology National Engineering Handbook. Chapter 7 - Hydrologic Soil Groups. <https://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=17757.wba>

US Department of Agriculture, Natural Resource Conservation Service (NRCS). 2002. Soil Survey of Cape May County, NJ. https://www.nrcs.usda.gov/Internet/FSE_MANUSCRIPTS/new_jersey/capemayNJ2002/capemayNJ2002.pdf

US Department of Agriculture, Natural Resources Conservation Service (NRCS). Technical Soil Services Handbook (TSSH), Part 624 Soil Quality. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_053400. Accessed March 30, 2016.

Wikipedia. Accessed May 30, 2013. Calcium chloride. http://en.wikipedia.org/wiki/Calcium_chloride

Internet Resources: Soils

NRCS New Jersey Office: <http://www.nj.nrcs.usda.gov/>

NRCS Soils Website: Helping People Understand Soils: <http://soils.usda.gov/>

NRCS Soil Data Mart (download soils data for GIS): <http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx>

NRCS Soils Online Study Guide: http://www.nrcs.usda.gov/wps/portal/nrcs/detail/nj/home/?cid=nrcs141p2_018928

Rutgers New Jersey Agricultural Experiment Station Soil Testing Laboratory Interpretation of Organic Matter Levels in New Jersey Soils: <http://njaes.rutgers.edu/soiltestinglab/pdfs/nj-om-interpret.pdf>

Web Soil Survey (online soils mapping): <http://websoilsurvey.nrcs.usda.gov/app/>

5: GROUND WATER & DRINKING WATER

A. Water Cycle/Ground Water Background

Even though the quantity of water on the earth is great, only a small portion can be used for drinking water and other human needs. Ninety-seven percent of the world's water supply is saltwater stored in the oceans. The remaining 3% is fresh water. However, most of this is unavailable for human use because it is frozen in the polar ice caps, glaciers, and icebergs; too difficult to tap (below 1.6 miles depth); or too polluted. This leaves 0.003% of water that is available as fresh surface or ground water that humans can use (Miller, 1988).

Surface water is water that is visible above the ground surface, such as creeks, rivers, ponds, lakes, and wetlands. *Ground water* is that portion of water beneath the land surface that is within the zone of saturation (below the water table) where pore spaces are filled with water. An *aquifer* is a water-bearing rock or geologic formation (including sediments such as unconsolidated sands) where water is present in usable quantities. Water is constantly recycled through the *hydrologic cycle*, also known as the *water cycle* (see **Figure 5a**). Precipitation falls on the ground and some travels on the surface of the land (called *surface runoff*), entering streams (where it can be seen as high flows after rain events), and eventually making its way back to the ocean. Some of the water from precipitation enters the ground but remains in the shallow layers where it is available for use by plants, and returns to the atmosphere through *transpiration* by plants, while some water re-enters the atmosphere directly through *evaporation* from surface water. Evaporation and transpiration combined are known as *evapotranspiration*. The water that migrates below the root zone travels underground and exits the system as stream flow, known as *ground water baseflow* or *ground water recharge*. Ground-water baseflow can be calculated by measuring stream flow during dry weather conditions. A smaller portion of the water penetrates deeper into the ground and




Figure 5a: The Water Cycle

Source: USGS, no date

enters (or recharges) the saturated zone of the fractured bedrock or other geologic formation, called the *aquifer*, where most wells obtain their water.

Pollutants can enter water as it travels the water cycle. Surface runoff can pick up chemicals and sediment on its way, depositing these pollutants in waterways. This is especially true of “uncontrolled runoff” on soils that are vulnerable to erosion. Water seeping into the soil can be cleansed of many pollutants by natural soil processes. However, if the pollutant is one that is resistant to break-down, or if the pollutant doesn’t get exposed to the soil long enough (such as by entering a bedrock fracture or by entering the ground water through sub-surface disposal), pollutants can spread underground and pollute sources of drinking water.

Movement of ground water is usually quite slow, on average, ranging from about one foot per day to perhaps ½ inch per month. Therefore, in some areas, it might take days for water to travel from the point where it enters the ground, to a point of discharge into a stream, or it might take millennia (Heath, 1983). Movement of water through an aquifer of unconsolidated sediment depends in part on the pore size between particles as well as the size and uniformity of the actual particles. Water is stored in and moves through connected pore spaces and larger particles of uniform size can more readily transmit water. Because hydraulic conductivity can be high, surficial (water table) aquifer systems of unconsolidated sediments may be susceptible to contamination (USGS, 2016).

An understanding of the water cycle emphasizes the connections between surface and ground water. While the Borough of West Cape May relies primarily on water from public wells (not individual wells) fed by ground water, the water is no less part of the natural water cycle, and is susceptible to human impacts and the influence of climate and geology.

B. The Aquifers in West Cape May

West Cape May’s drinking water comes predominantly from ground water (as does almost half of New Jersey’s drinking water), and originates as local or regional precipitation. The ground water moves through the aquifers, eventually discharging into streams or directly into the aquifer systems and is distributed by wells (USEPA, August 1992).

The density of housing and impervious surfaces can impact aquifers and may result in reduced recharge, lowered yields, increased interference (wells interfering with each other), and degradation of ground water quality. In any aquifer, if the rate of water use exceeds the recharge rate, well yields will decrease (see **Section 5D**). Furthermore, these changes can alter stream flow dynamics resulting in higher flows after storm events and lowered flows between events. In coastal areas, increased rates of water use may also result in saltwater intrusion into freshwater aquifers and wells.

Aquifers are typically described as being unconfined or confined. *Unconfined* aquifers are those aquifers where the ground water is directly connected to the atmosphere through the pores of the aquifer. *Confined* aquifers are water-bearing formations that are separated from the surface by a layer of rock or soil through which water cannot move (Dunne and Leopold, 1978).

The Borough of West Cape May is underlain by five main aquifers, described in the paragraphs below and summarized in **Table 5.1**. **Figure 5b** illustrates the unconfined (water table) aquifer (the Holly Beach water-bearing zone) that outcrops as the upper aquifer throughout most of Cape May County. **Figure 5c** presents a cross-sectional view of the unconfined and confined aquifers that occur beneath the county. These include Holly Beach water-bearing zone, estuarine sands, the Cohansey, Rio Grande water-bearing zone, and the Atlantic City 800-foot sand.

The recharge area (where precipitation or snowmelt enters the ground) for unconfined (surficial) aquifers is local, usually above the aquifer. In contrast, the recharge area for confined aquifers can be a distance away, wherever that aquifer eventually intersects with the surface of the ground. In the case of the Atlantic City 800-foot sand aquifer, the outcrop area is well to the west of Cape May County where the confining layer thins and the 800-foot sand joins the Kirkwood –Cohansey aquifer system (McAuley et al., 2001). Confined aquifers can also receive additional recharge from “leaking” from overlying aquifers.

West Cape May relies on groundwater from an aquifer system that is interconnected among all Atlantic coast counties such that withdrawals in lower Cape May County may affect water levels and water quality in Atlantic or Ocean County municipalities, and vice versa. Due to the interconnected nature of the coastal plain groundwater supply, the New Jersey Department of Environmental Protection (NJDEP) has recently completed a water supply plan for the

confined aquifers of the coastal plain that explores options for ensuring adequate freshwater supply as demand continues to increase in the coming years (NJDEP, 2017).

The following are the main aquifers of lower Cape May County, as described by French and Silvestri, 1999; Herman et al., 1998; Lacombe and Carleton, 2002; and Sugarman et al., 2016.

Holly Beach Water-Bearing Zone

The Holly Beach water-bearing zone is a water table (surficial) aquifer that is recharged directly from local precipitation. It is composed of orange and yellow gravels and sands (coarse to fine-grained) with minor silt and clay deposits. It includes alluvium, beach, dune, deltaic, and marine sands. Although present at the ground surface in wetland areas, it can be found more than 10 feet deep in upland regions. The aquifer thickness of the freshwater portion ranges from 10-80 feet in the southern part of the county. There are high iron and manganese concentrations locally, with elevated salinity near coastal areas. Well yields range from 3 – 300 gallons/minute (Gill, 1962 as cited in Lacombe and Carleton, 2002).

Estuarine Sand Aquifer

The estuarine sand aquifer is composed of poorly sorted gravel and medium- to fine-grained sand, with some lignite, and ranges from 25 – 160 feet thick. Well yields range from 3 – 230 gallons/minute, although one test well generated 700 gallons/minute (Sugarman et al., 2016). High salinity levels are found in Cape May City and Villas, some areas of Middle Township, as well as along barrier islands.

Cohansey Aquifer




The Cohansey aquifer is the deepest of the peninsula's shallow water aquifers and is confined in Cape May County. It is made up of fine- to medium-grained sand, with lenses of silt and clay. The aquifer is 60 – 180 feet thick, with well yields ranging from 5 – 700 gallons/minute, though can go as high as 1,500 gallons/minute. It is predominantly fresh water in the mainland sections of the county, but has elevated chloride levels near the coast and barrier island.

Rio Grande water-bearing zone

The Rio Grande water-bearing zone underlays the Cohansey aquifer and is comprised of fine- to medium-grained sand. It is typically 50 – 100 feet thick, being thicker in the southern part of the county. High chloride levels are found south of Middle Township and east, beneath barrier islands.

Atlantic City 800-foot sand

This aquifer is the deepest of the Cape May aquifers and consists of fine- to medium-grained sand, with silt and clay. There are high levels of chloride in the lower part of the Cape May peninsula and southern Lower Township. The aquifer averages 150 – 200 feet thick and is found 400 – 950 feet below the surface. It yields large quantities of water from Stone Harbor to Ocean City and on mainland from Middle to Upper Township, typically 500 – 1000 gallons/minute.



Prepared by
Kratzer
Environmental
Services

Data Sources: NJDEP



Note: Map accuracy is limited to the accuracy and scale of the original data.

Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Figure 5b: West Cape May Aquifer & Sole-source Aquifer

The water table (unconfined) aquifer for much of southern Cape May County, including West Cape May Borough, is the Holly Beach water-bearing zone. The location of ground water wells, discharge sites and monitoring stations serving West Cape May are also pinpointed on this map. See Figure 5c for an illustration of the aquifers in cross section.

28 Figure 5c: Hydrostratigraphy in Cape May County
Relative positions of the five aquifers in lower Cape May County. The top image shows the location of the cross section of hydrostratigraphy pictured in the bottom figure.



Source:
 Lacombe, Pierre J. and Glen B. Carleton. 2002. Hydrogeologic Framework, Availability of Water Supplies, and Saltwater Intrusion, Cape May County, New Jersey. U.S. Geological Survey Water-Resources Investigations Report 01-4246. Prepared in Cooperation with the NJDEP.

Table 5.1: Characteristics of West Cape May Aquifers

Aquifer Unit	Aquifer Rank [■]	Characteristics	Common Range*		
			Depth (ft)	Yield (gpm)	Thickness
UNCONSOLIDATED AQUIFERS					
Holly Beach water-bearing zone	C	Surficial aquifer of Quaternary age, orange and yellow sand and gravel, with thin clay and silt lenses and shell layers. It includes alluvium, beach, dune, deltaic, and marine sands. 20'-100 feet thick. Locally may be contaminated by human activities, and may be locally high in nitrate and iron with elevated salinity near coastal areas.	2 wells are <100' ^[S]	250-350 ^[S]	20-100 ^[S]
				Avg. = 110 Median = 100 ^[F]	
				3-300 ^[Gill 1962, as cited in L]	10 – 80 ^[L]
CONFINING UNIT					
Estuarine Sand	C	Fine to medium sands with some gravel and lignite. Variable thickness 20 – 150+ feet. High chlorides and sodium levels in Cape May City and barrier islands and parts of Lower Township. High iron in most areas.	80-225 ^[S]	700 from one well ^[S]	3-100 ^[S]
				Avg. = 360 Median = 250 ^[F]	
				3-230 ^[L]	25 -165 ^[L]
CONFINING UNIT					
Cohansey	A	Composed of medium- to coarse-grained sand with lenses of silty clay. High chloride levels beneath barrier islands and Cape May City. It can be thin, may be connections between confining units. 60-180 feet thick. Yields large quantities of water on the mainland.	200-300 in southern part of county ^[S]	Up to 1500 ^[S]	50-200 ^[S]
				Avg. = 514 Median = 500 ^[F]	
				5-700 ^[L]	60-180 ^[L]
CONFINING UNIT					
Rio Grande water-bearing zone	B	Fine to medium-grained sand. Typically 50-100 feet thick, being thicker in the southern part of the county. High chloride levels south of Middle Township. Upper layer within the Kirkwood Formation.		One well 1000 ^[S]	50-100 ^[S]
				Avg. = 460 Median = 300 ^[F]	
				1100 ^[L, one well]	30-170 ^[L]
CONFINING UNIT					
Atlantic City 800-foot sand	A	Fine to medium-grained sand, with silt and clay. High levels of chloride in the lower part of the Cape May peninsula and southern Lower Township. Averages 150 – 200' thick. Yields large quantities of water from Stone Harbor to Ocean City and on mainland from Middle to Upper Township.	400-950 ^[S]	500-1000 ^[S]	150-200 ^[S]
				Avg. = 579 Median = 700 ^[F]	
				600-900, or even > 1000 ^[L]	125-150 ^[L]
<p>■ Aquifer Rank is from NJGS GIS data. It is based on High Capacity Wells (such as water-supply, irrigation, and industrial-supply wells sited and tested for maximum yield. Many of the wells have boreholes exceeding the standard six-inch diameter for domestic wells. State Rank is best viewed on a relative basis, with “A” yielding the most water, and “E” the least. Median High Capacity Wells Yield (in gpm): [A] > 500; [B] 251 to 500; [C] 101 to 250; [D] 25 to 100; [E] <25</p> <p>Sources: Herman et al., 1998; USGS, January 14, 2013; except as noted below.</p> <p>* Various studies list differing results for the range of depth, yield and thickness of the units, and the sources are noted in brackets. Note that these characteristics vary from location to location.</p> <p>^[F] French and Silvestri, 1999</p> <p>^[L] Lacombe and Carleton, 2002</p> <p>^[S] Sugarman et al., 2016</p>					

C. Sole-Source Aquifer

The Safe Drinking Water Act (SDWA) of 1974 contains a provision in Section 1424(e) that provides for designating an aquifer that is the sole or principal drinking water source for an area and that, if contaminated, would create significant hazard to public health. As defined by the U.S. Environmental Protection Agency (EPA), *sole-source aquifers* (SSA) are those aquifers that contribute more than 50% of the drinking water to a specific area and the water would be impossible to replace if the aquifer were contaminated. Once designated, no Federal financial assistance may be approved for any project that may contaminate the aquifer through a recharge zone so as to create a significant hazard to public health (US EPA, August 1992). Therefore, the EPA must review any federally-funded project in an area that could affect ground water in a sole-source aquifer, including the *aquifer's recharge zone* (the area through which water recharges the aquifer) and its *stream-flow source zone* (the upstream area that contributes recharge water to the aquifer).

The Kirkwood-Cohansey Aquifer System met the technical requirements for SSA designation, and Notice of approval was published in the Federal Register 57 FR 39201, August 28, 1992. It covers much of the Inner and Outer coastal plain. The Kirkwood-Cohansey SSA in New Jersey is shown in **Figure 5b** (upper right inset) and includes water-bearing units that supply West Cape May's water (NJDEP, May 19, 1998).

D. Recharge

Ground water recharge is defined as water added to an aquifer (for example, precipitation that seeps into the ground deep enough to enter the saturated zone of the fractured bedrock). A *ground water recharge area* is the land area that allows precipitation to seep into the saturated zone. These areas are generally at topographically high areas with discharge areas at lower elevations, commonly at streams or other water bodies (i.e. the ground water returns to surface water). In general, ground water divides¹⁶ coincide with, or are slightly offset from surface water divides (Lewis-Brown and Jacobsen, 1995) (watersheds are described in **Section 6A** and shown in **Figure 6a**). Most ground water flows through the shallow layers of soil and weathered bedrock to the nearest stream. A smaller percentage penetrates deeper and recharges the aquifer.

Recharge rates are expressed in terms of the amount of precipitation that reaches the aquifer per unit of time (e.g. inches/year during a drought year is used in Figure 5d). New Jersey receives an average of about 40 to 51 inches of precipitation per year (lowest along the southeast coast, including West Cape May, and highest in the north-central parts of the state) (ONJSC, no date). Many factors affect the amount of recharge that will occur in a given area, including climate (e.g. the amount, intensity, and form of precipitation, and the effect of wind, humidity and air temperature on evapotranspiration), soil, surficial geology, and vegetation factors. In addition, recharge of ground water varies seasonally. During the growing season, precipitation is intercepted by plants and returned to the atmosphere through transpiration (part of the hydrologic cycle, see Section 5A). Likewise, evaporation is higher during the warmer months. Together, these are known as evapotranspiration. Therefore, most recharge occurs during late fall, winter, and early spring, when plants are dormant and evaporation rates are minimal (Heath, 1983). Relative to land use, recharge rates in forests are much higher than those in urban areas (Heath, 1983). This is because urban areas have large areas covered with impermeable surfaces, hastening runoff to surface water, instead of allowing precipitation to percolate into the ground.

¹⁶ A ground water divide is a line on a water table where on either side of which the water table slopes downward. It is analogous to a drainage divide between two drainage basins on a land surface.

http://www.srh.noaa.gov/jetstream/append/glossary_g.htm

To ensure that water is available during all weather conditions for human consumption as well as ecosystems dependent on water, the NJDEP established the Planning Threshold, or *dependable yield*, to be used for planning purposes. *Dependable yield* is defined as “the water yield maintainable by a ground-water system during projected future conditions, including both a repetition of the most severe drought of record and long-term withdrawal rates without creating undesirable effects.” The most severe drought on record was in the early 1960's (see **Section 2A**), and this is used in the Statewide Water Supply Plan. However, the Plan acknowledges that there is insufficient long-term precipitation data to prove that this is the worst drought that could occur in the future, in duration or severity, and recommends re-evaluation of safe-yield estimates and development of optimal strategies for severe droughts (NJDEP OEP, 1996). Robert Canace, formerly of the NJ Geological Survey, suggested that 20% of the estimated recharge should be used for planning purposes, representing the portion of recharge actually available for use during drought conditions (Canace, 1995).


In view of the importance of not exceeding the aquifers' safe yield, the New Jersey Geological Survey has completed studies quantifying recharge, as discussed in the following sections.

New Jersey Geological Survey Recharge Method GSR-32


N.J.S.A. 58:11A, 12-16 required the NJDEP to publish a methodology to map and rank aquifer-recharge areas. In addition, the legislation required the development of ground water protection practices designed to encourage ecologically sound development in aquifer-recharge areas (Charles et. al., 1993). To fulfill the requirements of this legislation, the NJ Geological Survey developed GSR-32, which estimates ground water recharge (but not aquifer recharge), and is useful for evaluating the relative effect of present and future land uses on recharge areas (Charles et. al., 1993). For this method, recharge was calculated based on data for precipitation, soil, land-use/land-cover¹⁷, surface runoff, and evapotranspiration. This method was then applied by NJGS to create a GIS coverage (see Figure 5d). There were a number of assumptions made for the calculations and model inputs that limit the accuracy of the method: 1) the calculated ground water recharge includes any water entering the ground (in actuality, lesser amounts actually enter the aquifer); 2) assumes that all water that migrates below the root zone recharges the aquifer (which does not happen); 3) addresses only natural ground water recharge, and does not include artificial recharge, withdrawals or natural discharge; 4) wetlands and water bodies were eliminated from analysis, because the direction of flow between ground water and surface water is site-specific and also varies seasonally, and this level of detail was beyond the scope of the study (these areas were assumed to provide no recharge or discharge); 5) stream baseflows used may not be representative of local streams (Charles et. al., 1993) and 6) does not consider topography, depth to bedrock, presence of impervious surfaces, and/or type of bedrock underlying soils. An additional limitation of the data is that they estimate long-term average annual recharge, which does not represent the reduced recharge during critical summertime conditions (NJ Water Supply Authority, 2002).

¹⁷ Land use/land cover data from 1995-1997 were used for this study. Changes in land use/land cover and impervious surfaces affect recharge, but are not shown on **Figure 5d**, because this involves complex calculations, and NJGS has not updated this GIS data layer.

Drought Recharge



Aquifer Rank/Ground Water Recharge Rank



Legend


- West Cape May
- Municipalities
- Roads
- Streams
- Water

Estimated Drought Recharge Value (in/yr)	
	0
	5
	6
	7
	8
	9

Aquifer Rank	High Yield Wells	Ground water recharge rank	Recharge
A	> 500 gpm	A	18 to 23 in/yr
B	250 to 500 gpm	B	10 to 17 in/yr
C	100 to 250 gpm	C	8 to 9 in/yr
D	25 to 100 gpm	D	1 to 7 in/yr
E	< 25 gpm	E	0 in/yr

L: hydric soils--no recharge calculated

W: wetlands and open water--no recharge calculated



Data Sources: NJDEP, NJDOT

Note: Map accuracy is limited to the accuracy and scale of the original data sets.

Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Prepared by Kratzer Environmental Services

Figure 5d: West Cape May Recharge and Aquifer Potential

Left: Drought recharge values are estimated in inches per year of subsurface recharge. Right: Recharge ranks range from A (highest) to E (lowest). The two ranks are combined in the format "ground water recharge rank/aquifer recharge rank". Higher ranks are positively correlated with higher yields.

Applying the GSR-32 method to West Cape May, the estimated average annual subsurface recharge rates range from 0 to 10 inches per year (excluding surface water, wetlands and hydric soils) and 0 to 9 inches per year during drought (shown on **Figure 5d**). Applying the 20% consumptive use limit to these figures results in usable recharge from 0 to 2 inches per year. As previously mentioned, only a portion of water entering the ground actually recharges the aquifer, but since GSR-32 did not attempt to quantify this amount, this method would be better described as *soil recharge*.

New Jersey Geological Survey Ground Water Potential

In 2005, also in response to N.J.S.A. 58:11A, 12-16, the NJ Geological Survey developed a qualitative representation of the potential for aquifer recharge. This was created by combining ground water recharge rankings and aquifer rankings.

NJGS assigned a relative rank based on the inches of ground-water recharge per year (mapped in **Figure 5d**), from A (highest recharge) to E (lowest recharge). Relative values of aquifer yield (based on high yield industrial wells and described in **Table 5.1**) were assigned to each aquifer, from A (highest yield) to E (lowest yield). For both data sets, areas of wetlands, open water and hydric soils were not ranked, since individual areas differ in whether they increase or decrease recharge, which varies seasonally. These two ranks are combined in the format "ground water recharge rank/aquifer recharge rank" and illustrated in **Figure 5d**. For example, A/A would be an area with the highest relative recharge and highest yield, and an area designated E/E would have the lowest recharge and lowest yield, while other combinations would lie somewhere in between (French, 2004). The State Ranks for the aquifers underlying West Cape May are C/D, D/D and E/D.

E. Ground Water Quality

Pollution, such as nitrates, bacteria, metals, pesticides and antibiotics, can enter ground water via non-point sources (including septic systems and runoff from fields and roads), point sources, and rain. The New Jersey Comparative Risk Project (2003) identified a number of possible human health risks from drinking water, including lead (which, when present, is usually from the plumbing) (NJDEP, August 17, 2017), radon, arsenic, MTBE (methyl-t-butyl ether, which is a fuel oxygenate), nitrates, and waterborne pathogens.

In addition, radioactive substances (including uranium, thorium, radium, and radon) from natural sources (see **Section 3D**) are frequently found in ground water in New Jersey. Almost all rocks and soil contain at least some radioactive substances, which can sometimes exceed safe levels in drinking water (no data was found specific to West Cape May, but background information may be found in **Internet Resources**).

Saltwater Intrusion

Saltwater intrusion is a major concern in coastal regions and is the predominant threat to the Cape May County water supply (Lacombe and Carleton, 2002). It occurs when excessive drawdown of the water table through freshwater withdrawals allows saltwater to enter supply wells. The rates of aquifer recharge and discharge determine the extent of saltwater intrusion (Dunne and Leopold, 1978).

Possible actions to halt or mitigate saltwater intrusion include; limiting water withdrawals, protecting aquifer recharge areas locally and regionally, enhancing groundwater recharge with the use of constructed recharge basins or artificially by injecting storm water or treated sewage effluent into aquifer(s), Aquifer Storage and Recovery (ASR), constructing saltwater desalinization plants, or any combination of the above. Each of these methods has costs

and benefits, with a goal of not exceeding the carrying capacity of the groundwater supply. See Lacombe and Carleton, 2002; and Lacombe et al., 2009 for a discussion of sustainable potable water-supply alternatives to meet current and future water supply demands in Cape May County.

Public Water Wells

The Cape May City Water Department serves the City and surrounding municipalities, including West Cape May, from the wells shown in **Figure 5b**. The two main wells in use today are located in the Atlantic City 800-foot sand aquifer. Due to their higher salt content, water from these wells is treated by the City's reverse osmosis water treatment facility constructed in 1998. In addition, some water from Well No. 5, drawn from the 560-foot-deep Cohansey aquifer, is blended with the clean water from the desalination plant. As saltwater intrusion continues to threaten the Cohansey wells in this region, other options are under consideration, such as recharging existing wells (e.g., No. 4) with clean water so that it could be used at future times (i.e. Aquifer Storage and Recovery (ASR)) (Degener, August 24, 2011) (see also Lacombe and Carleton, 2002; and Lacombe et al., 2009).

In addition to the reverse osmosis process, lime is used for pH adjustment and sodium hypo-chlorite is used for disinfection (West Cape May Water Department, 2016).

In the 2016 annual report (based on 2015 and some earlier water quality data), the West Cape May Water Department reported one positive Total Coliform Bacteria sample and follow-up sampling was negative. Coliform bacteria are naturally present in the environment and are monitored as an indicator that other, potentially-harmful, bacteria could be present. One water sample out of 10 in 2014 exceeded the lead action level. All other measured water quality parameters were below the Maximum Contaminant Levels (MCLs) for drinking water (West Cape May Water Department, 2016).

In 2016 to 2017, West Cape May Water Department has had 4 violations of type 27 (Monitoring and Reporting) for chlorine and 2 for trihalomethanes (TTHM) and one for type 71, violation of the Consumer Confidence Rule for submitting results late (NJ Drinking Water Watch, 2017). Water quality testing results are available online (see **Internet Resources**).

Private Wells

The New Jersey Private Well Testing Act (N.J.S.A. 58:12A-26 et seq.) became effective in September 2002, mandating private well testing upon the sale of a house. The number of wells tested in a municipality reflects the number of real estate transactions involving homes with private wells. Private wells in Cape May County are required to be tested for total coliform (plus fecal coliform or E.coli if total coliform is positive), nitrate, iron, manganese, pH, VOCs, lead, mercury, and gross alpha particle activity. Since West Cape May is served primarily by public water systems, just 28 private wells have been tested pursuant to this regulation from September 2002 to April 2014. Of these 28 wells tested, one well exceeded the standard for bacteria; 18 exceeded the manganese standard; 17 exceeded the pH standards; and 13 exceeded the iron MCL,(see **Table 5.2**) (NJDEP Division of Water Supply and Geoscience, August 17, 2017; NJDEP Division of Water Supply and Geoscience, January 19, 2017).

Table 5.2: NJ Private Well Testing Act Data Summary (September 2002 to April 2014) in West Cape May Borough

Parameter	Number of Wells Sampled	% of Wells Exceeding MCL	MCL
Nitrate	28	0.0%	10 mg/l
Arsenic	Testing is not required under PWTA in Cape May Co.		5 µg/l
Iron	28	46.4%	0.3 mg/l
Manganese	28	64.3%	0.05 mg/l
Gross Alpha	26	0.0%	(initial) ¹⁸ 5 pCi/L
Mercury	28	0.0%	2 µg/l
VOC	28	0.0%	*
Fecal coliform or E. coli	28	3.6%	0 colonies
pH	28	60.7%	6.5-8.5
* MCLs vary for the 26 Volatile Organic Compounds (VOCs) required by the PWTA.			
Source: NJDEP Division of Water Supply and Geoscience, January 19, 2017; Atherholt et. al., 2008			

F. Ground Water Quality Standards

The New Jersey Ground Water Quality Standards (GWQS; N.J.A.C. 7:9C) (last amended July 22, 2010) specify the quality criteria and designated uses for ground water, and serve as the basis for setting ground water discharge standards under the New Jersey Pollutant Discharge Elimination System program (see **Section 5H**), as well as for establishing standards for ground water cleanups and other relevant laws. The criteria are numerical values assigned to each constituent (pollutant). The GWQS also contain technical and general policies to ensure that the designated uses can be adequately protected.

Ground water within watersheds of FW1 surface waters (see **section 6D** for surface water classifications), state-owned Natural Areas, and the major aquifers of the Pinelands Area are designated *Class I*. The designated use for Class I ground water is the maintenance of special ecological resources, with secondary uses being potable, agricultural and industrial water. *Class II* waters are those not specifically designated Class I or Class III. The designated use of Class II ground waters is to provide potable water using conventional treatment. Class II criteria specify the levels of constituents above which the water would pose an unacceptable risk for drinking water. *Class III* ground waters can be used for anything other than for potable water (NJDEP Bureau of Environmental Analysis, Restoration and Standards, September 15, 2017).

West Cape May's waters are designated Class II (to provide potable water with conventional treatment). It should not be assumed that ground water quality everywhere meets the criteria for each classification area in view of natural variability and the possibility of localized pollution.

G. Ground Water Discharges

New Jersey regulates the discharge of pollutants to ground water under the authority of the New Jersey Water Pollution Control Act (WPCA) N.J.S.A. 58:10A. The New Jersey Pollutant Discharge Elimination System (NJPDDES) permit program regulations are contained in N.J.A.C. 7:14A (NJDEP, January 5, 2009).

NJPDDES permits are required for discharges to ground water of both sanitary and industrial wastes. These permits, which limit the mass and/or concentration of pollutants

¹⁸ Results greater than 5 pCi/L requires a second gross alpha count. The MCL for gross alpha is 15 pCi/L.

discharged, are issued to sanitary and industrial facilities that have ongoing, operational discharges of wastewater to ground water. The purpose is to restrict the discharge of pollutants to the ground waters of the state and protect the public health and the environment. Discharges from past activities may continue to be regulated under the Site Remediation Program or the Division of Solid and Hazardous waste.

There are 2 ground water discharges within West Cape May, described below in **Table 5.3** and shown on **Figure 5.b** (NJDEP DWQ BNPC, July 18, 2007).

Table 5.3: NJPDESGW in West Cape May

FACILITY	PI #	NJPDES #	DISCHARGE TYPE*
DEPOT TRAVEL PARK INC	49849	NJG0135402	T1
WESTWOOD MOBILE ESTATE/REEVES	47400	NJG0089265	T1
* T1 stands for Sanitary Subsurface Disposal and authorizes the discharge of sanitary sewage from facilities to a subsurface disposal (septic) system with a design volume in excess of 2,000 GPD			
Source: NJDEP, Division of Water Quality (DWQ), Bureau of Nonpoint Pollution Control (BNPC), July 18, 2007			

H. Contaminated Sites

On May 7, 2012, NJDEP adopted amendments, repeals, and new rules to implement site remediations through the *Site Remediation Reform Act (SRRA)*, N.J.S.A. 58:10C-1 et seq., and related amendments to the *Brownfield and Contaminated Sites Act (Brownfield Act)* N.J.S.A. 58:10B-1 et seq., the *Spill Compensation and Control Act (Spill Act)*, N.J.S.A. 58:23-11 35 seq., the *Industrial Site Recovery Act (ISRA)*, N.J.S.A. 13:1K-6 et seq., and the *Underground Storage of Hazardous Substances Act (UST Act)*, N.J.S.A. 58:10A-21 et seq. This major shift requires remediations of contaminated sites to proceed under the supervision of a *Licensed Site Remediation Professional (LSRP)* (hired by the property owner) instead of NJDEP (NJDEP Site Remediation Program, May 7, 2012).

The goal of these changes is to increase the pace of remediation, in order to decrease the threat of contamination to public health and safety and the environment, and to more quickly return properties to productive use that are underutilized due to contamination.

Some key provisions create a licensing board and a code of ethics (including penalties for violations) for LSRPs; establish obligations of each person responsible for conducting remediation; institute mandatory timeframes for the completion of key phases of site remediation; set forth the circumstances under which NJDEP would undertake direct oversight of a remediation; and require NJDEP to establish presumptive remedies for residential development, schools and childcare facilities to ensure that the remediation at these sites is protective of human health and safety and of the environment (NJDEP SRP, September 15, 2017).

The LSRP program does not apply to unregulated underground storage tanks (i.e. residential tanks) (see **Internet Resources**).

Known Contaminated Sites List (KCSL)

The NJDEP Site Remediation Program compiles a list of Known Contaminated Sites (KCS). The Known Contaminated Sites List 19 (non-homeowner) for New Jersey (as required

¹⁹ The GIS data is updated periodically (the most recent data currently available is from April 27, 2017. The tabular data is updated more frequently (see **Internet Resources**).

under N.J.S.A. 58:10-23.16-17 and also the New Residential Construction Off-Site Conditions Disclosure Act N.J.S.A 46:3C1 et seq.) contains sites defined as those sites and properties within the state where contamination of soil or ground water has been confirmed at levels equal to or greater than applicable standards. Sites identified in the Known Contaminated Sites list can undergo a variety of activities, ranging from relatively simple soil removals to highly complex remedial activities. It is important to note that the list may include sites where remediation is either currently under way, required but not yet initiated or has been completed (and no longer considered contaminated). In addition, new contaminated sites may have been identified since the creation of this list and are not included here (NJDEP SRP, February 14, 2017).

There are no active, pending, or closed contaminated sites within the Borough of West Cape May. Nor are there sites in West Cape May currently on the National Priorities (Superfund) List.

Homeowner sites are not included because they generally involve small heating oil discharges from leaking underground storage tanks (USTs) that are resolved relatively quickly (see Internet Resources for a link to NJDEP's grant program for removal and cleanup of USTs) (NJDEP SRP, April 25, 2017).

Classification Exception Area (CEA)

The Ground Water *Classification Exception Area* (CEA) dataset identifies those sites where ground water contamination has been identified and the NJDEP has established a Classification Exception Area (CEA). CEAs are institutional controls in geographically defined areas within which the New Jersey Ground Water Quality Standards (NJGWQS) for specific contaminants have been exceeded. When a CEA is designated for an area, the constituent standards and designated aquifer uses are suspended for the term of the CEA. This data is intended to provide information to the public regarding areas of contaminated ground water to prevent inappropriate well placement, preventing potential health risks and can minimize unintended contaminant plume migration (NJDEP SRP, March 29, 2016). There are no CEAs in West Cape May.

Deed Notice

A *Deed Notice* is defined by NJSA 58:10B-13a as a "...notice to inform prospective holders of an interest in the property that contamination exists on the property at a level that may statutorily restrict certain uses of, or access to, all or part of that property...." The purpose of the deed notice GIS layer is to minimize any chance of exposure to contaminants remaining on the property (NJDEP. August 28, 2017). There are no Deed Notices delineated within the Borough of West Cape May.

Remediated Sites

A current Data Miner search revealed that there are currently no sites undergoing remediation in West Cape May (NJDEP SRP, June 13, 2017).


I. Ground Water Level Monitoring

The *ground water level* is the distance from the land surface to the water in a well. Ground water level monitoring is critical for determining the current state of the ground water, identifying trends and predicting ground water drought. In addition to drought, over-withdrawal of ground water can occur in areas where more ground water is being pumped out of the aquifer than is replenished through recharge. This could lead to a drop in the ground water level, affecting well performance, sometimes causing wells to go dry, in addition to causing a decrease


in the baseflows of adjacent streams. In coastal areas like West Cape May, over-withdrawal of ground water also leads to saltwater intrusion into freshwater wells, a major concern.

The USGS maintains a nation-wide network of wells to monitor the effects of droughts and other climate variability on ground water levels. Data from these wells (among other factors) are used by the NJDEP to determine drought status (see **Internet Resources**). In West Cape May, the USGS has monitored a 293 foot deep well since 1957. A description of this site and a graph of ground water level for 2016-2017 is shown in **Table 5.4**. Its location is shown in **Figure 5b**. On average, the lowest groundwater levels occur during June-September, while highest levels occur January-March (USGS, 2017a; USGS 2017b).

Table 5.4: USGS Real-Time Ground Water Level Network

Site Number: 385607074555201 - 090150-- West Cape May 1 Obs	
Location:	Latitude 38°56'07", Longitude -74°55'56" NAD27 West Cape May Borough, Cape May County, New Jersey , Hydrologic Unit 02040302
Well Characteristics:	Depth: Well depth: 293 feet Land surface altitude: 6.60 feet above NGVD29.
Aquifer:	Well completed in "Northern Atlantic Coastal Plain aquifer system" national aquifer Well completed in "Cohansey Sand" local aquifer
Begin date (periodic record):	07/27/1957
Extremes for Period of Record:	<i>Highest Water Level:</i> 17.69' <i>Median:</i> 19.58' <i>Lowest Water Level:</i> 21.50'
Website:	https://waterdata.usgs.gov/nwis/inventory/?site_no=385607074555201&agency_cd=USGS https://groundwaterwatch.usgs.gov/AWLSites.asp?S=385607074555201
 <p>USGS 385607074555201 090150-- West Cape May 1 Obs</p> 	
Site Number: USGS 390156074533401 090333-- Pump Pond N Obs	
Location:	Latitude 39°01'56", Longitude 74°53'34" NAD27 Middle Township, Cape May County, New Jersey , Hydrologic Unit 02040206
Well Characteristics:	Well depth: 43 feet Hole Depth: 69 feet Land surface altitude: 20 feet above NGVD29.
	Well completed in "Northern Atlantic Coastal Plain aquifer system" (S100NATLCP) national aquifer. Well completed in "Holly Beach Water-Bearing Zone" (112HLBC) local aquifer Cape May County, New Jersey , Hydrologic Unit 02040302
Begin date (periodic record):	07/15/1992

Extremes for Period of Record: *Highest Water Level: 10.80' Median: 13.02' Lowest Water Level: 13.83'*
Website: https://waterdata.usgs.gov/nwis/inventory?agency_code=USGS&site_no=390156074533401
<https://groundwaterwatch.usgs.gov/AWLSites.asp?S=390156074533401>



Note: These sites are maintained by USGS New Jersey Water Science Center.
Sources: USGS, 2017a & b

References: Ground Water

A. Water Cycle/Ground Water Background

Heath, Ralph C. 1983. Basic Ground-Water Hydrology. U.S. Geological Survey Water-Supply Paper 2220. U.S. Geological Survey in cooperation with the North Carolina Department of Natural Resources and Community Development, 84 pages.

Miller, G. Tyler. 1988. Living in the Environment: An Introduction to Environmental Science. Wadsworth Publishing Company: Belmont, California, 603 pages.

US Geological Survey. 2016. Aquifer basics: Unconsolidated and semiconsolidated sand and gravel aquifers. <https://water.usgs.gov/ogw/aquiferbasics/uncon.html>. Accessed October 25, 2017.

US Geological Survey. No Date. Water Science for Schools: The Water Cycle. <http://ga.water.usgs.gov/edu/watercycle.html>

B. Aquifers

Dunne, Thomas and Luna B. Leopold. 1978. Water in Environmental Planning. W. H. Freeman and Company: New York, New York. 818 pages.

French, Mark and ShayMaria Silvestri. 1999. Groundwater Recharge and Aquifer Recharge Potential for Cape May County, New Jersey. N.J. Department of Environmental Protection and N.J. Geological Survey. Open-File Map OFM-31. <http://www.nj.gov/dep/njgs/pricelst/ofmap/ofm31.pdf>

Gill, H.E. 1962. Groundwater Resources of Cape May County, New Jersey, Salt-Water Invasion of Principal Aquifers. N.J. Department of Conservation and Economic Development, Division of Water Policy and Supply, Water Resources Special Report 18, 171 pages.

Gill, H.E., 1959, Geology and ground-water resources of the Cape May Peninsula, lower Cape May County, New Jersey: New Jersey Department of Conservation and Economic Development Water-Resources Circular 1, 19 p.

Herman, Gregory C., Robert J. Canace, Scott D. Sanford, Ronald S. Pristas, Peter J. Sugarman, Mark A. French, Jeffrey L. Hoffman, Michael S. Serfes, and William J. Mennel. 1998. Aquifers of New Jersey. Open-File Map OFM 24. <http://www.state.nj.us/dep/njgs/pricelst/ofmap/ofm24.pdf>. NJGS Digital Geodata Series DGS98-5. GIS data. <http://www.state.nj.us/dep/njgs/geodata/dgs98-5.htm>

Lacombe, Pierre J. and Glen B. Carleton. 2002. Hydrogeologic Framework, Availability of Water Supplies, and Saltwater Intrusion, Cape May County, New Jersey. U.S. Geological Survey Water-Resources Investigations Report 01-4246. Prepared In Cooperation With The New Jersey Department of Environmental Protection. 151 pages. <https://pubs.usgs.gov/wri/wri014246/pdf/wrir01-4246.pdf>

McAuley, Steven D., Julia L. Barringer, Gary N. Paulachok, Jeffrey S. Clark, and Otto S. Zapecza. 2001. Ground-water flow and quality in the Atlantic City 800-Foot Sand, New Jersey. New Jersey Geological Survey, Geological Survey Report GSR 41, West Trenton, New Jersey. <https://pubs.usgs.gov/unnumbered/70114185/report.pdf>

New Jersey Department of Environmental Protection (NJDEP). 2017. Appendix B- Water supply options, confined aquifers of the New Jersey Coastal Plain in New Jersey Water Supply Plan 2017-2022. <http://www.nj.gov/dep/watersupply/wsp.html>

Sugarman, Peter J., Donald H. Monteverde, Scott D. Stanford, Stephen W. Johnson, Yelena Stroitleva, Ronald S. Pristas, Kathleen Vandegrift, and Steven E. Domber. 2016. Geologic and Aquifer Map of Cape May County, New Jersey. N.J. Geological and Water Survey, NJ Department of Environmental Protection. GMS 16-1. <http://www.state.nj.us/dep/njgs/pricelst/gmseries/gms16-1.pdf>

U.S. Environmental Protection Agency (US EPA) Region 2. August 1992. Ramapo Aquifer Systems. Support Document. <http://www.epa.gov/region02/water/aquifer/ramapo/ramapo.htm>

U.S. Geological Survey (USGS). January 14, 2013. Aquifer and Well Characteristics in NJ. https://nj.usgs.gov/infodata/aquifers/table_1.html

C. Sole Source Aquifers

NJDEP, New Jersey Geological Survey (NJGS), Jeffrey L. Hoffman (ed.). May 19, 1998. NJDEP Sole-Source Aquifers in New Jersey. <http://www.state.nj.us/dep/njgs/geodata/dgs98-6.htm>

D. Recharge

Canace, Robert. 1995. New Jersey Geological Survey, Department of Environmental Protection, Division of Science and Research. Critique of the Franklin Township ground water resources.

Charles, Emmanuel G., Cyrus Behroozi, Jack Schooley, and Jeffrey L. Hoffman. 1993. A Method for Evaluating Ground-Water-Recharge Areas in New Jersey. NJ Geological Survey Report GSR-32. NJDEP Geological Survey. 95 pages. <http://www.state.nj.us/dep/njgs/pricelst/greport/gsr32.pdf>

French, Mark. 2004. Aquifer Recharge Potential for NJ Watershed Management Area 3, (Pompton, Wanaque and Ramapo Rivers), NJ. NJDEP, NJGS. GIS data. <http://www.njgeology.org/geodata/>

Heath, Ralph C. 1983. Basic Ground-Water Hydrology. U.S. Geological Survey Water-Supply Paper 2220. U.S. Geological Survey in cooperation with the North Carolina Department of Natural Resources and Community Development. 91 pages. <http://pubs.er.usgs.gov/publication/wsp2220>

Lewis-Brown, Jean C. and Eric Jacobsen. 1995. Hydrogeology and Ground-water flow, Fractured Mesozoic Structural-Basin Rocks, Stony Brook, Beden Brook, and Jacobs Creek Drainage Basins, West-Central New Jersey. US Geological Survey Water-Resources Investigations Report 94-4147. West Trenton, New Jersey, 89 pages. <http://pubs.usgs.gov/wri/1994/4147/report.pdf>

NJDEP, Office of Environmental Planning (OEP). June 1996. Water for the 21st Century: The Vital Resource, New Jersey Statewide Water Supply Plan. 205 pages. <http://rucore.libraries.rutgers.edu/rutgers-lib/32328/>

NJ Water Supply Authority. June 2002. Ground Water in the Raritan River Basin: A Technical Report for the Raritan Basin Watershed Management Project. 40 pages. http://www.raritanbasin.org/Alliance/ground_water.html

Office of the New Jersey State Climatologist (ONJSC). No Date. Climate Overview. <http://climate.rutgers.edu/stateclim/?section=usc&target=NJCoverview>

E. Ground Water Quality

Atherholt, Thomas B., Judith B. Louis, John Shevlin, Karen Fell, and Sandra Krietzman. April 2009. The New Jersey Private Well Testing Act: An Overview. NJDEP Division of Science, Research and Technology <http://www.state.nj.us/dep/dsr/research/pwta-overview.pdf>.

Degener, Richard. August 24, 2011. West Cape May gets \$3 million to improve water, sewer infrastructure. Atlantic City Press. http://www.pressofatlanticcity.com/news/press/cape_may/west-cape-may-gets-million-to-improve-water-sewer-infrastructure/article_d6124d02-cecb-11e0-9b26-001cc4c002e0.html. Accessed November 27, 2017.

Dunne, Thomas and Luna B. Leopold. 1978. Water in Environmental Planning. W. H. Freeman and Company: New York, New York. 818 pages.

Lacombe, Pierre J. and Glen B. Carleton. 2002. Hydrogeologic Framework, Availability of Water Supplies, and Saltwater Intrusion, Cape May County, New Jersey. U.S. Geological Survey Water-Resources Investigations Report 01-4246. Prepared In Cooperation With The New Jersey Department of Environmental Protection. 151 pages. <https://pubs.usgs.gov/wri/wri014246/pdf/wrir01-4246.pdf>

New Jersey Drinking Water Watch: West Cape May Water Department. 2017. https://www9.state.nj.us/DEP_WaterWatch_public/JSP/Violations.jsp?tinwsys=265. Accessed November 29, 2017.

NJDEP Division of Water Supply and Geoscience. August 17, 2017. Private Well Testing Act (PWTA). http://www.nj.gov/dep/watersupply/pw_pwta.html

NJDEP Division of Water Supply and Geoscience. January 19, 2017. NJ Private Well Testing Act Data Summary (Sep. 2002 to Apr. 2014). Story Map (interactive map): <http://njdep.maps.arcgis.com/apps/MapSeries/index.html?appid=826ec9fae77543caa582a787d5f088e7>

NJDEP Division of Water Supply and Geoscience. August 17, 2017. Lead in Drinking Water. <http://www.nj.gov/dep/watersupply/dwc-lead.html>. Accessed August 26, 2017.

Steering Committee of the New Jersey Comparative Risk Project. March 2003. Final Report of the New Jersey Comparative Risk Project. 213 pages. <http://www.state.nj.us/dep/dsr/njcrp/lacombe>

West Cape May Water Department. 2016. Annual Drinking Water Quality Report (PWS ID# NJ0512001). Results from the Year 2015. <http://www.westcapemay.us/LinkClick.aspx?fileticket=LTvaSavJvw%3d&tabid=1990>

F. Ground Water Quality Standards

N.J.A.C. 7:9C Ground Water Quality Standards. Date Last Amended: July 22, 2010. 27 pages. http://www.nj.gov/dep/rules/rules/njac7_9c.pdf

NJDEP Bureau of Environmental Analysis, Restoration and Standards. September 15, 2017. Ground Water Quality Standards N.J.A.C 7:9C. <http://www.nj.gov/dep/wms/bears/gwqs.htm>

G. Ground Water Discharges

NJDEP, Division of Water Quality (DWQ), Bureau of Nonpoint Pollution Control (BNPC). July 18, 2007. New Jersey Pollution Discharge Elimination System (NJPDES) Regulated Facility Locations, Edition 20070718 (Envr_NJEMS_NJPDES_all). GIS data. <http://njogis-newjersey.opendata.arcgis.com/datasets/njdep-non-point-wastewater-sites>

NJDEP. January 5, 2009. N.J.A.C. 7:14A-1 Pollutant Discharge Elimination System Date last amended: January 5, 2009. <http://www.state.nj.us/dep/dwq/714a.htm>

H. Contaminated Sites

[NJDEP Data Miner](https://www13.state.nj.us/DataMiner). June 13, 2017. <https://www13.state.nj.us/DataMiner>

NJDEP Site Remediation Program (SRP). September 15, 2017. [Site Remediation Program](http://www.nj.gov/dep/srp/). <http://www.nj.gov/dep/srp/>. Accessed October 26, 2017.

NJDEP Site Remediation Program (SRP). August 28, 2017. [Deed Notice Extent in New Jersey \(Envr_mon_soil_DNA\)](http://njogis-newjersey.opendata.arcgis.com/datasets/076a3d08a34348d2bdb3c109ec349e53_17). http://njogis-newjersey.opendata.arcgis.com/datasets/076a3d08a34348d2bdb3c109ec349e53_17

NJDEP Site Remediation Program (SRP). June 13, 2017. [Community Relations Site List](http://www.nj.gov/dep/srp/community/sites/). <http://www.nj.gov/dep/srp/community/sites/>. Accessed October 26, 2017.

NJDEP Site Remediation Program (SRP). April 25, 2017. [Underground Storage Tanks \(USTs\)](http://www.nj.gov/dep/srp/bust/). <http://www.nj.gov/dep/srp/bust/>. Accessed November 26, 2017.

NJDEP Site Remediation Program (SRP). February 14, 2017. [Closed Sites with Remediated Contamination](http://www.nj.gov/dep/srp/kcs-nj/). <http://www.nj.gov/dep/srp/kcs-nj/>. Accessed October 26, 2017.

NJDEP Site Remediation Program (SRP). March 29, 2016. [Classification Exception Areas-Well Restriction Areas for New Jersey \(Envr_mon_gw_CEA\)](http://www.state.nj.us/dep/gis/digidownload/zips/statewide/Envr_mon_gw_CEA.zip). GIS Data. http://www.state.nj.us/dep/gis/digidownload/zips/statewide/Envr_mon_gw_CEA.zip and Guidance documents: http://www.nj.gov/dep/srp/guidance/cea/cea_guide.htm

NJDEP Site Remediation Program (SRP). May 7, 2012. [Administrative Requirements for the Remediation of Contaminated Sites and Technical Requirements for Site Remediation](http://www.nj.gov/dep/rules/adoptions/adopt_20120507a.pdf). Adopted Amendments, Adopted Repeals and Adopted New Rules. 687 pages. http://www.nj.gov/dep/rules/adoptions/adopt_20120507a.pdf

I. Ground Water Level

USGS. 2017a. Real-Time Ground-Water-Level Network. Site Number: 390156074533401 - 090333-- Pump Pond N Obs. <https://groundwaterwatch.usgs.gov/AWLSites.asp?S=390156074533401>

USGS. 2017b. Groundwater Watch. Site Number: USGS 385607074555201 090150-- West Cape May 1 Obs https://waterdata.usgs.gov/nwis/inventory/?site_no=385607074555201&agency_cd=USGS

Internet Resources: Ground Water

Drinking Water

West Cape May Water Department –

Annual Drinking Water Report:

<http://www.westcapemay.us/borough-departments/annual-drinking-water-report.aspx>

NJ Drinking Water Watch:

https://www9.state.nj.us/DEP_WaterWatch_public/JSP/WSDetail.jsp?tinwsys=265

Your Utility and Its Water Quality: http://www.nj.gov/dep/watersupply/dwc_systems.html

Ground Water and Drinking Water (US EPA): <https://www.epa.gov/ground-water-and-drinking-water>

Lead in Drinking Water: <http://www.nj.gov/dep/watersupply/dwc-lead.html>

NJ and Federal Drinking Water Standards: http://www.nj.gov/dep/watersupply/pdf/dw_standards_2_2005.pdf

NJDEP Private Well Testing Act: http://www.nj.gov/dep/watersupply/pw_pwta.html

Story Map: NJ Private Well Testing Act Data Summary (Sep. 2002 to Apr. 2014)

<http://njdep.maps.arcgis.com/apps/MapSeries/index.html?appid=826ec9fae77543caa582a787d5f088e7>

Drought and Ground Water Level

NJDEP Drought Information: www.njdrought.org

USGS New Jersey Real-Time Groundwater Level Network

NJ: <https://groundwaterwatch.usgs.gov/NetMapT1L2.asp?ncd=rtn&sc=34>

Canal 5 Obs <https://groundwaterwatch.usgs.gov/AWLSites.asp?S=385748074553301&ncd=NJN>

West Cape May 1 <https://groundwaterwatch.usgs.gov/AWLSites.asp?mt=g&S=385607074555201&ncd=awl>

Ground Water Quality

NJDEP Ground Water Monitoring: <http://www.state.nj.us/dep/wmm/bfbm/groundwater.html>

Comparative Risk Project: <http://www.state.nj.us/dep/dsr/njcrp/>

[NJDEP grants for Underground Storage Tank removal & remediation: http://www.nj.gov/dep/srp/finance/ustfund/](http://www.nj.gov/dep/srp/finance/ustfund/)

Hydrogeology

Aquifer and Well Characteristics in New Jersey (USGS): http://wwwnj.er.usgs.gov/gw/table_1.html

NJGS Hydrogeologic Data (horizontal hydraulic conductivity values, transmissivity values, and vertical hydraulic conductivity values): <http://www.state.nj.us/dep/njgs/geodata/dgs02-1.htm>

Radioactivity in Ground Water

NJDEP: A South Jersey Homeowner's Guide to Radioactivity in Drinking Water:

<http://www.state.nj.us/dep/rpp/rms/agreedown/radwater.pdf>

US EPA: Basic Information about the Radionuclides Rule.

<http://water.epa.gov/lawsregs/rulesregs/sdwa/radionuclides/basicinformation.cfm>

Other

[NJDEP Data Miner: https://www13.state.nj.us/DataMiner](https://www13.state.nj.us/DataMiner)

NJDEP Laws & Rules: <http://www.nj.gov/dep/landuse/lawsregs.html>

NJDEP Rules & Regulations (current & proposed): <http://www.nj.gov/dep/rules/>

Site Remediation Reform Act, N.J.S.A. 58:10C.: <http://www.nj.gov/dep/srp/regs/statutes/srra.pdf>

Brownfield & Contaminated Sites Act, N.J.S.A. 58:10B: <http://www.nj.gov/dep/srp/regs/statutes/bcsra.pdf>

Spill Compensation and Control Act: http://www.nj.gov/dep/srp/regs/statutes/spill_act.pdf

Industrial Site Recovery Act: <http://www.nj.gov/dep/srp/regs/statutes/isra.pdf>

Underground Storage Tanks: http://www.nj.gov/dep/rules/rules/njac7_14b.pdf

Ground Water Quality Standards: http://www.nj.gov/dep/rules/rules/njac7_9c.pdf

NJDEP Division of Water Monitoring and Standards: <http://www.nj.gov/dep/wms/bears/index.html>

NJ Geological Survey Home Page: <http://www.state.nj.us/dep/njgs/index.html>

Underground Storage Tanks: <http://www.nj.gov/dep/srp/bust/>

USEPA Ground Water and Drinking Water: <https://www.epa.gov/ground-water-and-drinking-water>

USGS - New Jersey District - Ground Water Information (USGS): <https://waterdata.usgs.gov/nj/nwis/gw>

USGS - Water Resources of NJ: <https://nj.usgs.gov/>

6: SURFACE WATER

A. Introduction Surface Water

As described in **Section 5A**'s overview of the water cycle, *surface water* is water that is visible above the ground surface, including creeks, rivers, ponds, lakes, and wetlands. Surface water is generally hydraulically connected to



ground water, although the interactions are difficult to observe and are affected by variations in weather and human activities. Streams connect with ground water in three basic ways: Streams can gain water from ground water through the streambed, lose water to ground water through the streambed, or have both gaining and losing reaches (Winter et. al., 1998).

Watersheds

A *watershed* (or basin) is the land area within the confines of a drainage divide in which all surface runoff will drain into a river, river system, or body of water. The Borough of West Cape May is within the Delaware Bay and Atlantic coastal watersheds, which covers the Cape May peninsula of New Jersey.

Watershed Management Areas



Watershed management is the process of managing and protecting all of the water resources within the area of a watershed, rather than on a site-specific basis. The NJDEP recognizes that watersheds are “nature’s boundaries,” and has established a watershed management approach (NJDEP, January 1997). A watershed management approach is based on three key components: 1) a geographic focus; 2) continuous improvement based on sound science; and 3) partnerships/stakeholder involvement. More information concerning watershed management is presented in **Section 10D**. NJDEP has divided the state's watersheds into 20 *Watershed Management Areas (WMAs)*. West Cape May falls within *WMA 16: Cape May* (see top left inset in **Figure 6a**).

Hydrologic Unit Codes (HUC)

The classification system used by the NJDEP assigns each sub-watershed a *14-digit Hydrologic Unit Code (HUC14²⁰)*. The HUC14 is a hierarchical system where the first 2 digits refer to the USGS Water Resources Region and the first 4 digits (also known as a HUC4) refer to the major

"Water is vital to life and comprises an invaluable natural resource which is not to be abused by any segment of the State's population or economy."
(NJDEP NJAC 7:9B, April 4, 2011)

²⁰ The HUC14s have a minimum size of 3,000 acres, although some basins are defined with smaller areas. At other times, small subwatershed units are combined. Note, however, that the current HUC12 numbers are not consistent with this system.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.





Figure 6b: West Cape May Subwatersheds and SWQS
All of the surface waters within West Cape May are categorized as FW2-NT/SE1. The classification is used for waterways in which there may be a fresh/salt water interface. Some Pond Creek tributaries just west of the borough have a C1 designation, which indicates exceptional significance and provides additional protections for those streams.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

drainage basin, or sub-region. Therefore, a HUC2 of “02” is in the Mid-Atlantic Region, and a HUC4 of “0204” is in the Delaware-Mid Atlantic Coastal major drainage basin (USGS, June 23, 2017).

The area of land where water flows generally west towards the Delaware Bay is assigned a HUC8 of “02040206,” and every sub-watershed within this basin has a HUC that starts with “02040206.” The western half of West Cape May is in this area. Land that drains east to the Atlantic Ocean (Great Egg Harbor to Cape May) is assigned a HUC8 of “02040302,” including the eastern half of West Cape May. HUC14 sub-watersheds encompassing the Borough of West Cape May are shown in **Figure 6a** and listed in **Table 6.1**.

Table 6.1: Hydrologic Unit Codes for West Cape May’s Sub-watersheds

HUC4	HUC8	14-Digit Hydrologic Unit Code (HUC14)	Sub-watershed Name
0204	02040206	02040206230070	Pond Creek / Cape May Canal West
0204	02040302	02040302080090	Cape May Harbor & Bays (below Rt 47)
Source: NJDEP NJGS, February 25, 2011			

B. Floodplains

A *floodplain* is the land along a river or stream that is subject to periodic flooding when the river or stream overflows its banks. As required by the Flood Disaster Protection Act of 1973, the Federal Emergency Management Administration (FEMA) is responsible for delineating floodplains.

According to FEMA, "Everyone lives in some type of flood zone." (FEMA, June 21, 2007). FEMA defines these geographic areas based on studies of flood risk.

FEMA provides accurate flood hazard and risk data to states and communities to guide mitigation actions. The National Flood Insurance Program (NFIP) is the basis of the NFIP regulations and flood insurance requirements. Flood hazard mapping is an important part of the NFIP. FEMA uses the best available technical data, such as statistical information on river flows, to create the *Flood Insurance Rate Maps* (FIRMs) that show the flood zone boundaries (FEMA, June 29, 2017).

The flood zone boundaries shown in **Figure 6c** are produced using FEMA’s National Flood Hazard Layer (FEMA, October 4, 2017). Special Flood Hazard Areas (SFHAs) are defined as areas subject to inundation by a flood having, on average, about 1 in 100 chance in any given year, also referred to as the 1% annual chance flood²¹ (FEMA, March 7, 2017). Below are brief definitions of the FEMA flood zones that occur within West Cape May.

Zones with a high-risk of flooding, or SFHAs, include *Zone A* and *Zone AE*. *Zone A* corresponds to the 1% annual chance floodplains that are determined by approximate methods of analysis (i.e., not with Base Flood Elevations). *Zone AE* corresponds to the 1% annual chance floodplains that are determined by detailed methods of analysis, which includes detailed hydraulic analyses to determine Base Flood Elevations. In communities such as West Cape May that participate in the NFIP, all homeowners in Zones A and AE are required to get flood insurance in order to get a loan from a federally regulated lender (FEMA, March 7, 2017; FEMA, August 6, 2017).

²¹ Flood designations are based on statistical averages, not the number of years between big floods. The term "100-year flood" does not mean a flood that happens once every 100 years. It is a statistical designation that there is a 1 in 100 chance that a flood of any given size will be equaled or exceeded during any year. Changes and variability in climate and land use over time can change flood frequency (Dinicola, 2005).





Figure 6c: West Cape May Flood Zones

According to FEMA, approximately 56% of the borough is within the AE Zone (1% annual chance of flooding), while another 13% has a 0.2% annual chance of flooding.



Flood Facts

- Floods and flash floods happen in all 50 states.
- Hurricanes, winter storms and snowmelt are common (but often overlooked) causes of flooding.
- New land development can increase flood risk, especially if the construction changes natural runoff paths.
- Federal disaster assistance is usually a loan that must be paid back with interest.
- If you live in a Special Flood Hazard Area (SFHA) or high-risk area and have a Federally backed mortgage, your mortgage lender requires you to have flood insurance
- 20 to 25% of all flood claims are filed in low to moderate flood risk areas.

(FEMA, July 25, 2017; FEMA, June 21, 2007)

Areas in *Zone X*, which includes approximately 32% of West Cape May, have low to moderate risk of flooding and are not in the SFHAs. They correspond to areas outside the 1% annual chance floodplain, areas of 1% annual chance sheet flow²² flooding where average depths are less than 1 foot, areas of 1% annual chance stream flooding or where the contributing drainage area is less than 1 square mile. No Base Flood Elevations or depths are shown within this zone. Areas with a 0.2% annual chance of flooding (typically referred to as the 500 year flood) are not considered high risk. The zone includes areas of little hazard, such as those with average depths of less than 1 foot and minimal hazard, such as ponding and local drainage problems. Insurance purchase is not required in this zone (FEMA, March 7, 2017).

Floodplains in West Cape May are shown in **Figure 6c**, based on FEMA determinations (FEMA, October 4, 2017). Approximately 56% of the borough is within the 1% annual chance of flooding, while another 13% has a 0.2% annual chance of flooding.

Floodplain management is the operation of a community program of corrective and preventative measures for reducing flood damage. Community involvement is an important element in making flood insurance available to home and business owners. These measures may include zoning, subdivision, or building requirements, and special-purpose floodplain ordinances. Riparian buffer and wetlands protection regulations and ordinances can also reduce flood damage by protecting those areas most susceptible to flooding and providing natural flood control. These efforts benefit downstream areas, as well.



West Cape May has a flood hazard protection ordinance (Chapter XXIII of the Borough Code) and plan (see **Internet Resources**).

Tidal monitoring and flood forecast gages near West Cape May are shown in **Table 6.2**. Floods are covered in **Section 2**.



²² Sheet flow, or overland flow, is flow that occurs overland in places where there are no defined channels, so the flood water spreads out over a large area at a uniform depth.

Table 6.2: National Weather Service Advanced (NOAA) Hydrologic Prediction Service (flood predictions)



C. Wetlands

A *wetland* is a transitional area between aquatic and terrestrial ecosystems. Wetlands are those areas that are inundated or saturated by surface water or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions, commonly known as hydrophytic vegetation. To determine if an area is a wetland, the vegetation (plants that like wet conditions), soils (wetland types, which often show mottling) and hydrology (low spots or evidence of water) are evaluated. A *transition area*, or buffer, is an area of land adjacent to a freshwater wetland that minimizes adverse impacts on the wetland or serves as an integral component of the wetlands ecosystem (N.J.S.A. 13:9B-3 in NJDEP Division of Land Use Management, July 16, 1998).

In the past, wetlands were often regarded as wastelands – only useful when drained and filled. In contrast, a 1978 Tufts University study showed that one acre of wetland provides at least \$153,000 (1978 dollars) of public value, considering proven monetary benefits of flood protection, pollution reduction, water supply, recreation and aesthetics (Fair, 2004). Some of the benefits of wetlands include:

- Wetlands protect drinking water by filtering out pollutants and sediments that would otherwise obstruct and contaminate our waters.
- Wetlands soak up runoff from heavy rains and snow melts, providing natural flood control.
- Wetlands release stored waters during droughts.
- Wetlands provide critical habitats for a major proportion of the state’s fish and wildlife, including many endangered, commercial and recreational species.
- Wetlands provide high quality open space for recreation and tourism (NJDEP Land Use Regulation, August 16, 2017).

The value of wetlands was not broadly accepted until at least the 1970s and 1980s. By then, more than half of the country's wetlands had been destroyed (NJDEP Land Use Regulation, August 16, 2017). Loss of wetlands has resulted in erosion, flooding, sedimentation, and

decreased populations of many types of wildlife. Structures built in wetlands suffer from frost heaving and other structural problems.

New Jersey protects wetlands under the 1987 New Jersey Freshwater Wetlands Protection Act (N.J.S.A. 13:9B) and Rules (N.J.A.C. 7:7A) (NJDEP Division of Land Use Management, July 16, 1998 and July 17, 2017). Under these, NJDEP regulates virtually all activities proposed within wetlands and



transition areas or buffers around freshwater wetlands, including cutting of vegetation, dredging,

excavation or removal of soil, drainage or disturbance of the water level, and filling or discharge of any materials. Development that would impair the wetland's ability to provide the values listed above (filtration, flood control, etc.) is prohibited. There are limited exemptions for existing farming, ranching, or forestry operations.



On-site inspection (direct testing and observation of soils, hydrology and vegetation) by a qualified professional is needed prior to making any disturbance within a wetland or transition area. Only an official determination from NJDEP, called a *Letter of Interpretation* (LOI) can verify the presence, absence, or boundaries of freshwater wetlands and transition areas on a site. Copies of these maps are filed at the NJDEP and the township building, but unfortunately, NJDEP does not digitize these determinations into a GIS layer²³.

In addition to defining the boundary of the wetland, the LOI establishes the value of the wetland, which will determine the width of the regulated transition area. *Ordinary Value* wetlands, such as man-made drainage ditches and swales, have a 0 foot buffer. *Intermediate Value* wetlands have a 50 foot buffer, which includes those wetlands not included in the definitions of Ordinary or Exceptional value. *Exceptional Value* wetlands have a 150 foot buffer width. Exceptional Value wetlands include wetlands that provide habitat for endangered and threatened species and those contiguous with FW-1, FW-2 Trout Production waters and their tributaries, and Category 1 classified streams (see **Section 6D** for descriptions of stream classifications, below). A determination of threatened and endangered species habitat is provided by using the Landscape Project data (see **Section 7F**).

There are approximately 266 acres of wetlands within West Cape May, or 35% of the borough (NJDEP, February 17, 2015). The wetlands shown in **Figure 6d** were determined by selecting all wetlands land use types from NJDEP's 2012 Land Use GIS data. **Figure 6d** provides guidance on where wetlands are found in West Cape May. This dataset is intended to serve as a resource for analysis rather than regulatory delineations because it is derived from aerial photos rather than on-site surveys. A transition area (buffer) width of 150 feet is mapped in **Figure 6d**, because the GIS data does not determine the value of each wetland. The actual transition area width (0, 50 or 150') required by the NJDEP is determined in the LOI.

There are several types of freshwater wetlands in West Cape May, such as deciduous wooded wetlands, saline marsh (low and high), deciduous scrub/shrub wetlands, and agricultural wetlands (see **Section 7A** and **Figure 7a**).

²³ Digitizing involves giving latitude and longitude coordinates to areas and lines to depict mapped features.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Figure 6d: West Cape May Wetlands

There are approximately 266 acres of wetlands within West Cape May, or 35% of the borough. Buffer areas indicated on the map are based on the maximum width of 150 feet. Actual buffer requirements may be less, depending on the values of individual wetlands.

D. Surface Water Quality Standards

Surface Water Quality Standards (SWQS) are the rules in chapter N.J.A.C. 7:9B that set forth designated uses, use classifications, and water quality criteria for the State's waters based upon the uses, and the NJDEP's policies concerning these uses, classifications and criteria, which are necessary to protect the State's waters. The SWQS operate in conformance with the Federal Water Pollution Control Act (33 U.S.C. 1313(c)), commonly known as the Clean Water Act (CWA), and the Federal Water Quality Standards Regulation at 40 CFR 131.

According to the Surface Water Quality Standards N.J.A.C. 7:9B,

“Water is vital to life and comprises an invaluable natural resource which is not to be abused by any segment of the State’s population or economy. It is the policy of the State to restore, maintain and enhance the chemical, physical and biological integrity of its waters, to protect the public health, to safeguard the aquatic biota, protect scenic and ecological values, and to enhance the domestic, municipal, recreational, industrial, agricultural and other reasonable uses of the State’s waters.

“The restoration, maintenance and preservation of the quality of the waters of the State for the protection and preservation of public water supplies is a paramount interest of the citizens of New Jersey.... Toxic substances in waters of the State shall not be at levels that are toxic to humans or the aquatic biota, or that bioaccumulate in the aquatic biota so as to render them unfit for human consumption.... Human health-based ambient criteria have been established in freshwaters due to consumption of fish and water, and in saline water due to consumption of fish. For carcinogens, the criteria have been established at levels which would result in no greater than a one-in-one-million lifetime excess cancer risk. For non-carcinogens, the criteria have been established which would result in no appreciable risk of deleterious effect.” (NJDEP Land Use Management, Water Monitoring and Standards, April 4, 2011).

According to the designated uses under the SWQS, NJDEP assigns *surface water classifications* to each stream in order to group waters and assign water quality criteria. Designated uses include potable water, propagation of fish and wildlife, recreation, agricultural and industrial supplies, and navigation. The *criteria* are numerical targets for constituent concentrations (such as toxic pollutants) or narratives that describe in-stream conditions to be attained, maintained or avoided, so that the specified uses are protected for the different use classifications.

The SWQS are used by several NJDEP programs, including the New Jersey Pollutant Discharge Elimination System program, Site Remediation program, Stream Encroachment, Land Use Regulation Program and Total Maximum Daily Loads (TMDLs, see **Section 6E**).

Table 6.3 describes the definitions of the surface water classifications. In **Figure 6b**, “category” is shown, which is a compendium of all surface water classification designations for a given water body. Category describes a stream's surface water classification in terms of its general surface water class (e.g. SE1), its trout water status (e.g. NT is non-trout) and its antidegradation status (e.g. C1). All of the surface waters within West Cape May are categorized as FW2-NT/SE1 (as defined by N.J.A.C. 7:9B(f):

“FW2-NT/SE1 (or a similar designation that combines two classifications) means a waterway in which there may be a salt water/fresh water interface. The exact point of demarcation between the fresh and saline waters must be determined by salinity measurements and is that point where the salinity reaches 3.5 parts per thousand at mean high tide. The stream is classified as FW2-NT in the fresh portions (salinity less than or equal to 3.5 parts per thousand at mean high tide) and SE1 in the saline portions.” (NJDEP, April 4, 2011)

- **FW2-NT/SE1**
 - **Cape Island Creek** - Forms the eastern boundary of the borough



- **Cape Island Creek UNT** – Several small streams in the eastern portion of the borough are small unnamed tributaries of Cape Island Creek.

- **Pond Creek and Pond Creek UNT** – Approximately the western 40% of West Cape May drain to this watershed.

The *Category One (C1)* antidegradation designation provides streams with additional protections that help prevent water quality degradation and discourage development where it would impair or destroy natural resources and water quality. Waterways can be designated C1 because of exceptional ecological significance, exceptional water supply significance, exceptional recreational significance, exceptional shellfish resource, or exceptional fisheries resource (NJDEP Water Monitoring and Standards, June 15, 2017). There are no C1 streams within West Cape May, but Pond Creek and an unnamed tributary of Pond Creek, flowing from the western

part of the borough are designated FW2-NTC1/SE1.

The antidegradation provisions of the SWQS are triggered when an applicant proposes an activity that has the potential to lower water quality. Previously approved wastewater discharges authorized through the NJPDES program as well as existing developments are not subject to the antidegradation policies unless a new or expanded activity is proposed. Under the Stormwater Management rules (N.J.A.C. 7:8) and the Flood Hazard Area Control Act rules (N.J.A.C. 7:13), for certain activities proposed adjacent to waters designated as C1, 300 foot buffers must be maintained in a natural state adjacent to all C1 waters and upstream tributaries of C1 waters (including named and unnamed tributaries), unless the disturbance is less than one acre and new impervious surface is less than 0.25 acres. However, where the buffer is already disturbed, the width may be reduced in the disturbed area, but will not be permitted to extend less than 150 feet from either bank. The buffer will not affect existing development. The buffer requirement can also be adjusted to reflect local conditions through the approval of a stream corridor protection plan as part of a regional stormwater management plan (NJDEP Water Monitoring and Standards, June 15, 2017).

Most of the state's Category One waters were designated in 1985 amendments to the SWQS, however no specific basis for these upgrades was documented. C1 upgrades after that were documented, based on their value as trout production (FW2-TP) waters or exceptional ecological significance or exceptional water supply significance.

Table 6.3: Surface Water Quality Standards Classification

Category	Definition
Freshwater General Surface Water Class	
FW1	FW1 means those fresh waters, as designated in N.J.A.C. 7:9B-1.15(j), that are to be maintained in their natural state of quality (set aside for posterity) and not subjected to any man-made wastewater discharges or increases in runoff from anthropogenic activities. These waters are set aside for posterity because of their clarity, color, scenic setting, other characteristic of aesthetic value, unique ecological significance, exceptional recreational significance, exceptional water supply significance or exceptional fisheries resource(s).
FW2	FW2 means the general surface water classification applied to those fresh waters that are not designated as FW1 or Pinelands Waters. In all FW2 waters the designated uses are: 1. Maintenance, migration and propagation of the natural and established biota; 2. Primary contact recreation; 3. Industrial and agricultural water supply; 4. Public potable water supply after conventional filtration treatment (a series of processes including filtration, flocculation, coagulation, and sedimentation, resulting in substantial particulate removal but no consistent removal of chemical constituents) and disinfection; and 5. Any other reasonable uses.
Saline Water General Surface Water Class	
SC	Saline waters means waters having salinities generally greater than 3.5 parts per thousand at mean high tide. SC means the general surface water classification applied to coastal saline waters.
SE	SE means the general surface water classification applied to saline waters of estuaries.
Trout Water Status - this is for information only and does not affect the water quality criteria for those waters.	
TP	Trout production means waters designated at N.J.A.C. 7:9B-1.15I through (i) for use by trout for spawning or nursery purposes during their first summer.
TM	Trout maintenance means waters designated at N.J.A.C. 7:9B-1.15I through (i) for the support of trout throughout the year.
NT	Nontrout waters means fresh waters that have not been designated in N.J.A.C. 7:9B-1.15I through (h) as trout production or trout maintenance. These waters are generally not suitable for trout because of their physical, chemical, or biological characteristics, but are suitable for a wide variety of other fish species.
Antidegradation	
ONRW	Outstanding National Resource Waters means high quality waters that constitute an outstanding national resource (for example, waters of National/State Parks and Wildlife Refuges and waters of exceptional recreational or ecological significance). Waters classified as FW1 waters and Pinelands waters are Outstanding National Resource Waters.
FW1/Non-degradation	Nondegradation waters means those waters set aside for posterity because of their clarity, color, scenic setting, other characteristic of aesthetic value, unique ecological significance, exceptional recreational significance, or exceptional water supply significance. These waters include all waters designated as FW1.
C1	Category one waters means those waters designated in the tables in N.J.A.C. 7:9B-1.15(c) through (i), for purposes of implementing the antidegradation policies set forth at N.J.A.C. 7:9B-1.5(d), for protection from measurable changes in water quality based on exceptional ecological significance, exceptional recreational significance, exceptional water supply significance or exceptional fisheries resource(s) to protect their aesthetic value (color, clarity, scenic setting) and ecological integrity (habitat, water quality and biological functions).
C2	Category two waters means those waters not designated as Outstanding National Resource Waters or Category One at N.J.A.C. 7:9B-1.15 for purposes of implementing the antidegradation policies set forth at N.J.A.C. 7:9B-1.5(d).
Source: NJDEP Land Use Management, Water Monitoring and Standards, April 4, 2011	

E. Integrated List and Total Maximum Daily Loads

States are required by the Federal Clean Water Act (US Federal Water Pollution Control Act, January 4, 2011) to develop a biennial Water Quality Inventory Report (required under Section 305(b) of the act) and a List of Water Quality Limited Segments (required under Section 303(d)). Since 2001, the USEPA has recommended that states integrate these two, producing the

Integrated List. The goal is to provide an effective tool for maintaining high quality waters where designated uses (designated by the SWQS, discussed above in **Section 6D**) are attained, and improving the quality of surface waters that do not attain their designated uses (NJDEP BEARS, May 2017 (final)).

The Integrated List is subject to regulatory requirements, which include public participation and submission to the USEPA for approval and adoption. The Integrated List identifies the status of all applicable designated uses for every assessment unit (usually by HUC14²⁴ sub-watershed) by labeling the results of each designated use assessment as *Fully Supporting*, *Not Supporting*, or *Insufficient Information* (see **Table 6.4**).

The NJDEP is required to use all existing and readily available data to assess water quality for the Integrated List. A methods document summarizes each step in the assessment process; to evaluate stations and data quality, combine stations to evaluate an assessment unit, assess designated uses, rank and prioritize assessment units that do not attain designated uses, develop a monitoring and assessment plan and provide for public participation (NJDEP, DWMS, BEARS, February 2015).

The 2014 Integrated List, which summarizes whether or not the surface water quality of West Cape May’s two subwatersheds meet the SWQS, is shown in **Table 6.4** and **Figure 6e**. The water quality supports the recreational use of waters in the eastern portion of the borough, while other uses are either not supported or there is insufficient information to assess them. No changes were noted compared to the 2012 assessment. **Table 6.5** displays more information about the impaired waters within West Cape May.

Table 6.4: 2012 Integrated List (Overview)

Assessment Unit (HUC14)	Sub-watershed	Designated Use*					
		Aquatic Life General	Aquatic Life Trout	Recreation	Water Supply	Shellfish	Fish Consumption
02040302080090	Cape May Harbor & Bays (below Rt 47)	Not Supporting	N/A	Fully Supporting	N/A	Not Supporting	Insufficient Data
02040206230070	Pond Creek / Cape May Canal West	Not Supporting	N/A	Insufficient Data	N/A	Not Supporting	Not Supporting
Minimum Suite of Parameters Needed to Determine if Water Quality is “Fully Supporting” a Use: General Aquatic Life - Biological data Aquatic Life - Trout Biological data and Temperature and DO Recreation - Pathogenic Indicator Bacteria Shellfish Harvest for Consumption - Total Coliform Public Water Supply - Nitrate Fish Consumption - Fish tissue data For a full list of parameters for each designated use, see Appendix A of the 2014 methods document (NJDEP, DWMS, BEARS, February 2015)							
Source: NJDEP Water Monitoring and Standards, May 2017 (final); NJDEP, DWMS, BEARS, February 2015							

When surface waters do not meet the SWQS, *Total Maximum Daily Loads* (TMDLs) must be developed, as specified under Section 303(d) of the Federal Clean Water Act (US Federal Water Pollution Control Act, January 4, 2011). A TMDL identifies all the contributors to surface water quality impacts and sets goals for load²⁵ reductions for specific pollutants in

²⁴ HUC14 = 14-digit Hydrologic Unit Code (see **Section 6a** for definition)

²⁵ Load is the total amount of material (pollutants) entering the system from one or multiple sources; measured as a rate in weight per unit time (USEPA, 2017).





order to meet the SWQS. Regulations concerning TMDLs are contained in EPA's Water Quality Planning and Management Regulations (USEPA, 2017).

TMDLs represent the assimilative capacity of surface water for a given parameter of concern. The development of TMDLs includes balancing the impacts from point sources, nonpoint sources and natural background levels of a specific pollutant. The TMDL then quantifies the amount of a pollutant a water body can assimilate without violating a state's water quality standards and allocates that load capacity to known point and nonpoint sources in the form of waste load allocations (WLAs) for point sources, load allocations (LAs) for nonpoint sources, plus a margin of safety (MOS). Load allocations (for nonpoint source pollution) consist of identifying categories of nonpoint sources that contribute to the parameters of concern, followed by recommendations for implementation measures for specific load reductions. Examples include best management practices (BMPs), including structural (stormwater runoff controls) and non-structural (local ordinances for stormwater management and nonpoint source pollution control) mechanisms for addressing the water quality parameter(s) of concern (NJDEP Division of Watershed Management, March 9, 2017).

Table 6.5: Integrated Water Quality Assessment (Details)

Use	Attainment	Cause	First on 303(d) List	TMDL Priority*	Source
02040302080090 Cape May Harbor & Bays (below Rt 47)					
Aquatic Life	N	Oxygen, Dissolved	2010	Medium	Urban Runoff/ Storm Sewers
Shellfish	N	Total Coliform	2006	Completed Sublist 4	
02040206230070 Pond Creek / Cape May Canal West					
Fish Consumption	N	PCB in Fish Tissue	2006	Low Sublist L	Contaminated Sediments
Aquatic Life	N	Phosphorus	2004	Completed Sublist 4	Urban Runoff/ Storm Sewers
Shellfish	N	Total Coliform	2014	Completed Sublist 4	Urban Runoff/ Storm Sewers
*Designated Uses: N = Not Supporting					
*Medium priority = NJDEP expects to complete TMDL in the near future, but not within the next two years.					
Low priority = NJDEP does not expect to complete TMDL in the immediate or near future.					
Source: NJDEP Water Monitoring and Standards, May 2017 (final)					

Waters requiring TMDLs are identified and prioritized in the Integrated Water Quality Assessment. After the Integrated List is approved, the NJDEP writes a TMDL report, which is a proposed Water Quality Management Plan Amendment. When this is published in the NJ Register for public review and comment, the TMDL is considered *proposed*. NJDEP then considers comments received during public comment and finalizes the TMDL report, and the TMDL is considered *established* when it is formally submitted to the US EPA Region 2 for thirty-day review. The TMDL is considered *approved* when the US EPA Region 2 approves it.




Legend

West Cape May Borough

Sublists

- Sublist 2 - supporting use
- Sublist 3 - insufficient information
- Sublist 4 - not supporting use; TMDL completed
- Sublist 5 - not supporting use



Data Sources: NJDEP

Note: Map accuracy is limited to the accuracy and scale of the original data sets.

Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Prepared by Kratzer Environmental Services

Figure 6e: West Cape May 2014 Integrated List

The Clean Water Act requires the NJDEP to assess the state's surface waters with respect to whether the water quality is good enough to support the designated uses. The surface waters in and around West Cape May are designated to be used for aquatic life (general, i.e. not trout), shellfish consumption, recreation and fish consumption. The assessments are done on a HUC14 level (see Section 6A and Figure 6a). Assessment details are presented in Table 6.4 and Table 6.5. The water quality of waters in the eastern portion of the borough support recreational use. Other uses are either not supported or there is insufficient information to assess them.

Next, the TMDL is referred to as *adopted* when the EPA-approved TMDL is adopted by NJDEP as a water quality management plan amendment and the adoption notice is published in the NJ Register (NJDEP BEARS, March 9, 2017).

The NJDEP developed a TMDL to address total coliform (bacterial) impairments based on the 2004 Integrated List of Waterbodies in twenty waterbodies in WMA 16. In that list, waterbodies that did not fully support shellfish harvest according with National Shellfish Sanitation Program (NSSP) criteria were determined to be impaired. New Jersey Surface Water Quality Standards (SWQS) specify that shellfish waters must meet the stringent guidelines of the NSSP to protect against the harvest of shellfish in waters where the sanitary quality could have health risks for consumers. The TMDL outlines actions such as source trackdown and installation of best management practices (BMPs) for stormwater to achieve load reduction targets (NJDEP and Tetra Tech, Inc. September 27, 2006).

No other TMDLs for either of West Cape May's subwatersheds are scheduled for completion within the next two years (NJDEP Water Monitoring and Standards, May 2017 (final)).

Amendment to the Cape May County and Lower Delaware Water Quality Management Plans Ten Total Maximum Daily Loads for Total Coliform to Address Shellfish-Impaired Waters in Watershed Management Area 16 Atlantic Coastal Water Region. Approved: September 27, 2006. Prepared by: New Jersey Department of Environmental Protection And With assistance provided by: U.S. Environmental Protection Agency, Region 2

F. Point Source Pollution

Point source pollution (as defined by N.J.A.C. 7:9B Surface Water Quality Standards) refers to discernible, confined, and discrete conveyance, including, but not limited to, any pipe, ditch, channel, tunnel, conduit, well, discrete fissure, container, rolling stock, concentrated animal feeding operation, landfill leachate collection system, vessel, or other floating craft, from which pollutants are or may be discharged. This term does not include return flows from irrigated agriculture (NJDEP Land Use Management, Water Monitoring and Standards, April 4, 2011).

Point source discharges are regulated by NJDEP under the New Jersey Pollutant Discharge Elimination System (NJPDES). There are 2 existing discharges within or on the border of West Cape May (see **Table 6.6** and **Figure 6f**) (NJDEP, Division of Water Quality, Bureau of Surface Water Permitting. June 14, 2017).

G. Nonpoint Source Pollution

Nonpoint source or NPS pollution is any man-made or man-induced activity, factor, or condition, other than a point source, from which pollutants are or may be discharged. Nonpoint pollution may temporarily or permanently change any chemical, physical, biological, or radiological characteristic of water from what was or is the natural, pristine condition of such water.

Impervious surfaces are materials that prevent the infiltration of water into the soil (e.g. parking lots, roads, buildings, sidewalks and compacted soil). The construction of impervious surfaces disrupts the natural water cycle, and is one of the more significant landscape impacts attributable to urbanization (Hasse and Lathrop, December 2016). When water flows off impervious surfaces, it is known as *stormwater*. Nonpoint source pollution is directly associated with stormwater.

Table 6.6: NJ Pollutant Discharge Elimination System Surface Water Discharges

PI #	NJPDES ID. #	Facility Name	Status*	Discharge Type*	Receiving Waters
47725	NJ0108341	Cape May Desalination Project	E	Industrial Category B	Cape Island Creek
744321	NJG0262498	Former Cape May MGP Site	E	B4B General Petroleum Product Cleanup	Cape Island Creek
<p>*Notes for Above Codes (NJDEP's codes and definitions were used):</p> <p><i>Status:</i> E=Existing in the Point Source Permitting Regions; R=Revoked/Terminated - Pipe no longer permitted for discharge</p> <p><i>Discharge Type:</i> B=This individual NJPDES DSW permit is issued to those facilities that discharge treated and non-treated wastewater derived from, but not limited to process and non-process wastewater, contact and non-contact cooling water and storm water run-off. B4B=This general permit authorizes discharges of treated groundwater from petroleum leaks (i.e. fuel oil, diesel fuel, kerosene, aviation fuel, and gasoline) to select surface waterbodies.</p> <p>Source: NJDEP, Environmental Regulation, Division of Water Quality, June 14, 2017; NJDEP Bureau of Surface Water Permitting, January 6, 2016</p>					

An increase in impervious surface results in less water infiltrating to the soil and groundwater, which instead runs off the surface and gains velocity. As the velocity of water increases, the amount that can infiltrate into the soil and ground water is reduced and scouring and erosion increase. The stormwater eventually discharges into streams and rivers, carrying pollutants that it has picked up along the way (e.g. trash, used motor oil, sediments, fertilizers, pesticides, pet droppings, etc.). The transport of these pollutants into local water bodies can result in the destruction of fish, wildlife, and habitats; threats to public health due to contaminated food and drinking water supplies; and losses of recreational and aesthetic values. In addition, increased stormwater results in greater frequency and magnitude of floods (Hasse, and Lathrop, December 2016; Kaplan and Ayers, April 5, 2000).

Studies have shown that the level where impacts begin to be seen is above 10% impervious surfaces, and that impacts become severe over 25 to 30% (Kaplan and Ayers, April 5, 2000).

NJDEP determined approximate percent impervious surface based on particular land uses. Using the approximation calculated using the 2012 land use data, West Cape May is 21% impervious (161 acres) (see **Figure 6f**) (NJDEP, February 17, 2015).

The goals of New Jersey's Stormwater Management Rule (N.J.A.C. 7:8) include reducing runoff, flooding, erosion and non-point pollution for public safety as well as ecological and biological integrity. There are requirements for stormwater management measures and regional and municipal stormwater management planning (NJDEP, June 20, 2016).

The purpose of the Municipal Stormwater Regulation Program is to ensure a consistent approach to stormwater management statewide, reduce costs for regulated entities, and allow for a simple process for requesting authorization. All municipalities within the State are assigned either Tier A (more developed or coastal municipalities, including West Cape May) or Tier B (less developed and non-coastal) (NJDEP Bureau of Nonpoint Pollution Control, 2009).

The permits address stormwater quality related issues to new and existing development and redevelopment by requiring the preparation of a stormwater program and implementation of specific permit requirements referred to as Statewide Basic Requirements (SBRs). The Tier B Permit concentrates on new development and redevelopment projects and public education. The Tier A Permit has additional requirements aimed at controlling stormwater pollutants from existing development, such as public education, disposal of waste, solids and floatable controls, maintenance yard operations and employee training (NJDEP Bureau of Nonpoint Pollution Control, April 2004).





Figure 6f: West Cape May Percent Impervious Surface

Impervious surfaces are materials that prevent the infiltration of water into the soil such as parking lots, roads, buildings, sidewalks and compacted soil. Just over a fifth of West Cape May is impervious. Two point source pollution discharge sites are also located on the eastern border of the borough.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Chapter XXII Stormwater Control of the West Cape May Borough Code addresses stormwater management (see **Internet Resources**).

H. Surface Water Quality and Flow Monitoring

There are no routine NJDEP surface water quality monitoring sites within West Cape May. The NJDEP Bureau of Marine Water Monitoring collects samples from the coastal waters surrounding the Cape May Peninsula and analyzes them at its laboratory facilities at Leeds Point, NJ. Parameters such as salinity, dissolved oxygen, suspended solids and nutrients are measured to assess the ecological health of New Jersey's coastal waters See Internet Resources for a link to download the raw data for the Coastal Monitoring Network (BFBM, February 3, 2017).

Data collected by some sampling programs and from certain sites are input into EPA's national STORET²⁶ water quality database. These sites are indicated on **Figure 6g**.

I. Fish Consumption Advisories

When toxic pollutants are present in surface water, they are consumed by the organisms that live in the water. The process of *bioaccumulation* is when there is an increase in concentration of certain fat-soluble chemicals, such as DDT and PCBs, in successively higher trophic levels of a food chain or web. For example, insects living in contaminated sediments may have accumulated a certain amount of a toxin. Fish, by eating many of these insects, then ingest the toxin into their own bodies. Anything that eats that contaminated fish, including humans and other predators, will absorb the toxin. When the concentration of toxin becomes high enough, the individual's health will be impacted.

The NJDEP samples fish for certain toxic pollutants and, when necessary, issues state and regional *fish consumption advisories*, to reduce exposure to dioxin, PCBs and mercury. This information is intended to help individuals make an informed choice on the number of meals of fish to consume. The 2017 fish consumption advisories for fish caught anywhere in the state are listed in **Table 6.7**. See the **Internet References** for more information, such as fish preparation guidelines and annual updates.

²⁶ **STORET** is the U.S. Environmental Protection Agency (EPA) national water quality database for STORage and RETrieval (STORET). It includes physical, chemical, and biological data measured in waterbodies throughout the United States.

Legend

- West Cape May Borough
- Municipalities
- Canal
- Water
- Streams
- Storet stations*
- Tidal benchmarks
- ★ Head of Tide





Figure 6g: West Cape May Monitoring

Although there are a number of surface water quality monitoring stations in the vicinity of West Cape May, there are none located within the borough itself.



Data Sources: NJDEP
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

** STORET is the US EPA national water quality database for STORAge and RETrieval (STORET). It includes physical, chemical, and biological data measured in waterbodies throughout the United States.*

Table 6.7: 2017 Fish Consumption Advisories

LOCATION	SPECIES	ADVISORY/PROHIBITION(1)	
		General Population Range of Recommended Meal Frequency(2)	High-Risk Individuals (3)
		DO NOT EAT MORE THAN:	DO NOT EAT MORE THAN:
Delaware Estuary & Delaware Bay – Chesapeake & Delaware (C&D) Canal out to the mouth of Delaware Bay	Striped Bass	One meal per year	Do Not Eat
	White Perch	One meal per year	Do Not Eat
	White Catfish	One meal per year	Do Not Eat
	Channel Catfish	One meal per year	Do Not Eat
	America Eel	One meal per year	Do Not Eat
	Bluefish	Do Not Eat fish larger than 6lbs or 24 inches— One meal per year of fish less than 6lbs or less than 24 inches	Do Not Eat
	Weakfish	One meal per week	One meal per month
Atlantic Ocean – Sea Isle City to Cape May	Weakfish	One meal per week	One meal per month
Statewide Estuarine & Marine Waters – All coastal waterbodies except those under Waterbody Specific Advisories	Striped bass	One meal per month	Do Not Eat
	Bluefish (greater than 6 lbs/24 inches)	Six meals per year	Do Not Eat
	Bluefish (less than 6lbs/24 inches)	One meal per month	Do Not Eat
	American eel	Four meals per year	Do Not Eat
	American lobster	Do Not Eat the Green Gland (a.k.a., Tomalley or Hepatopancreas)	
Statewide Freshwater – All water bodies except those listed separately	Trout - (Brown, Brook Rainbow and Hybrid)	One meal per week	One meal per week
	Chain Pickerel	One meal per week	One meal per month
	Sunfish (bluegill, pumpkinseed and redbreast)	No restrictions	One meal per week
	Brown Bullhead	No restrictions	One meal per month
	Yellow Bullhead	No restrictions	One meal per month
	Common Carp	One meal per week	One meal per month
** Important Consumption Reminders **			
<p>(1) Eat only the fillet portions of the fish. Use proper trimming techniques to remove fat, and cooking methods that allow juices to drain from the fish (e.g., baking broiling, frying or grilling, and steaming). See text for full description.</p> <p>(2) One meal is defined as an eight-ounce serving.</p> <p>(3) High-Risk Individuals include infants, children, pregnant women, nursing mothers and women of childbearing age.</p>			
<p>Source: NJDEP Division of Science and Research, 2017 http://www.state.nj.us/dep/dsr/njmainfish.htm See interactive map for up to date fish advisories: http://njdep.maps.arcgis.com/apps/MapJournal/index.html?appid=922dff1885394cf19ccf1d9c8d52b4f0</p>			

References: Surface Water

A. Watersheds

NJDEP Land Use Management, Water Monitoring and Standards. April 4, 2011. N.J.A.C. 7:9B Surface Water Quality Standards. 115 pages. http://www.nj.gov/dep/rules/rules/njac7_9b.pdf

NJDEP Office of Environmental Planning. January 1997. Draft Statewide Watershed Management Framework Document for the State of New Jersey. <http://www.nj.gov/dep/watershedrestoration/docs/frame97fixed.pdf>

NJDEP, Geological Survey (NJGS). February 25, 2011. NJDEP 14 Digit Hydrologic Unit Code Delineations for New Jersey (DEPHUC14) (Version 20110225). GIS data. <http://www.state.nj.us/dep/gis/digidownload/zips/statewide/dephuc14.zip>

USGS. June 23, 2017. Hydrologic Unit Maps. <http://water.usgs.gov/GIS/huc.html> Accessed June 23, 2017.

Winter, T.C., J.W. Harvey, O.L. Franke, and W.M. Alley. 1998. "Natural Processes of Ground-Water and Surface-Water Interaction" in *Ground Water and Surface Water A Single Resource--USGS Circular 1139*. https://pubs.usgs.gov/circ/circ1139/htdocs/natural_processes_of_ground.htm. Accessed January 15, 2018.

B. Floodplains & Floods

Dinicola, Karen. 2005. The "100-Year Flood." US Geological Survey Fact Sheet 229-96. <http://pubs.usgs.gov/fs/FS-229-96/>

Federal Emergency Management Agency (FEMA). August 6, 2017. Community Status Book Report: New Jersey Communities Participating in the National Flood Program. <https://www.fema.gov/cis/NJ.html>. Accessed August 6, 2017.

Federal Emergency Management Agency (FEMA). July 25, 2017. FLOODSMART: The National Flood Insurance Program. http://www.floodsmart.gov/floodsmart/pages/flood_facts.jsp

Federal Emergency Management Agency (FEMA). June 29, 2017. National Flood Insurance Program: Flood Hazard Mapping. <https://www.fema.gov/national-flood-insurance-program-flood-hazard-mapping>. Accessed August 6, 2017.

Federal Emergency Management Agency (FEMA). March 7, 2017. Flood Zones. <https://www.fema.gov/flood-zones> (Definitions: <https://www.fema.gov/media-library-data/20130726-1535-20490-7429/appxd.pdf>). Accessed August 6, 2017.

Federal Emergency Management Agency (FEMA). October 4, 2017. FEMA's National Flood Hazard Layer (Official) (Cape May County, NJ (1:12,000)). GIS data: <https://msc.fema.gov>

Federal Emergency Management Agency (FEMA). June 21, 2007. Everyone Lives In A Flood Zone. Release Number: 1691/93-Floodsmart02. <https://www.fema.gov/news-release/2007/06/21/everyone-lives-flood-zone>

National Weather Service. October 2017a. Delaware Bay at Cape May. NOAA Advanced Hydrologic Prediction Service Hydrologic Prediction Service. <http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=cman4>. Accessed October 22, 2017.

National Weather Service. October 2017b. Cape May Harbor at Cape May Harbor. NOAA Advanced Hydrologic Prediction Service Hydrologic Prediction Service. <http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=capn4>. Accessed October 22, 2017.

West Cape May, NJ Borough Code. CHAPTER XXIII Flood Hazard Protection. <http://clerkshq.com/default.ashx?clientsite=westcapemay-nj>

C. Wetlands

Fair, Abigail. 2004. Freshwater Wetlands Protection in New Jersey: A Manual for Local Officials. Third Edition. Association of New Jersey Environmental Commissions. 52 pages. <http://www.anjec.org/WaterFreshwaterWetlands.htm>

NJDEP Division of Land Use Management. July 16, 1998. Freshwater Wetlands Protection Act N.J.S.A. 13:9B (last amended July 1998) 22 pages. http://www.nj.gov/dep/landuse/download/13_9b.pdf

NJDEP Division of Land Use Management. July 17, 2017. Freshwater Wetlands Protection Act Rules N.J.A.C. 7:7A (last amended July 17, 2017). 239 pages. http://www.nj.gov/dep/rules/rules/njac7_7a.pdf

NJDEP Land Use Regulation Program. August 16, 2017. Freshwater Wetlands Program Home Page. http://www.nj.gov/dep/landuse/fww/fww_main.html. Accessed October 22, 2017.

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice. 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

D. Surface Water Quality Standards

NJDEP Land Use Management, Water Monitoring and Standards. April 4, 2011. N.J.A.C. 7:9B Surface Water Quality Standards. 113 pages. Last Amended – April 4, 2011. http://www.nj.gov/dep/rules/rules/njac7_9b.pdf

NJDEP Water Monitoring and Standards. June 15, 2017. Category One Waters. <http://www.nj.gov/dep/wms/bears/c1waters.htm>

NJDEP, Bureau of Freshwater and Biological Monitoring (BFBM). December 2010. NJDEP Surface Water Quality Standards of New Jersey (Version 201012). GIS data. <http://www.state.nj.us/dep/gis/stateshp.html#SWQS>

E. Integrated List and Total Maximum Daily Loads (TMDLs)

NJDEP Division of Water Monitoring and Standards, Bureau of Environmental Analysis, Restoration and Standards (BEARS). May 2017 (final). 2014 New Jersey Integrated Water Quality Assessment Report. http://www.nj.gov/dep/wms/bears/docs/2014_final_integrated_report.pdf

NJDEP Division of Water Monitoring and Standards, Bureau of Environmental Analysis, Restoration and Standards (BEARS). March 9, 2017. Total Maximum Daily Loads home page. <http://www.nj.gov/dep/wms/bears/tmdl.html>. Accessed June 23, 2017.

NJDEP Division of Water Monitoring and Standards. September 27, 2017. Integrated List of Waters for New Jersey, 2014 (Integrated List). GIS Data. <http://njogis-newjersey.opendata.arcgis.com/datasets/2014-new-jersey-integrated-list-of-waters-integrated-list?geometry=-82.144,38.633,-61.237,41.573>

NJDEP Division of Water Monitoring and Standards, Bureau of Environmental Analysis, Restoration and Standards (BEARS). February 2015. 2014 Integrated Water Quality Monitoring and Assessment Methods. http://www.nj.gov/dep/wms/bears/docs/2014_final_methods_document_and_response_to_comments.pdf

NJDEP and Tetra Tech, Inc. September 27, 2006. Ten Total Maximum Daily Loads for Total Coliform to Address Shellfish-Impaired Waters in Watershed Management Area 16 Atlantic Coastal Water Region. http://www.nj.gov/dep/wms/bears/docs/Coastal_Pathogen_TMDLs_WMA16.pdf

USEPA. 2017. Clean Water Act Section 303(d): Impaired Waters and Total Maximum Daily Loads (TMDLs). <https://www.epa.gov/tmdl>. Accessed June 23, 2017.

US. January 4, 2011. Federal Water Pollution Control Act Title 33—Navigation And Navigable Waters Chapter 26 Water Pollution Prevention And Control [As Amended Through Pub.L. 111-378, January 4, 2011]† (33 U.S.C. § 1251 et seq.) 132 pages. http://www.waterboards.ca.gov/laws_regulations/docs/fedwaterpollutioncontrolact.pdf

F. Point Source Pollution

NJDEP Bureau of Surface Water Permitting. January 6, 2016. Surface Water Permitting Information. <http://www.nj.gov/dep/dwq/sw.htm>. Accessed October 22, 2017

NJDEP, Environmental Regulation, Division of Water Quality. June 14, 2017. NJPDES Surface Water Discharges in New Jersey, (1:12,000) Edition 20170614. GIS data. http://njogis-newjersey.opendata.arcgis.com/datasets/55e8b1a25706431490a8f3becf861f91_5

NJDEP Land Use Management, Water Monitoring and Standards. April 4, 2011. N.J.A.C. 7:9B Surface Water Quality Standards. Last Amended – April 4, 2011. 115 pages. http://www.nj.gov/dep/rules/rules/njac7_9b.pdf

G. Non-point Source Pollution

Hasse, John and Richard G. Lathrop. December 2016. Changing Landscapes in the Garden State: Land Use Change in New Jersey, 1986-2012. <http://www.crssa.rutgers.edu/projects/lc/urbangrowth/index.html>

Kaplan, Marjorie and Mark Ayers. April 5, 2000. Impervious Surface Cover Concepts and Thresholds. NJDEP and USGS. 10 pages. Cached copy: http://www.google.com/url?sa=t&rct=j&q=&esrc=s&frm=1&source=web&cd=3&cad=rja&ved=0CDoQFjAC&url=http%3A%2F%2Fmss3.libraries.rutgers.edu%2Fdlr%2Foutputs.php%3Fpid%3Drutgers-lib%3A37001%26ds%3DPDF-1&ei=BOOr7UdjbB_e34APUhoCQCg&usq=AFOjCNEcfJXaTLyII4pfltG4GedUWvTgma&sig2=hhz-DuRu2u8jFmIheg3btg

NJDEP Bureau of Nonpoint Pollution Control. April 2004. Tier A Municipal Stormwater Guidance Document. 143 pages. http://www.state.nj.us/dep/dwq/tier_a_guidance.htm

NJDEP Bureau of Nonpoint Pollution Control. 2009. Municipal Tier Assignments 2009 Under the NJPDES Municipal Stormwater Regulation Program. <http://www.nj.gov/dep/dwq/images/mun-tierA-assignments1.jpg>

NJDEP. June 20, 2016. N.J.A.C. 7:8 Stormwater Management Rule. 39 pages. Date last amended: June 20, 2016. http://www.nj.gov/dep/rules/rules/njac7_8.pdf

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice. 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

West Cape May, NJ Borough Code. CHAPTER XXII Stormwater Control. <http://clerkshq.com/default.ashx?clientsite=westcapemay-nj>

H. Surface Water and Flow Monitoring

NJDEP Bureau of Marine Water Monitoring. February 3, 2017. Basic Water Quality Measurements. <http://www.nj.gov/dep/bmw/delaware2.htm>. Accessed October 23, 2017.

I. Fish Consumption Advisories

NJDEP Division of Science and Research. 2017. Fish Smart, Eat Smart: A Guide to Health Advisories for Eating Fish and Crabs Caught in New Jersey Waters. <http://njdep.maps.arcgis.com/apps/MapJournal/index.html?appid=922dff1885394cf19ccf1d9c8d52b4f0>

Internet Resources: Surface Water

General Water Resources Protection

Natural Processes of Ground-Water and Surface-Water Interaction (USGS): https://pubs.usgs.gov/circ/circ1139/htdocs/natural_processes_of_ground.htm

SEEDS: The NJ Environmental Education Directory Website: <http://www.state.nj.us/dep/seeds/index.html>

Basic Watershed Information (Watershed Restoration Section): <http://www.nj.gov/dep/watershedrestoration/info.html>

The Clean Water Book: Choices for Watershed Protection:
http://www.nj.gov/dep/watershedrestoration/waterbook_tble.html

NJDEP Laws & Rules: <http://www.nj.gov/dep/landuse/lawsregs.html>

Water Quality Fact Sheets and Bulletins (NJ Agricultural Experiment Station Rutgers Cooperative Research & Extension): <http://njaes.rutgers.edu/pubs/subcategory.asp?cat=6&sub=50&order=LastRevised>

Floodplains & Floods

FEMA Flood Map Service Center: <http://msc.fema.gov/portal>

Flood Hazard Area Program (NJDEP Land Use Regulation) http://www.nj.gov/dep/landuse/fha_main.html

FloodSmart: The Official Site of the National Flood Insurance Program: <http://www.floodsmart.gov>

National Weather Service Advanced (NOAA) Hydrologic Prediction Service (flood predictions):

Delaware Bay at Cape May: <http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=cman4>

Cape May Harbor at Cape May Harbor:

<http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=capn4>

West Cape May, NJ Borough Code. CHAPTER XXIII Flood Hazard Protection.

<http://clerkshq.com/default.ashx?clientsite=westcapemay-nj>

Wetlands

Freshwater Wetlands Program (NJDEP Land Use Regulation): http://www.nj.gov/dep/landuse/fww/fww_main.html

Freshwater Wetlands Program: Before You Buy – Before You Build: <http://www.nj.gov/dep/landuse/bybob.html>

SWQS

Category One Waters: <http://www.nj.gov/dep/wms/bears/c1waters.htm>

Integrated List & TMDL

NJDEP Integrated WQ monitoring and Assessment Report: <http://www.nj.gov/dep/wms/bears/generalinfo.htm>

NJDEP Total Maximum Daily Load (TMDL): <http://www.nj.gov/dep/wms/bears/tmdls.html>

USEPA Laws and Regulations: <http://www2.epa.gov/laws-regulations>

Point Source Pollution

NJPDES Permitting: <http://www.nj.gov/dep/dwq/database.htm>

Non-Point Source Pollution / Stormwater:

NJDEP's Stormwater Website (includes links to all of the following, and more): <http://www.njstormwater.org/> or <http://www.nj.gov/dep/dwq/fd.htm>

NJDEP Municipal Stormwater Regulation Program: http://www.state.nj.us/dep/dwq/msrp_home.htm

Stormwater Best Management Practices Manual: http://www.njstormwater.org/bmp_manual2.htm

Clean Water NJ: <http://www.cleanwaternj.org/index.htm>

Multimedia Resources: <http://www.cleanwaternj.org/multimedia.html>

Green Infrastructure: <http://www.nj.gov/dep/gi/>

USEPA Nonpoint Source Pollution: <http://water.epa.gov/polwaste/nps/index.cfm>

Surface Water Quality and Flow

Coastal water quality data: <http://www.nj.gov/dep/bmw/delaware2.htm>

National Weather Service Advanced (NOAA) Hydrologic Prediction Service (flood predictions):

Delaware Bay at Cape May: <http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=cman4>

Cape May Harbor at Cape May Harbor:

<http://water.weather.gov/ahps2/hydrograph.php?wfo=phi&gage=capn4>

NJDEP Drought Information: <http://www.njdrought.org/>

USEPA STORET Database: <http://www.epa.gov/storet>

USGS Real-time flow data index of NJ sites: <http://waterdata.usgs.gov/nj/nwis/current/?type=flow>

Water Quality Data Portal: <https://www.waterqualitydata.us/>

Fish Advisories & Guides

NJ Division of Science & Research Fish Advisories Home Page: <http://www.state.nj.us/dep/dsr/njmainfish.htm>

Fish Smart Eat Smart: <http://www.state.nj.us/dep/dsr/fishsmart.pdf>

NJDEP Regulations:

NJDEP Rules & Regulations, current and proposed: <http://www.state.nj.us/dep/rules>

Phone Contacts:

NJ Drought Hotline: 1-800-4-ITS DRY (1-800-448-7379) or <http://www.njdrought.org/>

NJ Environmental Incident Hotline (hazardous spill, fire, explosion, illegal dumping, wildlife problem):
1-877-WARNDEP / 1-877-927-6337 (toll-free, 24 hours) or <http://www.nj.gov/dep/warndep.htm>

NJDEP Bureau of Coastal & Land Use Compliance & Enforcement: 1-609-292-1240

NJDEP Division of Land Use Regulation (Wetlands, Streams/Rivers, Flood Hazard Areas):

Technical Support Center: (609) 777-0454 or <http://www.nj.gov/dep/landuse/contact.html>

Forms: <http://www.nj.gov/dep/landuse/forms.html>

7: BIOLOGICAL RESOURCES

A. Dominant Vegetation (Land Cover)

The New Jersey Comparative Risk Project (March 2003) listed habitat fragmentation and habitat loss as the highest ranking stressors of Statewide ecological quality. Certain species that require large expanses of intact habitat are becoming less common. Other factors that impact ecological health include exotic species and exotic diseases, overpopulations of deer and geese, and pollution.



The 2012 Land Use/Land Cover (LU/LC) data layer was created by a consultant to NJDEP by comparing the 2007 LU/LC layer to 2012 color infrared imagery (2007 and 2012 aerial photos are shown in **Figure 1d** and **Figure 1e**, respectively) and delineating and coding areas of change with a 1 foot pixel resolution. The classification system used was a modified Anderson Classification System (USGS, 2010) that provided the parameters for proper and consistent coding of the LU/LC feature classes and subclasses. It should be noted that 1) changes since 2012 are not shown, and 2) the method is not 100% accurate. In addition, since it is based on interpretation of aerial photographs, it cannot provide information about the particular species found in an area. The land cover classifications are shown in **Figures 7a, 7b, and 7c**. The relative proportion of each within West Cape May is provided in the acreage figures in **Table 7.1** (NJDEP, 2010).

The largest portion of land in West Cape May is classified as Urban (43.72%). Another 13.88% is utilized for agriculture, while 5.61% of the borough's wetlands have been modified for agricultural, residential or other purposes. Together, these categories account for about 63% of the borough. The remaining 37% of the land is primarily either unmodified wetlands (29.70%) or forested land (5.63%). Open water and barren land each account for less than one percent of the total land use (NJDEP, 2010).

Roughly seventeen percent of the borough is classified as either a wetland (13.57%) or a forest (3.76%) land use type that is dominated by trees. In recognition of the many benefits of trees, the Borough of West Cape May has adopted two ordinances (494-2015 and 515-2016) for the preservation and protection of trees that add special value to the community because of their size, rarity, historical, environmental or aesthetic importance (Borough of West Cape May, Chapter XXX).

Table 7.1: Land Use/Land Cover (Anderson Classification) in West Cape May

Land Cover Code	Land Cover Name	Acres*	Percent of Borough
URBAN LAND USE TYPE			
1110	RESIDENTIAL, HIGH DENSITY OR MULTIPLE DWELLING	34.23	4.54%
1120	RESIDENTIAL, SINGLE UNIT, MEDIUM DENSITY	185.95	24.64%
1130	RESIDENTIAL, SINGLE UNIT, LOW DENSITY	36.49	4.84%

Land Cover Code	Land Cover Name	Acres*	Percent of Borough
1140	RESIDENTIAL, RURAL, SINGLE UNIT	10.23	1.36%
1200	COMMERCIAL/SERVICES	26.83	3.56%
1400	TRANSPORTATION/COMMUNICATION/UTILITIES	3.77	0.50%
1420	RAILROADS	0.88	0.12%
1600	MIXED URBAN OR BUILT-UP LAND	1.82	0.24%
1700	OTHER URBAN OR BUILT-UP LAND	10.42	1.38%
1800	RECREATIONAL LAND	16.57	2.20%
1804	ATHLETIC FIELDS (SCHOOLS)	2.72	0.36%
	Total Acres of URBAN:	329.91	43.72%
AGRICULTURE LAND USE TYPE			
2100	CROPLAND AND PASTURELAND	77.15	10.22%
2200	ORCHARDS/VINEYARDS/NURSERIES/HORTICULTURAL AREAS	12.9	1.71%
2400	OTHER AGRICULTURE	14.7	1.95%
	Total Acres of AGRICULTURE:	104.75	13.88%
FOREST LAND USE TYPE			
4110	DECIDUOUS FOREST (10-50% CROWN CLOSURE)	5.64	0.75%
4120	DECIDUOUS FOREST (>50% CROWN CLOSURE)	14.36	1.90%
4210	CONIFEROUS FOREST (10-50% CROWN CLOSURE)	4.49	0.59%
4220	CONIFEROUS FOREST (>50% CROWN CLOSURE)	0.01	0.00%
4312	MIXED FOREST (>50% CONIFEROUS WITH >50% CROWN CLOSURE)	2.05	0.27%
4322	MIXED FOREST (>50% DECIDUOUS WITH >50% CROWN CLOSURE)	1.88	0.25%
4410	OLD FIELD (< 25% BRUSH COVERED)	2.89	0.38%
4420	DECIDUOUS BRUSH/SHRUBLAND	9.51	1.26%
4440	MIXED DECIDUOUS/CONIFEROUS BRUSH/SHRUBLAND	1.64	0.22%
	Total Acres of FOREST:	42.47	5.63%
WATER LAND USE TYPE			
5300	ARTIFICIAL LAKES	2.92	0.39%
5410	TIDAL RIVERS, INLAND BAYS, AND OTHER TIDAL WATERS	3.09	0.41%
	Total Acres of WATER:	6.01	0.80%
WETLANDS LAND USE TYPE			
1750	MANAGED WETLAND IN MAINTAINED LAWN GREENSPACE	1.23	0.16%
2140	AGRICULTURAL WETLANDS (MODIFIED)	39.02	5.17%
2150	FORMER AGRICULTURAL WETLAND (BECOMING SHRUBBY, NOT BUILT-UP)	2.99	0.40%
6111	SALINE MARSH (LOW MARSH)	27.44	3.64%
6112	SALINE MARSH (HIGH MARSH)	21.54	2.85%
6141	PHRAGMITES DOMINATE COASTAL WETLANDS	27.91	3.70%
6210	DECIDUOUS WOODED WETLANDS	90.89	12.04%
6231	DECIDUOUS SCRUB/SHRUB WETLANDS	13.46	1.78%
6232	CONIFEROUS SCRUB/SHRUB WETLANDS	1.89	0.25%
6233	MIXED SCRUB/SHRUB WETLANDS (DECIDUOUS DOM.)	7.2	0.95%
6234	MIXED SCRUB/SHRUB WETLANDS (CONIFEROUS DOM.)	0.75	0.10%

Land Cover Code	Land Cover Name	Acres*	Percent of Borough
6240	HERBACEOUS WETLANDS	0.72	0.10%
6241	PHRAGMITES DOMINATE INTERIOR WETLANDS	17.72	2.35%
6251	MIXED WOODED WETLANDS (DECIDUOUS DOM.)	11.53	1.53%
7430	DISTURBED WETLANDS (MODIFIED)	1.32	0.17%
7440	DISTURBED TIDAL WETLANDS	0.82	0.11%
Total Acres of WETLANDS:		266.43	35.31%
BARREN LAND USE TYPE			
7500	TRANSITIONAL AREAS	5.06	0.67%
Total Acres of BARREN LAND:		5.06	0.67%
West Cape May Borough Totals:		754.63	100.00%
* Acreage from the GIS data may vary from acreage calculated based on tax maps.			
Source: NJDEP, 2010; USGS, 2010.			



B. Wildfire Fuel Hazard

The New Jersey Forest Fire Service (NJFFS), a division of NJDEP, assessed *Wildfire Fuel Hazard* (WFH) throughout New Jersey (see **Figure 7d** and **Table 7.2**). The purpose is to provide information for NJ Forest Fire Service personnel, government agencies, and others interested in assessing the risk of wildfires throughout New Jersey. Modified Anderson Land Use/Land Cover Classifications from the 2002 Land Use/Land Cover dataset were assigned Wildfire Fuel Hazard Rankings (0 = Water, 1 = Low, 2 = Moderate, 3 = High, 4 = Very High, 5 = Extreme, 6 = Urban, 7 = Agriculture, 8 = Barren Land). Areas with 30% or greater slope and Wildfire Fuel Hazard 1 to 4 were increased by 1 (e.g. Low became Moderate, etc.) (NJDEP, 2009).

The majority of West Cape May is either not rated or rated Low Wildfire Fuel Hazard. The highest wildfire risk areas in the borough are phragmites stands and high salt marsh.

Table 7.2: Wildfire Fuel Hazard

Fuel Hazard	Fire Description	Acres	Percent
0	water	4.6	1%
1	low	189.7	25%
2	moderate	61.1	8%
3	high	52.8	7%
4	very high	36.9	5%
5	extreme	3.9	1%
6	urban	303.7	40%
7	agriculture	94.2	12%
8	barren land	7.8	1%
Total		754.6	100%



Prepared by Kratzer Environmental Services

Data Sources: NJDEP, NJDOT


Note: Map accuracy is limited to the accuracy and scale of the original data sets.

Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Figure 7a: West Cape May

Land Cover 2012: Wetlands Type

Wetlands account for approximately 35% of the land cover in West Cape May Borough.




Legend

- West Cape May
- Parcels
- Water
- Roads

Land Cover 2012: Agriculture and Forest


- | | |
|--|---|
| 2100 Cropland and Pastureland | 4110 Deciduous Forest (10-50% closure) |
| 2200 Orchards/Vineyards/nurseries/Horticulture | 4120 Deciduous Forest (>50% crown closure) |
| 2400 Other Agriculture | 4210 Coniferous Forest (10-50% crown closure) |
| | 4312 Mixed Forest (>50% con. >50% closure) |
| | 4322 Mixed Forest (>50% dec. >50% closure) |
| | 4410 Old Field (< 25% brush covered) |
| | 4420 Deciduous Brush/Shrubland |
| | 4440 Mixed Dec./Con. Brush/Shrubland |



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

**Figure 7b: West Cape May
 Land Cover 2012: Agriculture & Forest**

Agricultural lands comprise nearly 14% of the area in West Cape May, and just under 6% of the borough is forested.




Legend

- West Cape May
- Parcels
- Water
- Roads

Land Cover (2012) Barren and Urban Types

- 1110 Residential, High Density or Multiple Dwelling
- 1120 Residential, Single Unit, Medium Density
- 1130 Residential, Single Unit, Low Density
- 1140 Residential, Rural, Single Unit
- 1200 Commercial/Services
- 1400 Transportation/Communication/Utilities
- 1420 Railroads
- 1600 Mixed Urban or Built-up Land
- 1700 Other Urban or Built-up Land
- 1800 Recreational Land
- 1804 Athletic Fields (schools)
- 7500 Transitional Areas (Barren Land)



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Prepared by Kratzer Environmental Services

**Figure 7c: West Cape May
 Land Cover 2012: Barren & Urban**

Roughly 44% of West Cape May is dedicated to urban uses. Less than 1% of the land in the borough is classified as barren.





Figure 7d: West Cape May Wildfire Fuel Hazard

The majority of West Cape May is Low Wildfire Fuel Hazard. Most of the areas ranked as Very High or Extreme are either high salt marsh habitat or phragmites coastal wetlands.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

C. Wildlife

New Jersey hosts 323 bird species, 89 mammal species, 44 reptile, 34 amphibian, 90 freshwater fish and over 300 marine finfish species (NJDEP 2004a, 2004b, 2014, 2016). This high diversity in such a small state is partly due to New Jersey's geographic position where northern ecosystems reach their southern limit and where southern ecosystems reach their northern limit. In addition, the state provides a wide variety of habitats including mountains, valleys, rolling hills, wetlands, pinelands, beaches, estuaries and rivers (NJDEP, January 19, 2012). The NJDEP website offers checklists for the birds, mammals, reptiles and amphibians of New Jersey; with notes on the status of each (e.g. common or rare) (see **Internet Resources**). A variety of plant and animal species enjoy West Cape May's diverse habitat types.

Birds

Southern Cape May County has long been known as the premier birding site in New Jersey. Its position between the Atlantic Ocean and the Delaware Bay make it a significant stopover for migratory shorebirds, songbirds and raptors, and its diverse habitat components offer suitable sites for many year-round resident species as well (USFWS, 1997). Cape Island, defined as the region south of the canal and west of Cape Island Creek, is listed as an Important Bird Area of New Jersey (see **Figure 7e**), and the small woodlots and fallow fields of West Cape May are specifically mentioned as part of the local habitat matrix that makes this region special (Frank, 2010).

To date, 427 species of birds have been documented in Cape May County (Sullivan et al., 2009), but the list may be revised with additional sightings and surveys (see **Appendix C.1**). Roughly a quarter of those species are only rare visitors to New Jersey, and are not included on the NJDEP list of state birds. Endangered, threatened and special concern birds are discussed in **Section 7E**, and non-indigenous birds in **Section 7G**. The Canada goose, a native species which may disproportionately affect the environment, is discussed below.

Canada Goose

The Canada goose (*Branta canadensis*) is one of New Jersey's most easily recognized birds, with its black head and neck, white cheek patch and undersides, brown back and large size (2'-3' tall, 10-12 lbs.). There are two distinct populations in NJ, migratory geese that visit the state in the winter and non-migratory geese, that nest in the state. Sources estimate the NJ population of resident Canada geese at approximately 83,000 to 96,800 (USDA, January 2003; NJDEP, March 2001).

While many people enjoy the sight of a few geese, this high population of non-migrating geese can cause the following problems:

- overgrazing of lawns and athletic fields, which impacts aesthetics and causes erosion
- damage to cropland, increasing erosion hazard and crop losses
- accumulations of feces on land, creating a health risk from disease-causing organisms
- degradation of water quality, from fecal bacteria, nitrogen and phosphorous
- hazards to aircraft at airports



- aggression and attacks on humans
- noise (USDA, January 2003; NJDEP, March 2001).

As migratory game species, Canada geese are afforded federal and state protection. Therefore, any management techniques involving handling nests, eggs or birds require a permit (NJDEP Division of Fish and Wildlife, No date).

Mammals

Twenty-eight of the 89 mammals listed on the NJDEP checklist are restricted to marine environments, leaving 61 species that may be observed at various locations around the state (**Appendix C.2**). Some of the listed mammals are limited to specific regions within the state, while others enjoy a wide range. A mammal list is not available specifically for either Cape May County or West Cape May Borough. Endangered mammals are discussed in **Section 7E**, and non-indigenous mammals in **Section 7G**. One large native mammal that frequently clashes with the human population is discussed below.

White-tailed Deer

The white-tailed deer (*Odocoileus virginianus*), the largest herbivore living wild in New Jersey, is seen throughout all but the most urbanized areas of the state. Although the deer is a large animal, individuals tend to stay in a one square mile or less home range, one of the smallest ranges among wild ruminants (Burnett, 2004).

Biologists have estimated that before the arrival of European settlers, there were about 8-11 white-tailed deer per square mile. By the early 1900's, New Jersey's deer herd was reduced to a handful by unregulated hunting. However, efforts to protect the deer herd were so successful that deer were considered over-populous by the 1920's (Latham et al, 2005). In addition, deer have been able to adapt to human-altered habitats. Studies have shown that deer densities of over 10-15 per square mile have negative impacts on the diversity of understory vegetation and on the native songbird and wildflower populations that depend on a diverse understory, while deer populations in excess of 20 per square mile prevent tree regeneration (Latham et al, 2005).

Where deer are overabundant, this results in excessive damage to agricultural crops, gardens and residential landscaping; an increased incidence of deer/vehicle collisions; prevention of forest regeneration (which impacts plants and animals dependent on the forest); and the potential for reduced deer health due to inadequate nutrition and the spread of disease (Honachefsky, 2000; Latham et al, 2005; Sauer, 1998). Despite these impacts, deer remain a natural part of the ecosystem, and are not solely responsible for diversity loss and habitat degradation.



Documentation of deer population numbers is not available for Cape May County or West Cape May Borough, therefore it is unknown whether the population exceeds either the number that can be sustained over an extended period (*ecological carrying capacity*) or the number that can coexist compatibly with local human populations (*cultural carrying capacity*) (NJDEP, 1999).

The state is divided into 70 Deer Management Zones (DMZs), with differing deer hunting regulations applied to different DMZs. West Cape May Borough falls within DMZ 34, which includes much of Cape May County and portions of Cumberland County and utilizes Regulation Set 0. Public lands open to deer hunting in Zone 34 include Beaver Swamp

Wildlife Management Area (WMA), Belleplain State Forest, Cape May National Wildlife Refuge, Cape May Wetlands WMA, Dennis Creek WMA, Heislerville WMA, Higbee Beach WMA, Lizard Tail Swamp Preserve, and Tuckahoe WMA. Data from the 2015-16 season indicate that 367 deer were harvested in Zone 34 (NJDEP Division of Fish and Wildlife, March 1, 2017).

Reptiles and Amphibians

Of the 39 non-marine reptile species recorded in New Jersey, 29 have ranges which include Cape May County (**Appendix C.3**). Similarly, there are 34 species of amphibians in the state, 21 of which range into Cape May County (**Appendix C.4**). Although lists of reptiles and amphibians (collectively known as herptiles) are not maintained by the county, their potential presence in the Cape May area can be extrapolated from the species range maps provided by the state's Division of Fish and Wildlife (NJDEP, Division of Fish & Wildlife, June 6, 2016). Endangered, threatened and special concern herptiles are discussed in **Section 7E**, and the non-indigenous species are covered in **Section 7G** (USGS January 26, 2017).

Wildlife of Vernal Pools

Vernal pools are defined as confined depressions, either natural or man-made, that maintain ponded water for part of the year, have no permanent outflow, and are devoid of breeding fish populations. These temporary wetlands provide habitat to many species of amphibians, several of which breed exclusively in vernal pools, as well as a multitude of insects, reptiles, plants, and other wildlife. Certification of a vernal pool may be achieved by documenting breeding activity of obligate vernal pool species (such as wood frogs or spotted salamanders; see **Table 7.3**) or by documenting both the presence of facultative species and photographic evidence that the pool goes dry or demonstrating the absence of fish (Tesauro, no date).

Table 7.3: Obligate And Facultative Fauna Species Found In Vernal Habitats

Obligate Vernal Pool Breeding Species	Facultative Vernal Pool Breeding Amphibians	Reptiles that Inhabit Vernal Pools on a Seasonal Basis
Eastern tiger salamander <i>ENDANGERED</i>	Green frog	Wood turtle <i>THREATENED</i>
Marbled salamander <i>Special Concern</i>	Bullfrog	Spotted turtle <i>Special Concern</i>
Spotted salamander	Pickerel frog	Mud turtle
Jefferson salamander <i>Special Concern</i>	Southern leopard frog	Eastern painted turtle
Blue-spotted salamander <i>ENDANGERED</i>	Carpenter frog <i>Special Concern</i>	Common snapping turtle
Jefferson x Blue-spotted salamander <i>ENDANGERED</i>	Spring peeper	
Wood frog	Eastern cricket frog	(These reptiles visit vernal pools primarily to eat the eggs and larvae of amphibians.)
Eastern spadefoot toad	New Jersey chorus frog	
Fairy shrimp (order Anostraca)	Upland chorus frog	
	Northern gray treefrog	
	Cope's gray treefrog <i>ENDANGERED</i>	
	Pine barrens treefrog <i>ENDANGERED</i>	
	Four-toed salamander	
	Long-tailed salamander <i>THREATENED</i>	
Note: Species in black are either known to occur in West Cape May Borough or their ranges include Cape May County; species in gray have ranges that do not include Cape May County, therefore it would be unlikely to find them in West Cape May.		
Sources: Kenney et al, no date; Gessner and Stiles, February 2001; N.J.A.C 7:7A, Appendix 1.		

There are currently no documented vernal pool habitats within West Cape May. One site just east of the Borough has been identified as a potential vernal pool, and a portion of the associated habitat extends into West Cape May (see **Figure 7f**). Furthermore, vernal pools have




Legend

- West Cape May
- Municipalities
- Roads

Important Bird Areas: Priority

- Continental
- Global
- State



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.


Figure 7e: West Cape May & Vicinity Important Bird Areas (Audubon)

Cape Island is listed as a Globally Important Bird Area. Portions of West Cape May which are part of the habitat matrix that makes this region special include Cape May Meadows, the Cape Island Creek area, and the small woodlots and fallow fields in the western part of the borough.



Figure 7f: West Cape May Landscape Project version 3.3

The Landscape Project indicates that 91.6% of the borough is part of a habitat patch known to support endangered animal species. Smaller patches of land indicate the presence of suitable habitat for additional rare species. Two vernal pools and a portion of vernal habitat associated with an offsite pond are also found in the borough.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized

been reported within a complex of permanent and temporary ponds in the center of the borough that are utilized by breeding frogs (Remington, Vernick & Walberg, 2003).

Fish

The New Jersey Division of Fish and Wildlife (2016) currently reports a total of 90 freshwater fish species in the state (**Appendix C.5**), although one of those (the longnose gar) is considered extirpated. Roughly two-thirds of those species are native to the state, while the others have been introduced either accidentally or deliberately. Some species introduced as game fish have become naturalized, while others do not readily reproduce and are repeatedly stocked for recreational purposes (NJ Division of Fish & Wildlife, 2016). Endangered fish species are discussed in **Section 7E**, and non-indigenous fish in **Section 7G**.

The location of West Cape May Borough does not lend itself to recreational freshwater fishing. In fact, no streams or rivers are listed as public fishing sites anywhere within Cape May County, and public fishing in ponds or lakes is only available north of the canal (NJDEP, July 22, 2016). Similarly, no trout stocked waters are located in the immediate vicinity of West Cape May (NJDEP Division of Fish and Wildlife, June 4, 2013).

D. Endangered, Threatened and Special Concern Species

The health of an area's animal and plant populations can be an indicator of the health and sustainability of the environment for people. The decline or disappearance of one (or more) species may signal the deterioration of the habitat. Other species, and human health and welfare, may soon follow. Preserving the future of endangered and threatened species helps preserve our own species, benefiting human health and quality of life by protecting watersheds, preserving land in its natural state, and restoring wildlife habitat. Many people also place an intrinsic value on all species (Conserve Wildlife Foundation, 2002).

Many species are naturally rare in parts of their range, especially at the periphery. New Jersey often lies at the southern periphery of the range for many "northern" species and at the northern edge of the range of many "southern" species. Therefore, a species considered rare or imperiled within the state of New Jersey is not necessarily in danger of extinction worldwide. In addition, many rare species depend on large tracts of continuous undisturbed habitat to survive. If these habitats are interrupted by developed areas, the patches may become too small to support certain species.

The NJ Endangered Species Conservation Act was signed into law on December 14, 1973 (N.J.S.A. 23:2A-1 - 15), preceding the federal Endangered Species Act by two weeks. This milestone legislation established laws to protect and restore the state's endangered and threatened wildlife whose survival in New Jersey is imperiled by loss of habitat, over-exploitation, pollution, or other impacts (NJDEP, October 6, 2004). In February 2012, NJDEP updated the Endangered and Nongame Species rules (N.J.A.C. 7:25), revising the species list based on science, upgrading the status of some recovering species and adding some declining species to the list (NJDEP Division of Fish and Wildlife, April 2, 2012 and January 18, 2011).

Table 7.4 presents the definitions used by NJDEP in describing the status of species. In order to better document the status or change in status of species, NJDEP solicits information from the general public concerning sightings of endangered, threatened and special concern species. People should use the appropriate reporting forms (see **Internet Resources** and **Appendix D.1 and D.2**).

Table 7.4: Definitions of Species Status

STATE STATUS	STATE STATUS DEFINITIONS
<p>Animals: Two animal lists provide state status codes after the Endangered and Nongame Species Conservation Act of 1973 (N.J.S.A. 23:2A-13 et. seq.): the list of endangered species (N.J.A.C. 7:25-4.13) and the list defining status of indigenous, nongame wildlife species of New Jersey (N.J.A.C. 7:25-4.17(a)). The status of animal species is determined by the Endangered and Nongame Species Program (ENSP), with the review and approval of the Endangered and Nongame Species Advisory Committee. Status for animals separated by a slash(/) indicate a dual status. First status refers to the state breeding population, and the second status refers to the migratory or winter population.</p>	
E	An endangered species is one whose prospects for survival within the state are in immediate danger due to one or many factors - a loss of habitat, over exploitation, predation, competition, disease. An endangered species requires immediate assistance or extinction will probably follow.
T	A threatened species is a species that may become endangered if conditions surrounding the species begin to or continue to deteriorate.
SC	The term Special Concern applies to animal species that warrant special attention because of some evidence of decline, inherent vulnerability to environmental deterioration, or habitat modification that would result in their becoming a Threatened species. This category would also be applied to species that meet the foregoing criteria and for which there is little understanding of their current population status in the state.
S	A stable species is one whose population is not undergoing any long-term increase/decrease within its natural cycle.
U	An undetermined species is one about which there is not enough information available to determine the status.
<p>Plants: Plant taxa listed as endangered are from New Jersey's official Endangered Plant Species List (N.J.A.C. 7:5C – 5.1).</p>	
E	Native New Jersey plant species whose survival in the State or nation is in jeopardy.
FEDERAL STATUS	FEDERAL STATUS DEFINITIONS
LE	Taxa formally listed as endangered .
LT	Taxa formally listed as threatened .
REGIONAL STATUS	REGIONAL STATUS CODES FOR PLANTS AND ECOLOGICAL COMMUNITIES
LP	Indicates taxa listed by the Pinelands Commission as endangered or threatened within their legal jurisdiction. Not all species currently tracked by the Pinelands Commission are tracked by the Natural Heritage Program. A complete list of endangered and threatened Pineland species is included in the NJ Pinelands Comprehensive Management Plan.
HL	Indicates taxa or ecological communities protected by the Highlands Water Protection and Planning Act within the jurisdiction of the Highlands Preservation Area.
GLOBAL & STATE CODE	GLOBAL ELEMENT RANK DEFINITIONS
G1	Critically imperiled globally because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) making it especially vulnerable to extinction.
G2	Imperiled globally because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it very vulnerable to extinction throughout its range.
G3	Either very rare and local throughout its range or found locally (even abundantly at some of its locations) in a restricted range (e.g., a single western state, a physiographic region in the East) or because of other factors making it vulnerable to extinction throughout its range; with the number of occurrences in the range of 21 to 100.
G4	Apparently secure globally ; although it may be quite rare in parts of its range, especially at the periphery.
G5	Demonstrably secure globally ; although it may be quite rare in parts of its range, especially at the

	periphery.
GH	Of historical occurrence throughout its range i.e., formerly part of the established biota, with the expectation that it may be rediscovered.
STATE CODE	STATE ELEMENT RANK DEFINITIONS
S1	Critically imperiled in New Jersey because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres). Elements so ranked are often restricted to very specialized conditions or habitats and/or restricted to an extremely small geographical area of the state. Also included are elements which were formerly more abundant, but because of habitat destruction or some other critical factor of its biology, they have been demonstrably reduced in abundance. In essence, these are elements for which, even with intensive searching, sizable additional occurrences are unlikely to be discovered.
S2	Imperiled in New Jersey because of rarity (6 to 20 occurrences). Historically many of these elements may have been more frequent but are now known from very few extant occurrences, primarily because of habitat destruction. Diligent searching may yield additional occurrences.
S3	Rare in state with 21 to 100 occurrences (plant species and ecological communities in this category have only 21 to 50 occurrences). Includes elements which are widely distributed in the state but with small populations/acreage or elements with restricted distribution, but locally abundant. Not yet imperiled in state but may soon be if current trends continue. Searching often yields additional occurrences.
S4	Apparently secure in the state, with many occurrences.
S5	Demonstrably secure in state and essentially ineradicable under present conditions.
SH	Elements of historical occurrence in New Jersey. Despite some searching of historical occurrences and/or potential habitat, no extant occurrences are known. Since not all of the historical occurrences have been field surveyed, and unsearched potential habitat remains, historically ranked taxa are considered possibly extant, and remain a conservation priority for continued field work with the expectation they may be rediscovered.
B	Refers to the breeding population of the element in the state.
N	Refers to the non-breeding population of the element in the state.
Note: To express <i>uncertainty</i> , the most likely rank is assigned and a question mark added (e.g., G2?). A range is indicated by combining two ranks (e.g., G1G2, S1S3).	
Source: NJDEP Division of Fish and Wildlife, March 22, 2010	

Endangered, Threatened & Special Concern Animals

The NJDEP Division of Fish and Wildlife, Endangered and Nongame Species Program’s (ENSP) mission is: “To actively conserve New Jersey’s biological diversity by maintaining and enhancing endangered and nongame wildlife populations within healthy functioning ecosystems.” The program is responsible for the protection and management of New Jersey’s wildlife, including 50 endangered, 36 threatened and 100 species currently listed as special concern (NJDEP Division of Fish and Wildlife, April 2, 2012 and February 21, 2012). For state-wide species lists, see **Internet Resources**.

A search of NJDEP Division of Parks and Forestry *Natural Heritage Database* in December 2016 revealed the documented presence of 19 special animal species in West Cape May (see **Tables 7.4** for code definitions and **Table 7.5** for list). Special Concern animal species, which warrant concern due to evidence of decline or vulnerability, include six birds. State Threatened species include five birds. State Endangered species within West Cape May include two amphibians and four birds. Two insect species, one moth and one beetle, are being tracked by the Endangered and Nongame Species Program, but do not have any special designation. Also included on the special animal report is a Migratory Raptor Concentration Site, which



has been designated as Endangered due to its importance for the protection of rare species.

Three additional animal species tracked through the Natural Heritage Program are known from the immediate vicinity of West Cape May, but have not been documented within the Borough. Nesting areas for the Federally Threatened and State Endangered piping plover (*Charadrius melodus*) and the Special Concern American oystercatcher (*Haematopus palliatus*) occur within a quarter mile of the Borough. A State Endangered butterfly, the bronze copper (*Lycaena hyllus*) has also been recorded as a casual flyby in the area. A summary of the habitat requirements for each species discussed in this section is provided in **Appendix D.3**.

Table 7.5: Natural Heritage Database Animal Species in West Cape May

Class	Common Name	Scientific Name	Feature Type	LP Rank	Protection Status	Global Rank	State Rank
Amphibia	Cope's Gray Treefrog	<i>Hyla chrysocelis</i>	Occupied habitat	4	State Endangered	G5	S1
	Cope's Gray Treefrog	<i>Hyla chrysocelis</i>	Vernal pool breeding	4	State Endangered	G5	S1
	Eastern Tiger Salamander	<i>Ambystoma t. tigrinum</i>	Occupied habitat	4	State Endangered	G5T5	S1
Aves	Bald Eagle	<i>Haliaeetus leucocephalus</i>	Foraging	4	State Endangered	G5	S1B, S2N
	Bald Eagle	<i>Haliaeetus leucocephalus</i>	Nest	4	State Endangered	G5	S1B, S2N
	Barred Owl	<i>Strix varia</i>	Non-breeding sighting	3	State Threatened	G5	S2B, S2N
	Black Skimmer	<i>Rynchops niger</i>	Foraging	4	State Endangered	G5	S1B, S1N
	Black-crowned Night Heron	<i>Nycticorax nycticorax</i>	Foraging	3	State Threatened	G5	S2B, S3N
	Bobolink	<i>Dolichonyx oryzivorus</i>	Breeding sighting	3	State Threatened	G5	S2B, S3N
	Brown Thrasher	<i>Toxostoma rufum</i>	Breeding sighting	2	Special Concern	G5	S3B, S4N
	Cattle Egret	<i>Bubulcus ibis</i>	Foraging	3	State Threatened	G5	S2B, S3N
	Common Tern	<i>Sterna hirundo</i>	Foraging	2	Special Concern	G5	S3B, S4N
	Glossy Ibis	<i>Plegadis falcinellus</i>	Foraging	2	Special Concern	G5	S3B, S4N
	Least Tern	<i>Sternula antillarum</i>	Foraging	4	State Endangered	G4	S1B, S1N
	Little Blue Heron	<i>Egretta caerulea</i>	Foraging	2	Special Concern	G5	S3B, S3N
	Migratory Raptor Concentration Site	<i>Raptor Winter concentration Area</i>	Concentration Site	4	State Endangered	GNR	SNR
	Osprey	<i>Pandion haliaetus</i>	Foraging	3	State Threatened	G5	S2B

Class	Common Name	Scientific Name	Feature Type	LP Rank	Protection Status	Global Rank	State Rank
	Osprey	<i>Pandion haliaetus</i>	Nest	3	State Threatened	G5	S2B
	Red-shouldered Hawk	<i>Buteo lineatus</i>	Breeding sighting	4	State Endangered	G5	S1B, S3N
	Snowy Egret	<i>Egretta thula</i>	Foraging	2	Special Concern	G5	S3B, S4N
	Tricolored Heron	<i>Egretta tricolor</i>	Foraging	2	Special Concern	G5	S3B, S3N
Insecta	Precious Underwing	<i>Catocala p. pretiosa</i>				G4T2	S2S3
	Southeastern Tiger Beetle	<i>Cicindela dorsalis media</i>				G4T3T4	S1S2
Note: See Table 7.4 for Global and State Rank definitions and Table 7.7 for Landscape Project Rank definitions.							
Source: Natural Heritage Program, December 15, 2016							

Of the 427 bird species that have been documented in Cape May County, 72 (about 17%) are listed as Endangered, Threatened or Special Concern (**Appendix D.4**). The location of West Cape May relative to the Cape May Migratory Bird Refuge Natural Heritage Priority Site (discussed in **Section 7F**) would make the Borough attractive to a variety of bird species. In addition to the fifteen rare bird species currently documented in the vicinity via the Natural Heritage Database, it is likely that a number of additional rare birds utilize habitat within West Cape May.

Endangered, Threatened & Special Concern Plants

The Endangered Plant Species List Act (N.J.S.A. 13:1B-15.151) was enacted in 1989, defining endangered plants as "any native plant species whose survival in the State or the nation is in jeopardy... and any species having five or fewer extant populations within the State." The Division of Parks and Forestry has the responsibility of creating the list of NJ endangered plant species (N.J.A.C. 7:5C-1.1). While the rule does not provide any protection for officially listed species, several regulatory agencies within NJDEP responsible for protecting plant habitat have incorporated the Endangered Plant Species List into their criteria for review of permits (NJDEP Division of Parks and Forestry, January 4, 2007).

Information on the rare plants and natural communities throughout the state is tracked in the *New Jersey Natural Heritage Database* by the NJDEP Office of Natural Lands Management (ONLM). A search of the Natural Heritage Database in December 2016 revealed records of four special concern plants in West Cape May Borough (see **Table 7.4** for code definitions and **Table 7.6** for the species list) (NJDEP ONLM, December 2016). In addition to the plants described in the table below, the Natural Heritage Database listed 27 rare plants known from the immediate vicinity of the Borough.

Appendix D.4 provides a list of all rare plant species known from Cape May County. Species known from West Cape May, and those known from the immediate vicinity, are identified as subsets within the list. Rare plants that have been documented at other locations in the county could be present in West Cape May if suitable habitat is present within the Borough.

Table 7.6: Natural Heritage Database Plant Species recorded in West Cape May.


Scientific Name	Common Name	Federal Status	NJ Protection	
<i>Diodia virginiana</i> var. <i>virginiana</i>	Larger Buttonweed	-	E	
Regional Status	Global Rank	State Rank	Last Observed	
LP, HL	G5T5	S1 (critically imperiled)	8/21/1991	
Madder Family (Rubiaceae)				
Description: plant 2-8 dm high, leaves opposite, entire and narrow, flowers white or pink with 4 petals and 2 sepals				
Habitat: moist, open, low ground of watersides and ditches				
Description from Gleason & Cronquist (1991).				
Scientific Name	Common Name	Federal Status	NJ Protection	
<i>Hydrocotyle verticillata</i> var. <i>verticillata</i>	Whorled Marsh-pennywort	-	-	
Regional Status	Global Rank	State Rank	Last Observed	
HL	G5T5	S3 (rare)	2009	
Parsley Family (Apiaceae)				
Description: stem creeping, leaves to 6 cm high, peltate, rounded and lobed; inflorescence a forking spike of tiny white flowers with five petals				
Habitat: wet or moist ground of bogs, swampy woods, watersides				
Description from Gleason & Cronquist (1991).				
Scientific Name	Common Name	Federal Status	NJ Protection	
<i>Malus angustifolia</i> var. <i>puberula</i>	Spiny wild Crabapple	-	-	
Regional Status	Global Rank	State Rank	Last Observed	
HL	G5?T2T4	S2 (imperiled)	2010	
Rose Family (Rosaceae)				
Description: tall shrub or low tree with thorny branches, leaves alternate, ovate and toothed, flowers pink to white with 5 petals. Variety distinguished by dense, felty look of twigs and pedicels.				
Habitat: woods and thickets				
Species description from Gleason & Cronquist (1991). Variety description from Virginia Botanical Associates (2017).				

Image credit to Robert H. Mohlenbrock, hosted by the USDA-NRCS PLANTS Database / USDA SCS. 1989. Midwest wetland flora: Field office illustrated guide to plant species. Midwest National Technical Center, Lincoln.

Image credit to USDA-NRCS PLANTS Database / Britton, N.L., and A. Brown. 1913. An illustrated flora of the northern United States, Canada and the British Possessions. 3 vols. Charles Scribner's Sons, New York. Vol. 2: 649.

Image (species only) credit to Eleanor, North Carolina State University/A&T State University Cooperative Extension, Plant Database: *Malus angustifolia*. Available online at <https://plants.ces.ncsu.edu/plants/all/malus-angustifolia/>

Scientific Name	Common Name	Federal Status	NJ Protection
<i>Sesuvium maritimum</i>	Seabeach purslane	-	-
Regional Status	Global Rank	State Rank	Last Observed
HL	G5	S2 (imperiled)	2010
Carpetweed Family (Aizoaceae)			
Description: plant succulent and sprawling, leaves opposite, oblanceolate and smooth, flowers lacking petals but with 5 sepals			
Habitat: sea beaches			
Description from Gleason & Cronquist (1991).			
Note: For status and rank definitions, refer to Table 7.4 .			
Source: NJDEP ONLM, December 15, 2016			




Image credit to USDA-NRCS PLANTS Database / Britton, N.L., and A. Brown. 1913. An illustrated flora of the northern United States, Canada and the British Possessions. 3 vols. Charles Scribner's Sons, New York. Vol. 2: 34.

E. Protecting Endangered, Threatened and Special Concern Species

Wildlife Action Plan

NJDEP Division Fish and Wildlife prepared a Wildlife Action Plan (WAP) in 2008, required by the US Fish and Wildlife Service in order to qualify for future federal funds through the State Wildlife Grants program. This program provides federal funds to states for the conservation of species that are endangered, threatened, or have special conservation needs. A 25% match, provided by citizen contributions, is required. NJ has received approximately \$1.2 million dollars of State Wildlife Grants funding each year (NJDEP, January 23, 2008).

The report states,

"The greatest threats to NJ's natural resources include habitat loss, destruction, alteration, and fragmentation. This has been a recurring theme within NJ for years as it is the most densely populated state in our nation with an annually increasing population requiring additional homes, roads, commercial buildings, schools, etc. Additional threats include, but are not limited to, invasive species (flora and fauna, aquatic and terrestrial), pollution, and unsustainable land management practices." (NJDEP, January 23, 2008)

A Wildlife Action Plan specific to each region identifies habitats, wildlife of greatest conservation need, and threats. Conservation goals and actions are identified and prioritized, and potential partnerships are outlined with landowners, the public and conservation organizations, wildlife professionals and local, state and federal agencies. West Cape May is in the Peninsula conservation zone within the Delaware Bays landscape region.

The Landscape Project

The state's *Landscape Project* (see **Figure 7f**) is a pro-active, ecosystem-level approach to the long-term protection of rare species and their important habitats in New Jersey. Its goal is to protect New Jersey's biological diversity by maintaining and enhancing rare wildlife populations within healthy, functioning ecosystems. It provides users with peer reviewed, scientifically sound wildlife data that is easily accessible and can be used by state, county, and local governments, as well as nongovernmental conservation organizations and private land

owners for planning, open space acquisition, and land-use regulation (NJDEP Division of Fish and Wildlife, 2012).

The NJDEP, Division of Fish and Wildlife, Endangered and Nongame Species Program is responsible for the Landscape Project. Version 3.1 was released in 2012. The dataset was created by intersecting endangered, threatened and priority species data with the 2007 Land Use/Land Cover GIS layer, which was derived from aerial photography. The resulting data layer identifies, delineates and ranks (based on the conservation status of species present) critical habitat statewide. **Table 7.7** lists rank definitions. Each habitat patch is coded for the number of special concern, state threatened, state endangered and federally listed species present.

Table 7.7: Landscape Project Habitat Rank Definitions

Rank	Definition
0	No Suitable Habitat
1	Suitable Habitat – Rank 1 is assigned to patches that meet habitat-specific suitability requirements such as minimum size criteria for endangered, threatened or priority wildlife species, but that do not intersect with any confirmed occurrences of such species.
2	Special Concern – Rank 2 is assigned to patches containing one or more occurrences of species considered to be species of special concern.
3	State Threatened – Rank 3 is assigned to patches containing one or more occurrences of State threatened species.
4	State Endangered – Rank 4 is assigned to patches with one or more occurrences of State endangered species.
5	Federally Listed – Rank 5 is assigned to patches containing one or more occurrences of wildlife listed as endangered and threatened pursuant to the Federal Endangered Species Act of 1973.

Source: NJDEP Division of Fish and Wildlife, 2012

Over 90% of West Cape May is ranked as habitat for priority species according to the Landscape Project Version 3.3, while 4.4% is Rank 1 for potential habitat (see **Table 7.8** and **Figure 7f**).

Table 7.8: Landscape Project version 3.3

Land Use Type	Acres	Percent
No suitable habitat	30.26	4.0%
Rank 1	33.12	4.4%
Rank 2	0.00	0.0%
Rank 3	0.00	0.0%
Rank 4	691.25	91.6%
Rank 5	0.00	0.0%
Total Acres	754.63	100.0%

Source: NJDEP Division of Fish and Wildlife, May 9, 2017

Natural Heritage Grid and Priority Sites

The NJDEP Office of Natural Lands Management (ONLM) has developed the Natural Heritage Grid Map (see **Figure 7g**²⁷), which provides a general representation of the locations of rare plant species and natural communities, including both historically and recently documented

²⁷ The Natural Heritage Database search results (2013) differed from the most recent GIS data (2009) for the Natural Heritage Grid, therefore the search information (more recent) is shown.

habitat. The purpose of the Grid Map is to document rare plant species and natural community habitats to inform decision-makers who need to address the conservation of natural resources. The map identifies potentially sensitive areas, and indicates where custom database searches are needed for land use decision-making. Seven different quadrangles on the grid map include portions of West Cape May, and a total of 40 rare plants or communities are noted as occurring within those sections (NJ-GeoWeb, 2017). A custom search of the Natural Heritage Program Database conducted on December 15, 2016 revealed that four of those species records are from within West Cape May Borough (see **Table 7.6** for descriptions), and 27 other rare plant species have been recorded within a quarter mile of the borough (identified with green text in Appendix D.5). The Grid Map does not include habitat for animal species, and not all areas have been surveyed (NJDEP ONLM, November 2009).

In addition, the Natural Heritage Program makes lists of New Jersey rare plant species and ecological communities by county (see **Appendix D.5**). If suitable habitat exists in the borough, it is possible that some of those species could be found in West Cape May (NJDEP ONLM NHP, July 30, 2008).


Natural Heritage Priority Sites have been identified by the ONLM as areas critically important for preservation of New Jersey's biological diversity. These are considered some of the best and most viable occurrences of endangered and threatened plant species and natural communities, but other occurrences of endangered and threatened plant species may exist. A large Natural Heritage Priority Site is located at the tip of the Cape May peninsula, and it includes portions of West Cape May Borough (see **Figure 7g and Appendix D.6**). Features that contribute to the importance of the site include a high concentration of migratory birds, state imperiled plant and animal species, globally imperiled animal species, high quality coastal plant communities, and vernal habitat.

F. Invasive Nonindigenous Species

Non-native species (also called alien, exotic or introduced species) are those species that have been introduced outside their natural geographic range as a result of human actions, whether intentionally (e.g. as sources of food, for landscaping purposes or the release of unwanted pets) or unintentionally (e.g. in the ballast of a ship or in a load of lumber). Executive Order 13112 defines an *invasive species* as a species that is non-native to the ecosystem and whose introduction causes or is likely to cause economic or environmental harm or harm to human health (USDA, February 3, 1999). The most problematic of these displace native species, contribute to local elimination of species or even extinctions, alter the community structure, and may eventually disrupt ecosystem processes (Snyder et al, 2004). Preliminary research in NJ has documented over 1,200 species of nonindigenous plant species, or as much as 62% of the state's total vascular flora (Snyder et al, 2004).

Flora

Native plants can be susceptible to introduced diseases, which they have not evolved resistance to. The chestnut blight fungus was an accidental introduction that destroyed all mature American chestnut (*Castanea dentata*) trees, once one of the dominant trees in the New Jersey landscape. Another introduced fungus, Dutch elm disease, destroyed the American elm (*Ulmus americana*).



Legend


- West Cape May
- Water
- Streams
- Parcels
- Roads

Natural Heritage Priority Site

- Cape May Migratory Bird Refuge

Natural Heritage Grid Map (2009)

- Generalized Locations of Rare Plant Species and Ecological Communities
- S - Documented Location Known Precisely
 - M - Documented Location Known Within 1.5 Miles
 - Both - Both 'M' and 'S' Occurrences



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Figure 7g: West Cape May Natural Heritage Priority Sites & Grid
The Natural Heritage Grid Map indicates the general locations of rare plant species and communities. Natural Heritage Priority Sites are areas that have been identified as critically important for preservation of New Jersey's biological diversity.

In addition, native plants may have little resistance to certain introduced insects, and/or these insects may have no natural enemies in their new surroundings, allowing them to rapidly reach pest proportions. Introduced insects which may be impacting West Cape May's trees include the southern pine beetle, the gypsy moth, and the emerald ash borer (NJ Forest Service, 2010). The pests weaken their host trees, which often succumb to successive years of infestation, to diseases carried by the insects, or other environmental stresses.

For these reasons, the Final Report of the New Jersey Comparative Risk Project, which evaluated the relative risks of environmental problems to the people and ecosystems of New Jersey identified invasive species (including plants, insects, and other organisms) as one of the state's top environmental problems (Steering Committee of the New Jersey Comparative Risk Project, 2003).


While there is no official invasive species list for New Jersey, An Overview of Nonindigenous Plant Species in New Jersey (Snyder et al, 2004) profiles 27 nonindigenous plant species that aggressively invade natural plant communities in New Jersey, which are summarized in **Table 7.9**. Some of the most problematic long-established invasive exotic species observed in West Cape May include English ivy, multiflora rose, Japanese honeysuckle, and common reed. The first three are illustrated in the table below, but the last is not.

Table 7.9: Invasive Nonindigenous Plants

Scientific Name	Common Name	Problems Caused	Illustration	Illus. Source
<i>Acer platanoides</i>	Norway maple	Dispersed seeds easily sprout in shade, crowding out native plants. Canopy produces deep shade and roots produce a toxic substance preventing growth of wildflowers and other trees under its canopy.		Jan Samanek, State Phytosanitary Administration, Bugwood.org
<i>Ailanthus altissima</i>	tree of heaven	Aggressive in disturbed areas, crowding out native plants.		Britton and Brown, 1913, Vol. 2: 446.
<i>Alliaria petiolata</i>	garlic mustard	Aggressive in shady habitats, crowding out native plants.		Deborah J. Kratzer

<i>Artemisia vulgaris</i>	Mugwort or common wormwood	Aggressive, crowding out native plants.		Deborah J. Kratzer
<i>Berberis thunbergii</i>	Japanese barberry	Can grow so thick in the understory of open forests that it shades out indigenous understory plants. Affects soil properties, particularly pH, which can affect plant establishment. Can form nearly impenetrable thorny thickets that impact the recreational value of natural lands.		Deborah J. Kratzer
<i>Celastrus orbiculatus</i>	Oriental bittersweet	The vine twines around surrounding plants, impeding sap flow. Also makes host plants too heavy, increasing wind, snow & ice damage.		Deborah J. Kratzer
<i>Cirsium arvense</i>	Canada thistle	Competes with crops and degrades pastures (inedible to livestock).		Deborah J. Kratzer
<i>Dipsacus fullonum</i>	wild teasel	Highway mowing equipment and discarded dried teasel heads from flower arrangements can lead to the establishment of new colonies, often forming a monoculture that displaces native communities.		Steve Dewey, Utah State University, Bugwood.org
<i>Elaeagnus umbellata</i>	autumn olive	Sprouts vigorously in disturbed areas, produces shade, preventing sprouting of native trees.		Deborah J. Kratzer

<i>Euonymus alatus</i>	burning bush	Grows well in many sites, especially upland forests and pastures, crowding out native plants.		James H. Miller, USDA Forest Service, Bugwood.org
<i>Hedera helix</i>	English ivy	Grows vigorously in deep shade, inhibiting growth of native woodland plants. Vines up tree trunks, adding to weight, and increasing likelihood of wind damage.		Deborah J. Kratzer
<i>Ligustrum vulgare</i>	common privet	Crowds out more desirable native plants.		USDA PLANTS Database, Bugwood.org
<i>Lonicera japonica</i> Thunberg	Japanese honeysuckle	Spreads aggressively in disturbed habitats, crowding out native plants. Aggressive roots can decrease the growth of native trees and vines. Vines engulf small trees and shrubs, causing them to collapse. Leaves out very early in spring, which could inhibit flowering by spring ephemerals.		Deborah J. Kratzer
<i>Lythrum salicaria</i>	purple loosestrife	Spreads aggressively in wetlands, eliminating open water habitats and crowding out native plants. Contributes to the loss of wildlife that depend on native wetland plants.		John D. Byrd, Mississippi State University, Bugwood.org
<i>Microstegium vimenium</i>	Japanese stiltgrass	Spreads aggressively in disturbed, moist, shady areas, crowding out native plants. May raise pH and reduce organic soil horizon.		Deborah J. Kratzer

<i>Myriophyllum spicatum</i> L.	Eurasian water-milfoil	An aquatic plant that begins growing earlier in spring than most indigenous aquatic plants, it quickly overtops, outshades, and outcompetes surrounding vegetation.		Britton and Brown, 1913, Vol. 2: 614.
<i>Miscanthus sinensis</i>	Chinese silver grass	Escapes from ornamental plantings and can form large clumps along disturbed areas, crowding out native vegetation. It is also extremely flammable and increases fire risks where it grows.		James H. Miller, USDA Forest Service, Bugwood.org
<i>Phyllostachys aurea</i>	Golden bamboo	Forms dense monocultural thickets that crowd out other plants. Difficult to eradicate once established.		Chuck Barger, Univ. of Georgia, Bugwood.org
<i>Polygonum cuspidatum</i>	Japanese knotweed	Spreads aggressively in disturbed, sunny areas, especially river banks and wetlands, crowding out native plants.		Tom Heutte, USDA Forest Service, Bugwood.org
<i>Potamogeton crispus</i> L.	curly leaf pondweed	An aquatic plant that begins growing earlier in spring than most indigenous aquatic plants, it quickly overtops, outshades, and outcompetes surrounding vegetation. Can form dense mats that disrupt boating, swimming, and fishing.		Mohlenbrock, 1995
<i>Rosa multiflora</i>	multiflora rose	Spreads everywhere, except standing water, crowding out native plants and degrading pastures.		James H. Miller, USDA Forest Service, Bugwood.org

<i>Rubus phoenicolasius</i>	wineberry	Forms an extensive, nearly impenetrable understory layer in favorable locations such as moist soils in forests over dolomite, marble, shale, diabase, and traprock, crowding out native plants.		Jill M. Swearingen, USDI National Park Service, Bugwood.org
<i>Viburnum plicatum</i>	Japanese viburnum	Shade tolerant shrub considered highly threatening to native plant communities.		Richard Webb, Self-employed horticulturist, Bugwood.org
<i>Vinca minor</i>	periwinkle	Spreads in shady forests, crowding out native plants.		Jill M. Swearingen, USDI National Park Service, Bugwood.org
<i>Wisteria floribunda</i> and <i>W. sinensis</i>	Japanese and Chinese Wisteria	Aggressive climbing vines that girdle tree trunks and branches. Dense canopies weigh down branches and shade underlying areas.		Ted Bodner at USDA-NRCS PLANTS Database
Sources: Snyder & Kaufman, 2004; Swearagain et al., 2002; Courtney, 1997; Center for Invasive Species and Ecosystem Health (bugwood.org), 2010; Britton and Brown, 1913; Mohlenbrock, 1995; Bodner at USDA-NRCS PLANTS Database				

Common reed (*Phragmites australis ssp. australis*) has sometimes been excluded from invasive species lists due to its strong resemblance to a native subspecies (*Phragmites australis ssp. americanus*). The two subspecies have recently been found to be morphologically distinguishable (Sarver et al., 2008). While the invasive subspecies is widely established throughout New Jersey, the native subspecies is only known from Atlantic County (Kartesz, 2013). Because the large grass forms large, monotypic stands that completely exclude other wetland vegetation it is considered highly threatening to native communities.



The New Jersey Invasive Species Strike Team (NJISST) focuses on preventing the spread of newer invasive species throughout the state (NJISST, 2016). The team is tracking 35 invasive species in Cape May County, 34 of which are plants (**Appendix E**). The strike team provides links to fact sheets which provide information regarding identification, threat levels and control measures for each species tracked in their system (NJISST, 2016). Although their focus is on eradicating newly introduced species before they can establish and spread, fact sheets are also available for many of the widespread invasives. Four species that NJISST is currently tracking at multiple locations in West Cape May Borough are invasive vines (**Table 7.10**).




Table 7.10: Some Recent Invasives Tracked in West Cape May

Scientific Name	Common Name	Description	Threat
<i>Pueraria montana var. lobata</i>	Kudzu	climbing or trailing semi-woody vine with alternate compound leaves, long clusters of purple flowers, brown, hairy flattened seed pods	Highly threatening to native communities
<i>Ampelopsis brevipedunculata</i>	Porcelain-berry, amur peppervine	climbing woody vine with alternate, lobed and coarsely-toothed leaves, small greenish white flowers, mature fruits bright blue	Highly threatening to native communities
<i>Clematis terniflora</i>	Sweet autumn virginsbower, Japanese clematis, leatherleaf clematis	climbing woody vine with opposite compound leaves, small white flowers with 4 petals, fruit with long feathery plumes	Highly threatening to native communities
<i>Persicaria perfoliata</i>	Mile-a-minute	Climbing annual prickly vine grows up to 6" per day, alternate, toothless, triangular leaves with saucer-shaped leaf surrounds the stem at base of leaf stem and base of flower, small white flowers, small round fruit turns blue when mature	Widespread and highly threatening to native plant communities
<i>Albizia julibrissin</i>	mimosa, Persian silk tree, powder puff tree, silk tree, silky acacia, Japanese mimosa	small deciduous tree with bipinnate (double compound) leaves, flowers are showy, tight clusters of pink stamens with a white base, looking like silky threads, fruit is a flat brown pod	regionally common or abundant, moderately threatening to native communities

Source: NJISST, 2016

Fauna

As with exotic plant species, the introduction of non-native animal species can have a devastating effect on natural communities. This most often occurs due to competition with native animals for limited resources such as food and shelter, but it may also be due to predation on native species. An example of the latter is the feral cat (*Felis domesticus*), which is the mammal tracked by the New Jersey Invasive Species Strike Team that is most likely to occur in West Cape May Borough. Feral cats are a widespread problem around the state, and are considered highly threatening to native communities (NJISST, 2016). The nutria (*Myocastor coypus*) has a more limited presence, but is occasionally seen in the southern part of the state (Van Clef, 2009).

Fact sheets with information about identification, threat levels and control measures for invasive animal species are also provided by the Strike Team (NJISST, 2016). All five of the invasive bird species tracked are commonly encountered throughout New Jersey, including Cape May County (see **Appendix C.1**), and are likely to be present in West Cape May. Both the house finch and the house sparrow are classified as mildly threatening to natural communities. The European starling, which poses a moderate threat to native species, is frequently seen in large flocks: the high count for this species in Cape May County was estimated at 5,000 birds in 2013 (Sullivan et. al., 2009). Both the mute swan and brown-headed cowbird are rated as highly threatening to native communities. In addition to competing with native birds for resources, the brown-headed cowbird is a brood parasite, laying its eggs in the nests of other bird species, which then raise the young cowbirds at the expense of their own offspring.

The sole invasive reptile likely to be encountered in Cape May County is the red-eared slider, which is classified as highly threatening. This turtle is widespread in New Jersey, and may be found in ponds, lakes, swamps, streams, or slow-flowing rivers.

No amphibians are currently tracked by NJISST. Although included on the DEP's list of New Jersey amphibians, the barking treefrog (*Hyla gratiosa*) is not considered native to the state. The species was at one time established and breeding in Cape May County southern New Jersey, but was thought to be introduced and is currently presumed extirpated (USGS 2017).

The state Division of Fish and Wildlife (2016) lists nine species that pose a serious threat to freshwater resources, and which must be destroyed when encountered. Those invasive fish include the swamp eel, grass carp, bighead carp, silver carp, flathead catfish, brook stickleback, green sunfish, warmouth, and oriental weatherfish. The NJISST tracks all of those species, and one additional freshwater fish. The northern snakehead (*Channa argus*) is not included on the state list of freshwater fish provided in **Appendix C.5**, although the Strike Team fact sheet states that it is widespread in New Jersey. No threat level is given for the snakehead.

The Invasive Species Strike Team additionally tracks a number of invertebrate species in the state, including 18 insects, 12 mollusks, 4 crustaceans and 1 worm. Fact sheets are also available for the invasive invertebrates (NJISST, 2016).

References: Biological Resources

Dominant Vegetation (Land Cover)

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice. 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

NJDEP, Bureau of Geographic Information Systems (BGIS). February 17, 2015. Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague 1:2,400. GIS Data. <http://www.nj.gov/dep/gis/listall.html>

Steering Committee of the New Jersey Comparative Risk Project. March 2003. Final Report of the New Jersey Comparative Risk Project. 213 pages. <http://www.state.nj.us/dep/dsr/njcrp/njcrp-final.pdf>

US Geological Survey (USGS). 2010. Edited by NJDEP, Bureau of Geographic Information Systems (BGIS) 1998, 2000, 2001, 2002, 2005, 2007, 2010. NJDEP MODIFIED ANDERSON SYSTEM 2007 derived from: A Land Use and Land Cover Classification System for Use with Remote Sensor Data, USGS Professional Paper 964, 1976. 35 pages. <http://www.state.nj.us/dep/gis/digidownload/metadata/lulc07/anderson2007.html>

Wildfire Fuel Hazard

NJDEP, New Jersey Forest Fire Service (NJFFS). April 17, 2009. 2002 NJFFS Wildfire Fuel Hazard for Cape May County, New Jersey. GIS Data. <http://www.state.nj.us/dep/gis/njffh.html#CAP>

Wildlife

Burnett, Andrew. 2004. White-tailed Deer – Natural History and Autumn Behavior. NJDEP Division of Fish and Wildlife. 1 page. <http://www.njfishandwildlife.com/deerart.htm>

Frank, C.A. 2010. Important Bird Areas of New Jersey. Edited by C.D. Tonkinson. New Jersey Audubon, Bernardsville, New Jersey.

Gessner, Jackie and Eric Stiles. February 2001. Field Guide to Reptiles and Amphibians of New Jersey. Funded By: Conserve Wildlife Funds And U.S. Fish & Wildlife Service's Partnerships For Wildlife. 1st Edition, Herp Atlas Field Guide. <http://www.state.nj.us/dep/fgw/ensp/herpguide.htm>

Honachefsky, William B. 2000. Ecologically Based Municipal Land Use Planning. Lewis Publishers: New York. 255 pages.

Kenney, Leo P., Matthew R. Burne, with adaptations and modifications for New Jersey by Jason Tesauro, Kris Schantz and Melissa Craddock. No date. Salamanders, Frogs and Turtles of New Jersey's Vernal Pools: A Field Guide. NJDEP Division of Fish and Wildlife, Endangered and Nongame Species Program. 54 pages.

Latham, Roger Earl and, Jan Beyea, Merlin Benner, Cindy Adams Dunn, Mary Ann Fajvan, Ronald R. Freed, Marrett Grund, Stephen B. Horsley, Ann Fowler Rhoads, and Bryon P. Shissler. January 2005. Managing White-tailed Deer in Forest Habitat From an Ecosystem Perspective. Pennsylvania Case Study Report of the Deer Management Forum convened by Audubon Pennsylvania and the Pennsylvania Habitat Alliance in Harrisburg, Pennsylvania. 340+ pages. <http://pa.audubon.org/sites/default/files/documents/pa-deermanagement--full--textoffullreport.pdf>

NJDEP Division of Fish and Wildlife. No date. Canada Goose Control Information. 4 pages. http://www.njfishandwildlife.com/pdf/goose_control_info.pdf

NJDEP Division of Fish, Game and Wildlife with the Department of Agriculture. 1999. The Governor's Report on Deer Management in New Jersey. 32 pages. <http://www.state.nj.us/dep/fgw/pdf/govdrrpt.pdf>

NJDEP Division of Fish and Wildlife. 2004a. Checklist of NJ Birds. <http://www.state.nj.us/dep/fgw/chkbirds.htm>

NJDEP Division of Fish and Wildlife. 2004b. Checklist of NJ Mammals. <http://www.state.nj.us/dep/fgw/chkmamls.htm>

NJDEP Division of Fish and Wildlife, Endangered and Nongame Species Program (ENSP). February 21, 2012. NJDEP Species Based Habitat, Vernal Habitat (Version 3.1, 20120221). GIS data. <http://www.nj.gov/dep/gis/landscape.html>

NJDEP Division of Fish and Wildlife. March 1, 2017. White-tailed Deer in New Jersey. <http://www.njfishandwildlife.com/deer.htm>

NJDEP Division of Fish and Wildlife. July 22, 2016. Places to Fish. <http://www.state.nj.us/dep/fgw/fishplc.htm>

NJDEP Division of Fish and Wildlife. 2016. Checklist of NJ Fish. <http://www.njfishandwildlife.com/chkfish.htm>

NJDEP Division of Fish and Wildlife. June 6, 2014. Online Field Guide for Reptiles and Amphibians. http://www.state.nj.us/dep/fgw/ensp/fieldguide_herps.htm

NJDEP Division of Fish and Wildlife. June 4, 2013. Fishing Access Locations on Trout Waters - County Listing. <http://www.state.nj.us/dep/fgw/accesscnty.htm>

NJDEP Division of Watershed Management. March 2001. Management of Canada Geese in Suburban Areas: A Guide to the Basics. DRAFT. 26 pages. http://www.state.nj.us/dep/watershedmgt/DOCS/BMP_DOCS/Goosedraft.pdf

NJENSP (Endangered and Nongame Species Program). November 2014. The Leopard Frogs of New Jersey. http://www.njfishandwildlife.com/ensp/pdf/leopardfrog_guide.pdf

Remington, Vernick & Walberg Engineers. 2003. Environmental Resource Inventory, Borough of West Cape May.

Cape May County, New Jersey.

Sauer, Leslie Jones. 1998. The Once and Future Forest: A Guide to Forest Restoration Strategies. Island Press: Washington, D.C. 382 pages.

Sullivan, B.L., C.L. Wood, M.J. Iliff, R.E. Bonney, D. Fink, and S. Kelling. 2009. eBird: a citizen-based bird observation network in the biological sciences. *Biological Conservation* 142: 2282-2292.
<http://ebird.org/ebird/nj/places>. Accessed March 2017.

Tesauro, Jason. No date. New Jersey's Vernal Pools. NJDEP Division of Fish and Wildlife.
<http://www.state.nj.us/dep/fgw/vpoolart.htm>

USDA Animal and Plant Health Inspection Service. January 2003. Fact Sheet: Managing Canada Goose Damage. 9 pages. http://www.aphis.usda.gov/wildlife_damage/states/pennsylvania/fact_sheet_goose.shtml

USFWS (U.S. Fish & Wildlife Service). 1997. Significant Habitats and Habitat Complexes of the New York Bight Watershed: Cape May Peninsula. https://nctc.fws.gov/resources/knowledge-resources/pubs5/web_link/text/cmp_form.htm

Endangered, Threatened and Special Concern Species: Animals
Conserve Wildlife Foundation and NJDEP Division of Fish and wildlife. 2002. Endangered and Threatened Wildlife of New Jersey. 2 pages. <http://www.state.nj.us/dep/fgw/ensp/pdf/e&tbroch.pdf>

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program. December 15, 2016. Letter Re: West Cape May Borough Natural Resource Inventory. Natural Heritage Database and the Landscape Project habitat mapping search results.

NJDEP Division of Fish and Wildlife. April 2, 2012. New Jersey's Endangered and Threatened Wildlife.
<http://www.njfishandwildlife.com/tandespp.htm>

NJDEP Division of Fish and Wildlife. February 21, 2012. NJ Endangered and Nongame Species Program Special Concern – Species Status Listing. 4 pages. <http://www.njfishandwildlife.com/ensp/pdf/spclspp.pdf>

NJDEP Division of Fish and Wildlife. January 18, 2011. Notice of Readoption with Amendments: Division of Fish and Wildlife rules; Endangered Species List, Nongame Species List, N.J.A.C. 7:25-4.1, 4.13, 4.17
<http://www.nj.gov/dep/rules/notices/011811b.html>

NJDEP Division of Fish and Wildlife. March 22, 2010. Explanations of Codes Used in Natural Heritage Reports.
http://www.nj.gov/dep/parksandforests/natural/heritage/spplant_ap1.html

NJ Division of Fish and Wildlife. October 6, 2004. Endangered Species Conservation Act: 30 Years of Protection.
<http://www.state.nj.us/dep/fgw/ensp/somhome.htm>

Endangered, Threatened and Special Concern Species: Plants

Gleason, H.A. and A. Cronquist. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. Second Edition. The New York Botanical Garden, Bronx, New York.

NJDEP Division of Parks and Forestry. January 4, 2007. Endangered Plant Species Program.
<http://www.state.nj.us/dep/parksandforests/natural/endplants.html>

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program. December 15, 2016. Letter Re: West Cape May Borough Natural Resource Inventory. Natural Heritage Database and the Landscape Project habitat mapping search results.

Virginia Botanical Associates. 2017. Digital Atlas of the Virginia Flora: *Malus angustifolia* (Aiton) Michx.
<http://vaplantatlas.org/index.php?do=plant&plant=2980> .

Protecting Endangered, Threatened and Special Concern Species

NJDEP Division of Fish and Wildlife. May 9, 2017. NJDEP Species Based Habitat, Vernal Habitat, Version 3.3 (Envr_hab_ls_v3_3_vernalhabitat). GIS Data: <http://www.state.nj.us/dep/gis/landscape.html#geodatabase>

NJDEP Division of Fish and Wildlife. May 9, 2017. NJDEP Species Based Habitat, Vernal Pools, Version 3.3 (Envr_hab_ls_v3_3_vernalpools). GIS Data: <http://www.state.nj.us/dep/gis/landscape.html#geodatabase>

NJDEP Division of Fish and Wildlife. May 9, 2017. NJDEP Species Based Habitat, Delaware Bay Region, Version 3.3 (Envr_hab_ls_v3_3_delbay). GIS Data: <http://www.state.nj.us/dep/gis/landscape.html#geodatabase>

NJDEP Division of Fish and Wildlife. May 9, 2017. NJDEP Species Based Habitat, Atlantic Coastal Region, Version 3.3 (Envr_hab_ls_v3_3_coastal). GIS Data: <http://www.state.nj.us/dep/gis/landscape.html#geodatabase>

NJDEP Division of Fish and Wildlife. January 23, 2008. New Jersey Wildlife Action Plan for Wildlife of Greatest Conservation Need. Overview: <http://www.state.nj.us/dep/fgw/ensp/wap/pdf/wapintro.pdf>

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program (NHP). December 15, 2016. Letter Re: West Cape May Borough Natural Resource Inventory. Natural Heritage Database and the Landscape Project habitat mapping search results.

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program (NHP). March 21, 2013. Natural Heritage Grid Map. GIS Data. <http://www.state.nj.us/dep/gis/stateshp.html#NHPGRID>

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program (NHP). March 1, 2007. NJDEP Natural Heritage Priority Sites. GIS Data. <http://www.state.nj.us/dep/gis/stateshp.html#PRIORITY>

NJDEP Office of Natural Lands Management (ONLM) Natural Heritage Program (NHP). July 30, 2008. New Jersey Rare Plant Species and Ecological Community Lists By County. <http://www.state.nj.us/dep/parksandforests/natural/heritage/countylist.html>

NJ-GeoWeb. 2017. NJDEP Bureau of GIS. Interactive Mapping Application. <http://www.state.nj.us/dep/gis/geoweb splash.htm>. Accessed March 9, 2017.

Invasive Species

Bodner, Ted. 2005. USDA-NRCS PLANTS Database / James H. Miller and Karl V. Miller. Forest plants of the southeast and their wildlife uses. University of Georgia Press., Athens.

Britton, N.L., and A. Brown at USDA-NRCS PLANTS Database. 1913. An illustrated flora of the northern United States, Canada and the British Possessions. 3 vols. Charles Scribner's Sons, New York. <http://plants.usda.gov>

Center for Invasive Species and Ecosystem Health. 2010. Invasive and Exotic Species of North America. University of Georgia. Images by Chuck Barger, Univ. of Georgia; John D. Byrd, Mississippi State University; Steve Dewey, Utah State University; Tom Heutte, USDA Forest Service; James H. Miller, USDA Forest Service; Jan Samanek, State Phytosanitary Administration; and Richard Webb, Self-employed horticulturist. <http://www.invasive.org> and <http://bugwood.org>

Courtney, John Mark. 1997. Fact Sheets: Invasive Exotics. Bowman's Hill Wildflower Preserve, New Hope, PA. <http://www.bhwp.org/resources/Invasive-Plants.htm>

Kartesz, J.T. 2013. The Biota of North America Program (BONAP). North American Plant Atlas. Chapel Hill, N.C. [maps generated from Kartesz, J.T. 2013. Floristic Synthesis of North America, Version 1.0. Biota of North America Program (BONAP). (in press)]. <http://bonap.net/tdc>

Mohlenbrock, Robert H. at USDA-NRCS PLANTS Database / USDA NRCS. 1995. Northeast wetland flora: Field office guide to plant species. Northeast National Technical Center, Chester. <http://plants.usda.gov>

NJDEP Division of Fish and Wildlife. 2016. Checklist of NJ Fish. <http://www.njfishandwildlife.com/chkfish.htm>

NJ Forest Service. 2010. Forest Health. http://www.state.nj.us/dep/parksandforests/forest/njfs_forest_health.html

NJISST (New Jersey Invasive Species Strike Team Team). 2016. Searchable Invasive Species Map and Invasive Species Fact Sheets. <http://www.njisst.org/> Accessed March 2017.

Sarver, M., A. Treher, L. Wilson, R. Naczi and F.B. Kuehn. 2008. Mistaken Identity? Invasive plants and their native look-alikes: an identification guide for the Mid-Atlantic. Delaware Department of Agriculture. http://www.nybg.org/files/scientists/rnaczi/Mistaken_Identity_Final.pdf

Snyder, David and Sylvan R. Kaufman. 2004. An overview of nonindigenous plant species in New Jersey. New Jersey Department of Environmental Protection, Division of Parks and Forestry, Office of Natural Lands Management, Natural Heritage Program: Trenton, NJ. 107 pages. <http://www.state.nj.us/dep/parksandforests/natural/heritage/InvasiveReport.pdf>

Steering Committee of the New Jersey Comparative Risk Project. 2003. Final Report of the New Jersey Comparative Risk Project. 213 pages. <http://www.state.nj.us/dep/dsr/njcrp/>

Sullivan, B.L., C.L. Wood, M.J. Iliff, R.E. Bonney, D. Fink, and S. Kelling. 2009. eBird: a citizen-based bird observation network in the biological sciences. *Biological Conservation* 142: 2282-2292. <http://ebird.org/ebird/nj/places>. Accessed March 2017.

Swearingen, J., K. Reshetiloff, B. Slattery, and S. Zwicker. 2002. Plant Invaders of Mid-Atlantic Natural Areas. National Park Service and U.S. Fish & Wildlife Service, Washington, D.C. 82 pp. <http://www.nps.gov/plants/alien/pubs/midatlantic/toc.htm>

USDA. February 3, 1999. Executive Order 13112 of February 3, 1999 - Invasive Species. Federal Register: Feb 8, 1999 (Volume 64, Number 25) <http://www.invasivespeciesinfo.gov/laws/execorder.shtml>

USGS (U.S. Geological Survey). January 26, 2017. Nonindigenous Aquatic Species: *Hyla gratiosa*. <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=55>

Van Clef, M. 2009. New Jersey Strategic Management Plan for Invasive Species: The Recommendations of the New Jersey Invasive Species Council to Governor Jon S. Corzine. Pursuant to New Jersey Executive Order #97. <http://www.nj.gov/dep/njisc/docs/Final%20NJ%20Strategic%20Management%20Plan%20for%20Invasive%20Species%2011.09.pdf>

Internet Resources: Biological Resources

Wildlife and Plants

Backyard Habitats & Conservation:

Deer Tolerant/Resistant Native Plants: http://www.bhwp.org/cms/files/file_ID96121.pdf

Gardening for Butterflies: http://www.state.nj.us/dep/fgw/ensp/pdf/literature/butterfly_gardening.pdf

National Audubon Society: http://www.audubon.org/bird/at_home/

New Jersey Audubon Society: <http://www.njaudubon.org/SectionBackyardHabitat/Welcome.aspx>

NJDEP Outdoor Classroom links: <http://www.state.nj.us/dep/seeds/syhart/outclass.htm>

USDA NRCS:

http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/newsroom/features/?cid=nrcs143_023574

Checklists

Birds of NJ: <http://www.state.nj.us/dep/fgw/chkbirds.htm>

Butterflies of NJ: <http://www.naba.org/chapters/nabanj/butterflies.html>

Endangered & Threatened Wildlife of NJ: <http://www.njfishandwildlife.com/tandespp.htm>

Freshwater Fish Of NJ: <http://www.njfishandwildlife.com/chkfish.htm>

Mammals of NJ: <http://www.state.nj.us/dep/fgw/chkmamls.htm>

Reptiles and Amphibians of NJ: http://www.state.nj.us/dep/fgw/ensp/fieldguide_herps.htm

Species of Special Concern of NJ: <http://www.njfishandwildlife.com/ensp/pdf/spclspp.pdf>

Cornel Lab of Ornithology, All About Birds: <http://www.birds.cornell.edu/AllAboutBirds/BirdGuide/>

Native Plants:

Bowman's Hill Wildflower Preserve: <http://www.bhwp.org>

Native Plant Society of NJ: <http://www.npsnj.org/>

USDA Plants Database: <http://plants.usda.gov>

NJDEP:

Conserve Wildlife Foundation of New Jersey: <http://www.conservewildlifenj.org/>
Division of Fish and Wildlife Home Page: <http://www.njfishandwildlife.com/wildlife.htm>
Environmental Rules: http://www.nj.gov/dep/rules/nj_env_law.html
Endangered and Nongame Species Program Home Page: <http://www.state.nj.us/dep/fgw/ensphome.htm>
Landscape Project: <http://www.state.nj.us/dep/fgw/ensp/landscape/>
NJ Wildlife Action Plan: http://www.state.nj.us/dep/fgw/ensp/wap/wap_outline.htm
Rare Plants & Communities: <http://www.state.nj.us/dep/parksandforests/natural/index.html>

North American Butterfly Association, North Jersey Butterfly Club: <http://www.naba.org/chapters/nabanj/>
Long dash (*Polites mystic*): http://www.naba.org/chapters/nabanj/butterflies/long_dash.html

Rare Wildlife Sighting Form: <http://www.njfishandwildlife.com/ensp/rprtform.htm>

Rare Plant Report Form:

http://www.state.nj.us/dep/parksandforests/natural/heritage/natherrareplantspeciesreportform1_2008.doc

Invasive Species

Invasive Species – New Jersey: <http://www.invasivespeciesinfo.gov/unitedstates/nj.shtml>

Native Plant Society of New Jersey – Invasive Species: http://www.npsnj.org/pages/nativeplants_Plant_Lists.html

New Jersey Invasive Species Strike Team (NJISST): <http://www.njisst.org/>

Forest Health: http://www.state.nj.us/dep/parksandforests/forest/njfs_forest_health.html

8: OPEN SPACE & RECREATION

A. Purposes & Funding

The purposes of open space preservation include:

- provide adequate active and passive recreation;
- provide recreational and open space opportunities on an equal and accessible basis for all citizens;
- maintain water quality and groundwater recharge areas;
- protect sensitive environmental features;
- protect historic areas;
- maintain biodiversity;
- minimize erosion or damage from flooding;
- maintain rural character (ANJEC, no date).

Funding for open space comes from a variety of sources, including municipal, county, state and federal sources and private land trusts. Private land trusts are non-profit organizations that “can often act faster and be more creative in their real estate transactions than established government agencies” according to Howe (1989). Landowners are able to reap tax benefits through charitable donations to a land trust. Many successful open space purchases combine a number of funding sources and strategies.

Cape May County assesses a tax which is designated for open space and agricultural land preservation, parks and recreation projects and historic preservation. The rate, not to exceed one cent per \$100 of total County equalized real property valuation, provides approximately \$1.3 million annually (Cape May County Department of Open Space and Farmland Preservation, 2017).

The Garden State Preservation Trust Act provides state funds for land acquisition and park development through the Green Acres program. West Cape May is one of three municipalities in Cape May County that has not yet utilized the program (NJDEP Green Acres Program, January 3, 2017).

Private land trusts working to preserve land in southern New Jersey and the Association of New Jersey Environmental Commissions (ANJEC) are sources for in-depth information concerning open space preservation through various funding, planning, and zoning techniques (see **Internet Resources**).



B. Greenway Establishment & Maintenance

A *greenway* is a corridor of undeveloped land or open space, which often protects environmental features, such as a stream corridor, floodplain, forested ridgeline, or animal migration route, but which can also preserve a scenic view and provide recreational opportunities, such as parks or biking/hiking trails. Greenway corridors also have the potential for positive economic impacts, by creating jobs, enhancing property values, expanding local businesses, attracting new businesses, increasing local tax revenues, decreasing local government

expenditures, and promoting a local community. The publication Economic Impacts of Protecting Rivers, Trails and Greenway Corridors outlines procedures for analyzing economic impacts of a greenway project, and provides examples. Decision makers can benefit from recognition of potential economic impacts as well as intrinsic values of greenways in support of decisions that enhance the well-being of the community (National Park Service, 1995).

Garden State Greenways is an online planning tool designed for all those involved in conserving open space, farmland, and historic areas in New Jersey. It uses GIS to identify *hubs* (larger areas of undeveloped land with important natural resource values) and linear *connectors* between these hubs. The goal of the program is to help coordinate efforts of both private groups and government agencies (NJ Conservation Foundation, 2005).

Local governments often use a variety of planning and zoning techniques for establishing greenways, including creating a greenway map and adopting it as part of the Master Plan, creating a Greenway Overlay District, cluster zoning and Transfer of Development Rights. These strategies can be combined with land preservation, private land trusts, and conservation easements to meet the Borough's open space and recreation goals (Howe, 1989).

Before a greenway is established, issues of maintenance, public access and monitoring of easements must be addressed to ensure long-term success of the project (Howe, 1989).

A significant portion of the Cape Island region at the tip of Cape May County is included in the Garden State Greenways. Large swaths of preserved open space are present adjacent to West Cape May Borough, including Higbee Beach Wildlife Management Area, Cape May Point State Park, and South Cape May Meadows (Garden State

Greenways, 2004). A small portion of the Cape May Meadows Preserve falls within the Borough. West Cape May currently has roughly 130 acres of preserved farmland (West Cape May Sustainability Plan, 2009) that is associated with the greenway. In addition to the land already preserved, the western part of the Borough has a matrix of undeveloped emergent and forested wetland habitat that has not been permanently protected. The New Jersey Green Infrastructure Assessment (NJGIA) has identified a portion of those wetlands as conservation priorities in order to connect West Cape May's preserved farmlands to preserved open spaces within Cape May Point Borough (Garden State Greenways, 2004), and expansion of the greenway is included in West Cape May's long-term plan (Remington, Vernick & Walberg Engineers, 2009).



C. Open Space and Recreation in West Cape May

An updated inventory of the preserved open space and recreation properties within the Borough is presented in **Figure 8a**. Using the acreage figures calculated by GIS, a total of 18.8 acres of Open Space have been established in West Cape May (**Table 8.1**), which is approximately 2.5% the Borough's 755 acres.

Combined totals of preserved open space and farmland yield a collective 151 acres that have been protected in West Cape May, which is 20% of the land in the borough. Plans for the preservation of additional borough lands for greenway expansion and for the development of an eco-park are outlined in West Cape May's Master Plan (Remington, Vernick & Walberg Engineers, 2009).

Table 8.1: Open Space in West Cape May

Block	Lot	Acres*	Owner
33	1	16.58	The Nature Conservancy
49	1.01, 1.02, 1.03	1.76	NJDEP
3	1	0.47	West Cape May
Total:		18.80	
*Acreage calculated by GIS may vary from deed acreage.			

Municipal Open Space

Wilbraham Park and Arboretum is a narrow half acre lot donated to the Borough by John W. Wilbraham in 1921. The site, which is across the street from the historic Wilbraham Mansion, is presently maintained by the West Cape May Shade Tree Commission. The park serves as a small oasis in the downtown section of the borough, offering a beautifully landscaped setting with benches and picnic tables. The location of the park is depicted in **Figure 8b**.

Private Land Trust Open Space

South Cape May Meadows is a Nature Conservancy Preserve over 200 acres in size, and about 16.5 acres of the protected site fall within West Cape May Borough (**Figure 8c**). The land provides foraging and resting habitat for migrating birds, and the preserve's trail system provides hiking and wildlife viewing opportunities to an estimated 90,000 visitors per year (The Nature Conservancy, 2017).

State Open Space

Nearly two acres of state-owned land is located at the southwestern corner of 5th Avenue and State Street in West Cape May (**Figure 8d**). The site is managed as an ancillary parcel of the Higbee Beach Wildlife Management Area (MWA). The primary preserve is about 1,100 acres in size, providing critical habitat for threatened and endangered species and migrating birds. Visitors may enjoy wildlife from two viewing platforms and a network of nature trails (NJDEP Division of Fish & Wildlife, 2016). The remote parcel in West Cape May Borough is undeveloped.

Preserved Farmland

Farmland accounts for about 28% of the total land in West Cape May, with a total of 210.12 acres actively farmed (Cape May County Property Tax Records, 2017). More than half of the farmland in the borough has been preserved (**Table 8.2 and Figure 8a**). The 132 acres of preserved farmland accounts for 17.5% of the land in West Cape May.



Table 8.2: Preserved Farmland in West Cape May.

NJDA/ SADC Number	Preserved as BLOCK	Preserved as LOT	Current Tax ID	Acres*	Date Preserved	Listed Owner
9619	55	25	B 55 L 25	10.329	6/15/2009	Vasser
9630	64	1	B 64 L 1	2.917	4/25/2002	Rea
9632	64	2		1.658		
9633	64	3		0.182		
9634	64	4		1.053		
9624	65	1		2.526		
9626	65	2		1.9		
9628	65	3		0.164		
9629	65	4		0.654		
9622	66	1		0.971		
9631	Borough Hall Alley+			0.362		
9627	Fourth Avenue+			0.947		
9623	Seaview Drive+			0.699		
9625	Sees Alley+			0.385		
9621	73	9		B 73 L 9		
9620	73	10	B 73 L 10	31.28	4/25/2002	Rea
9618	73	10		3.462		
9615	74	2	B 74 L 2	23.739	6/6/2007	Bray Wilde
9616	74	3	B 74 L 3	27.565	4/25/2002	Rea
9617	74	3		9.074		
Total:				132.374		
*Acreage calculated by GIS may vary from deed acreage						
+ indicates undeveloped roadways						




Figure 8a: West Cape May Preserved Open Space and Recreation
 An overview of the 132 acres of Preserved Farmland and 19 acres of Preserved Open Space in West Cape May.


0 0.1 0.2 Miles
 Prepared by Kratzer Environmental Services

Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.



Legend

- West Cape May
- Parcels
- Roads
- Wilbraham Park (WCM Borough)




Prepared by Kratzer Environmental Services

Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.


**Figure 8b: West Cape May
 Wilbraham Park**

A detail map showing the location of the municipal park in the eastern section of the borough



Legend

- West Cape May
- Parcels
- Roads
- South Cape May Meadows Preserve (The Nature Conservancy)



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.


Prepared by Kratzer Environmental Services

**Figure 8c: West Cape May
 South Cape May Meadows Preserve**
A detail map showing the portion of the Nature Conservancy preserve that falls within the borough.



Legend

- Parcels
- Roads
- Water
- State Open Space - Higbee Beach WMA
- Preserved Farmland



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

**Figure 8d: West Cape May
 Part of Higbee Beach WMA**

A detail map showing the location of a parcel that is managed by the State of New Jersey in conjunction with a larger park west of the borough.

References: Open Space

- Association of New Jersey Environmental Commissions. No Date. Resource Paper: Open Space Plan. 12 pages. <http://anjec.org/pdfs/OpenSpacePlan.pdf>
- Cape May County Department of Open Space and Farmland Preservation. 2017. County Website: <http://capemaycountynj.gov/596/Open-Space-Farmland-Preservation>. Accessed March 10, 2017
- Cape May County Property Tax Records. 2017. Cape May County, New Jersey, Free Public Records Directory - Property Search. http://publicrecords.onlinesearches.com/NJ_CapeMay.htm. Accessed March 2017.
- Garden State Greenways. 2004. New Jersey Conservation Foundation. County Maps: Cape May. <http://www.gardenstategreenways.org/documents/capemayGSG.pdf>
- Howe, Linda. 1989. Keeping our Garden State Green: A Local Government Guide for Greenway and Open Space Planning. Association of New Jersey Environmental Commissions. 57 pages.
- National Park Service. 1995. Economic Impacts of Protecting Rivers, Trails, and Greenway Corridors: A Resource Book. Fourth Edition. 154 pages. http://www.nps.gov/pwro/rta/econ_all.pdf
- NJ Conservation Foundation. 2005. Garden State Greenways. <http://www.gardenstategreenways.org>
- NJ Department of Agriculture, SADC. January 8, 2013. New Jersey Farmland Preservation Program (NJFPP). GIS Data. <http://www.nj.gov/agriculture/sadc/farmpreserve/resources/njfpp.zip>
- NJDEP, Division of Fish and Wildlife. 2016. Higbee Beach Wildlife Management Area. <http://www.state.nj.us/dep/fgw/ensp/higbee.htm>
- NJDEP, Green Acres Program. May 27, 2016. NJDEP State Owned, Protected Open Space and Recreation Areas in New Jersey. GIS Data. http://njogis-newjersey.opendata.arcgis.com/datasets/b75fe11ed90543c1b4ee87e66af63b8b_1
- NJDEP Green Acres Program. January 3, 2017. Recreational and Open Space Inventory (ROSI). <http://www.nj.gov/dep/greenacres/openspace.html>. Accessed March 10, 2017.
- Regional Plan Association. 1996. A Region At Risk: A Summary of the Third Regional Plan for the New York-New Jersey-Connecticut Metropolitan Area. 22 pages. <http://library.rpa.org/pdf/RPA-Plan3-A-Region-at-Risk-Summary.pdf>
- Remington, Vernick & Walberg Engineers. 2009. Comprehensive Master Plan Update for the Borough of West Cape May. Prepared for the Borough of West Cape May and Submitted December 30, 2005. http://www.westcapemay.us/LinkClick.aspx?fileticket=UXgh_ymUJcs%3D&tabid=2031
- The Nature Conservancy. 2017. Our Initiatives: New Jersey - South Cape May Meadows. <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/newjersey/placesweprotect/south-cape-may-meadows.xml>
- West Cape May Sustainability Plan. 2009. Prepared by West Cape May Comprehensive Planning Studio, Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey. <http://www.westcapemay.us/LinkClick.aspx?fileticket=15iRI2xnNfl%3D&tabid=2024>

Internet Resources: Open Space

Association of New Jersey Environmental Commissions: <http://anjec.org>

Cape May County Open Space Program Guide: Updated 3/9/2017
<http://capemaycountynj.gov/DocumentCenter/Home/View/3883>

Garden State Greenways <http://www.gardenstategreenways.org>

Native Plant Society of New Jersey: <http://www.npsnj.org>

New Jersey Conservation Foundation: <http://www.njconservation.org/>

New Jersey Green Acres Program: <http://www.nj.gov/dep/greenacres/index.html>

New Jersey Natural Lands Trust: <http://www.njnlt.org/>

NJDEP, Division of Fish and Wildlife. 2016. Higbee Beach Wildlife Management Area.
<http://www.state.nj.us/dep/fgw/ensp/higbee.htm>

NY-NJ Trail Conference: <http://www.nynjtc.org/>

Open Space Institute: http://www.osiny.org/site/PageServer?pagename=Places_NewJersey

Rain Garden Manual: http://www.npsnj.org/pages/nativeplants_Rain_Gardens.html

Rutgers New Jersey Agricultural Experiment Station (NJAES) – information & links for farmers, gardeners, & consumers: <http://njaes.rutgers.edu/>


South Jersey Land and Water Trust: <http://sjlandwater.org/>

The Nature Conservancy. 2017. Our Initiatives: New Jersey - South Cape May Meadows.
<http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/newjersey/placesweprotect/south-cape-may-meadows.xml>

9: HISTORIC RESOURCES

A. Resource Inventory – Historic Resources

The earliest known inhabitants of southern Cape May County were members of the Kechemeche Tribe of the Lenni-Lenape Nation (Stevens, 1897). It is generally believed that the native tribes spent much of the year in the Delaware Valley, utilizing the coastal areas seasonally for hunting and fishing (Skinner and Schrabish, 1913). An archaeological site in West Cape May reportedly produced a number of artifacts, including "A bead beaten from a nugget of native gold" (Skinner and Schrabish, 1913). There is currently a state archaeological grid site (Cell ID BP331) centered in the northeastern corner of West Cape May Borough (see **Figure 9a**) (NJDEP HPO, September 22, 2016a), and Native American artifacts are frequently discovered in neighboring towns on Cape Island (Newsworks, July 20, 2016).



According to Stevens (1897), Cape May was christened in 1623 by the Dutch explorer and Captain Cornelius Jacobsen Mey, who named it after himself. Cape May County was established in 1692 by an Act of the General Assembly of the Province of West New Jersey (Cape May County, 2017). In 1723, the County Court created three precincts that divided the county along both geographic and religious lines: the Upper Precinct (Quaker), Middle Precinct (Baptist) and Lower Precinct (Presbyterian) (Fulginiti, 2017). The three precincts became townships by an act of the New Jersey Legislature in 1798, and West Cape May Borough was formed out of Lower Township in 1884 (Fulginiti, 2017).

B. Historic Districts

The NJDEP Historic Preservation Office (HPO) defines Historic Districts as areas that possess a significant concentration, linkage, or continuity of buildings, sites, structures, or objects united historically or aesthetically by plan or physical development. Historic Districts include National Historic Landmarks; areas on the New Jersey or National Registers of Historic Places; areas determined eligible for inclusion in the registers; districts designated as Local Historic Districts by a local government; or areas that have been identified through a cultural resource survey or other documentation on file at the HPO. Historic Districts in and adjacent to West Cape May are listed in **Table 9.1** and **Table 9.2** and shown in **Figure 9a** (NJDEP HPO, September 22, 2016b).

West Cape May has one region and one building that are listed on the National Register of Historic Places (State of NJ, 2017). The Cape May Historic District includes both the western end of Cape May City and the southeastern portion of West Cape May (see **Table 9.2** and **Figure 9a**). The district, which was added to both the state and national registers in 1970, is best known for its concentration of structures built during the late Victorian Period (1870-1901). The Whilldin-Miller House is located within the Cape May Historic District, but has also been

individually registered since 2002 in the state and 2003 nationally (NJDEP HPO, November 20, 2017) (see **Table 9.2** and **Figure 9a**). The original timber frame portion of the house was reputedly built by Joseph Whilldin around 1715, and the newer part constructed in 1860 (Wikipedia, 2017).

In addition to the registered sites, there is a 1977 opinion of eligibility record for a historic milestone on the south side of Sunset Boulevard in West Cape May. The milestone was a marker for the Cape Island Turnpike, circa 1848-1851. State Historic Preservation Officer (SHOP) opinions are issued in response to federally funded activities that would have an effect on historic properties not listed on the National Register (NJDEP HPO, November 20, 2017).

Table 9.1: Historic Districts In and Adjacent to West Cape May

ID	Name	Status ¹	Date of Status Update	NHL	List Date ²	Period of Significance	Survey Name ³	Survey Date ⁴
474	Cape May Historic District (NR/SR)	LISTED	1970-12-29	NO	NR: 1970-12-29 SR: 1970-12-10	1850-1910		
475	Cape May Historic District (NHL)	NHL	1976-05-11	YES	NR: 1970-12-29 SR: 1970-12-10			
476	West Cape May Historic District	ELIGIBLE	2006-10-26	NO				
643	Cape May Historic District [proposed NR/SR boundary revision, 2013]	LISTED	1970-12-29	NO	NR: 1970-12-29 SR: 1970-12-10	1850-1910	Phase I Archaeological Investigation and Structure Inventory ⁵	2013-01-01
780	Cape May Local Historic District	Locally Designated		NO				
788	Atlantic City Railroad Cape May Division Historic District	ELIGIBLE	2005-06-23	NO			Historic District Management Study ⁶	2010-05-07

Notes:

¹ NHL = National Historic Landmark; LISTED = Listed in the NJ and/or National Registers of Historic Places; ELIGIBLE = Formally determined eligible for listing in the NJ and/or National Registers of Historic Places (SHPO Opinion, COE, DOE); LOCALLY_DESIGNATED = Designated as an historic district by municipal ordinance.

² Title of most recent cultural resource survey report

³ NR = National Register; SR = State Register

⁴ Date of inclusion in most recent cultural resource survey; Either the report date or the individual survey date recorded on the survey form for the resource

⁵ Phase I Archaeological Investigation and Structure Inventory (Revised), Peckman River Flood Damage Reduction Project, Borough of Woodland Park (formerly West Paterson) and Townships of Little Falls and Cedar Grover, Pasasic and Essex Counties, New Jersey

⁶ Historic District Management Study for the Atlantic City Railroad Cape May Division Historic District; Camden, Atlantic, and Cape May Counties, New Jersey

Source: NJDEP HPO, September 22, 2016b

Table 9.2 West Cape May Sites on the National Register of Historic Places.

Data from State of New Jersey (2017), Locations from NJ GeoWeb (2017).

Cape May Historic District (added 1970 - #70000383)	
Cape May City and Borough of West Cape May, Cape May	
Historic Significance:	Event, Architecture/ Engineering
Architect, builder or engineer:	Multiple
Architectural Style:	Late Victorian
Area of Significance:	Conservation, Entertainment/ Recreation, Architecture
Period of Significance	1850-1875, 1875-1899, 1900-1924
Owner:	Private, Local
Historic Function:	Domestic, Health Care
Historic Sub-function:	Camp, Hotel, Resort
Current Function:	Domestic, Health Care
Current Sub-function:	Camp, Hotel, Resort
Whilldin-Miller House , also known as Fow House (added 2003 - #03000012)	
416 South Broadway, Borough of West Cape May, Cape May	
Historic Significance:	Architecture/ Engineering
Architectural Style:	Mid 19th Century Revival, Italianate
Area of Significance:	Architecture
Period of Significance	1850-1874
Owner:	Private
Historic Function:	Domestic
Historic Sub-function:	Single Dwelling
Current Function:	Commerce/Trade
Current Sub-function:	Restaurant
Source: NJDEP HPO, November 20, 2017	




Figure 9a: West Cape May Archeological and Historical Resources

The southeast corner of West Cape May includes a part of the larger Cape May Historic District, listed in 1970, as well as the West Cape May Historic District which was designated as Eligible for Listing in 2006. A part of the Atlantic City Railroad Cape May Division Historic District occupies the northeast corner of the borough. Inset: The northeast end of West Cape May is also included in a state archaeological project grid.

Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

0 0.25 0.5 Miles
 Prepared by Kratzer Environmental Services

C. Historic Properties

According to the NJDEP HPO, Historic Properties are buildings, sites, structures or objects that are evaluated as historically significant. These include properties that are on the National Historic Landmarks; included in the state or National Registers of Historic Places; determined Eligible for inclusion in the registers through the state or federal HPO processes; designated as Local Landmarks by local government; or identified through cultural resource survey or other documentation on file at the HPO. Properties on the current HPO inventory are listed in **Table 9.3** and shown on **Figure 9b** (NJDEP HPO, February 21, 2017a; NJDEP, February 21, 2017b).

Table 9.3: Historic Properties in West Cape May

ID	Name/Address	Status	Survey Name
Historic District: Atlantic City Railroad Cape May Division Historic District			
69113	Cape Island Creek Trestle Atlantic City Railroad, Cape May Division over Cape Island Creek; Atlantic City Railroad, Cape May Division over Cape Island Creek	ELIGIBLE_HD	Historic District Management Study for the Atlantic City Railroad Cape May Division Historic District; Camden, Atlantic, and Cape May Counties, New Jersey (2010-05-07)
90953	Cape Island Creek Trestle Atlantic City Railroad, Cape May Division over Cape Island Creek	ELIGIBLE_HD	Historic District Management Study for the Atlantic City Railroad Cape May Division Historic District; Camden, Atlantic, and Cape May Counties, New Jersey (2010-05-07)
Historic District: Cape May Historic District (NR/SR)			
511	Whilldin-Miller House 416 South Broadway	LISTED_INDV 2003-02-12	A Survey of the Heavy Timber Frame Buildings of Cape May County, New Jersey (2003-10-01)
77706	109 SECOND AVENUE	LISTED_HD	
77707	201 BROADWAY	LISTED_HD	
77708	141 PEARL AVENUE	LISTED_HD	
77709	105 W. GRANT ST.	LISTED_HD	
77710	196 SOUTH BROADWAY	LISTED_HD	
77711	100 GRANT STREET	LISTED_HD	
77712	110 ELDREDGE AVE	LISTED_HD	
77713	133 ELDREDGE AVE	LISTED_HD	
77714	402 BROADWAY	LISTED_HD	
77715	121 SIXTH AVENUE	LISTED_HD	
77716	4 1/2 E. MECHANIC ST.	LISTED_HD	
77717	156 LEAMING AVENUE	LISTED_HD	
77718	SUNSET BLVD.&1ST AVE	LISTED_HD	
77719	FIRST AVENUE	LISTED_HD	
77720	222 BROADWAY	LISTED_HD	
77721	113 PEARL AVE	LISTED_HD	
77722	301 BROADWAY	LISTED_HD	
77723	117 PEARL AVE	LISTED_HD	
77724	121 PEARL AVE	LISTED_HD	
77725	127 GOLDBEATEN ALLEY	LISTED_HD	
77726	125 PEARL AVE	LISTED_HD	
77727	129 PEARL AVE	LISTED_HD	
77728	135 PEARL AVE	LISTED_HD	
77729	220 BROADWAY	LISTED_HD	
79409	137 ELDREDGE AVE	LISTED_HD	
79410	129 ELDREDGE AVE	LISTED_HD	
79411	131 EMERALD AVE	LISTED_HD	
79412	117 EMERALD AVE	LISTED_HD	
79413	123 EMERALD AVE	LISTED_HD	
79414	135 EMERALD AVE	LISTED_HD	
79415	409 BROADWAY	LISTED_HD	
79416	131 THIRD AVENUE	LISTED_HD	
79417	EMERALD AVE	LISTED_HD	
79418	414 PARK BLVD	LISTED_HD	
79419	137 EMERALD AVE	LISTED_HD	

79420	124 THIRD AVENUE	LISTED_HD	
79421	605 PARK BLVD	LISTED_HD	
79422	9 E MECHANIC ST	LISTED_HD	
79423	409 PACIFIC AVE	LISTED_HD	
79424	16 E MECHANIC ST	LISTED_HD	
79425	112 PEARL AVE	LISTED_HD	
79426	114 PEARL AVE	LISTED_HD	
79427	116 PEARL AVE	LISTED_HD	
79428	120 PEARL AVE	LISTED_HD	
79429	122 PEARL AVE	LISTED_HD	
79430	124 PEARL AVE	LISTED_HD	
79431	128 PEARL AVE	LISTED_HD	
79432	111 SECOND AVENUE	LISTED_HD	
79433	132 PEARL AVE	LISTED_HD	
79434	136 PEARL AVE	LISTED_HD	
79435	140 PEARL AVE	LISTED_HD	
79585	114 SIXTH AVENUE	LISTED_HD	
79586	141 ELDREDGE AVE	LISTED_HD	
79587	126 LEAMING AVENUE	LISTED_HD	
79588	124 FOURTH AVENUE	LISTED_HD	
79589	122 ELDREDGE AVE	LISTED_HD	
79590	120 FOURTH AVENUE	LISTED_HD	
79591	423 BROADWAY	LISTED_HD	
79592	310 BROADWAY	LISTED_HD	
79593	420 PARK BLVD	LISTED_HD	
79594	411-413 BROADWAY	LISTED_HD	
79595	125 EMERALD AVE	LISTED_HD	
79596	310 PACIFIC AVE	LISTED_HD	
79597	316 BROADWAY	LISTED_HD	
79598	125 THIRD AVENUE	LISTED_HD	
79599	404 FOW AVE.	LISTED_HD	
79600	407 FIRST AVENUE	LISTED_HD	
79601	400 FIRST AVENUE	LISTED_HD	
79602	400 FOW AVENUE	LISTED_HD	
79603	217 MYRTLE AVE	LISTED_HD	
79604	324 FIRST AVENUE	LISTED_HD	
79605	325 FIRST AVENUE	LISTED_HD	
79606	6 CONGRESS ST	LISTED_HD	
79607	402, 404, 586-604 MYRTLE AVE	LISTED_HD	
79608	400 MYRTLE AVE	LISTED_HD	
79609	320 FIRST AVENUE	LISTED_HD	
79610	398 MYRTLE AVE	LISTED_HD	
79611	396 MYRTLE AVE	LISTED_HD	
79612	314 FOW AVE	LISTED_HD	
79613	318 FIRST AVENUE	LISTED_HD	
79614	317 FIRST AVENUE	LISTED_HD	
79615	316 FIRST AVENUE	LISTED_HD	
79616	313 FOW AVE	LISTED_HD	
79617	310 FIRST AVENUE	LISTED_HD	
79672	309 FIRST AVENUE	LISTED_HD	
79673	315 FOW AVENUE	LISTED_HD	
79674	312 FOW AVE	LISTED_HD	
79675	209 BROADWAY	LISTED_HD	
79676	216 BROADWAY	LISTED_HD	
79677	113 SECOND AVENUE	LISTED_HD	
79678	132 YORK AVE	LISTED_HD	
79679	111-117 SUNSET BLVD	LISTED_HD	
79680	119 SIXTH AVENUE	LISTED_HD	
79681	115 YORK AVE	LISTED_HD	
79682	103 SECOND AVENUE	LISTED_HD	

79683	204 BROADWAY	LISTED_HD	
79684	129 YORK AVE	LISTED_HD	
79685	116 SECOND AVENUE	LISTED_HD	
79686	133 YORK AVE	LISTED_HD	
79687	139 YORK AVE	LISTED_HD	
79688	112-114 SECOND AVENUE	LISTED_HD	
79689	139 BROADWAY	LISTED_HD	
79690	131 BROADWAY	LISTED_HD	
79691	110 SECOND AVE	LISTED_HD	
79692	123 PACIFIC AVE.	LISTED_HD	
79693	110 YORK AVE	LISTED_HD	
79694	108 SECOND AVENUE	LISTED_HD	
79695	112 YORK AVE	LISTED_HD	
79696	106 SECOND AVENUE	LISTED_HD	
79697	114 YORK AVE	LISTED_HD	
79698	116 YORK AVE	LISTED_HD	
79699	104 SECOND AVENUE	LISTED_HD	
79700	124 YORK AVE	LISTED_HD	
79701	114 FOURTH AVENUE	LISTED_HD	
79702	615 BROADWAY	LISTED_HD	
79703	122 BROADWAY	LISTED_HD	
79704	127 BROADWAY	LISTED_HD	
79705	MYRTLE AVE & PARK	LISTED_HD	
79782	128 YORK AVE	LISTED_HD	
79783	111 MYRTLE AVE	LISTED_HD	
79784	110 BROADWAY	LISTED_HD	
79785	126 SUNSET BLVD.	LISTED_HD	
79786	BROADWAY & SUNSET BLVD	LISTED_HD	
79787	MYRTLE AVE & PARK BLVD	LISTED_HD	
79788	118 SIXTH AVENUE	LISTED_HD	
79789	132 LEAMING AVE	LISTED_HD	
79790	605 BROADWAY	LISTED_HD	
79791	123 E MECHANIC STREET	LISTED_HD	
79792	308 PACIFIC AVENUE	LISTED_HD	
79793	312 PACIFIC AVENUE	LISTED_HD	
79794	608-616 MYRTLE AVENUE	LISTED_HD	
79795	15-17 PARK BLVD	LISTED_HD	
79796	121-125 YORK AVE	LISTED_HD	
79797	127 YORK AVE	LISTED_HD	
79798	151-153 YORK AVE	LISTED_HD	
79799	211-213 PARK BLVD	LISTED_HD	
79800	305 PARK BLVD	LISTED_HD	
79801	313 PARK BLVD	LISTED_HD	
79802	114 ELDREDGE AVE	LISTED_HD	
79803	115-117 ELDREDGE AVE	LISTED_HD	
79804	119-121 ELDREDGE AVENUE	LISTED_HD	
79805	EAST MECHANIC ST.	LISTED_HD	
79806	120 LEAMING AVE UNIT A & B	LISTED_HD	
79807	107 PARK BLVD	LISTED_HD	
79808	145 YORK AVE	LISTED_HD	
79809	SIXTH AVENUE	LISTED_HD	
79810	4 LANDIS AVE	LISTED_HD	
79811	620 BROADWAY	LISTED_HD	
79812	8 LANDIS AVE	LISTED_HD	
79813	6 LANDIS AVE	LISTED_HD	
79814	2 LANDIS AVE	LISTED_HD	
79895	616 BROADWAY	LISTED_HD	
79896	626 BROADWAY	LISTED_HD	
79897	135 SIXTH AVE	LISTED_HD	
79898	619 BROADWAY	LISTED_HD	

79899	133 SIXTH AVE	LISTED_HD	
79900	127 SIXTH AVENUE	LISTED_HD	
80633	105-107 SECOND AVENUE	LISTED_HD	
80634	102 GOLDBEATEN ALLEY	LISTED_HD	
80635	107 YORK AVE	LISTED_HD	
80636	212 BROADWAY	LISTED_HD	
80637	111 YORK AVE	LISTED_HD	
80638	304 FIRST AVENUE	LISTED_HD	
80639	422 BROADWAY	LISTED_HD	
80640	172 LEAMING AVE	LISTED_HD	
80641	5 E. MECHANIC ST	LISTED_HD	
80642	307 BROADWAY	LISTED_HD	
80643	120 THIRD AVENUE	LISTED_HD	
80644	145 ELDREDGE AVE	LISTED_HD	
80645	125 ELDREDGE AVE	LISTED_HD	
80646	129 THIRD AVENUE	LISTED_HD	
80647	127 THIRD AVENUE	LISTED_HD	
80648	136 LEAMING AVE	LISTED_HD	
80649	108-110 SIXTH AVENUE	LISTED_HD	
80650	140 LEAMING AVENUE	LISTED_HD	
80651	108 FOURTH AVENUE	LISTED_HD	
80652	Thomas Eldredge House 417 BROADWAY	LISTED_HD 1970-12-29	A survey of the Heavy Timber Frame Buildings of Cape May - Phase II (2006-11-01)
80726	324 BROADWAY	LISTED_HD	
80727	134 THIRD AVE	LISTED_HD	
80728	130 EMERALD AVE	LISTED_HD	
80729	170 LEAMING AVENUE	LISTED_HD	
80730	300 FIRST AVENUE	LISTED_HD	
80731	FOW AVE.	LISTED_HD	
80732	FIRST AVE & GRANT ST.	LISTED_HD	
80733	302 SOUTH BROADWAY	LISTED_HD	
80734	119 MYRTLE AVE	LISTED_HD	
80735	123 BROADWAY	LISTED_HD	
80736	136 YORK AVE	LISTED_HD	
80737	120 BROADWAY	LISTED_HD	
80738	140 YORK AVE	LISTED_HD	
80739	103-109 SUNSET BLVD	LISTED_HD	
80740	121 BROADWAY	LISTED_HD	
80741	116 BROADWAY	LISTED_HD	
80742	113 MYRTLE AVE	LISTED_HD	
80743	133 MYRTLE AVE	LISTED_HD	
80744	119 THIRD AVE	LISTED_HD	
80745	109 EMERALD AVE	LISTED_HD	
80746	117 THIRD AVE	LISTED_HD	
80747	135 THIRD AVENUE	LISTED_HD	
80748	115 EMERALD AVE	LISTED_HD	
80749	115 THIRD AVENUE	LISTED_HD	
80750	116 THIRD AVENUE	LISTED_HD	
80751	121 E MECHANIC ST	LISTED_HD	
80752	112 THIRD AVENUE	LISTED_HD	
80753	306 PARK BLVD	LISTED_HD	
80754	207 W. GRANT ST.	LISTED_HD	
80755	201 W GRANT ST	LISTED_HD	
80756	115 W GRANT STREET	LISTED_HD	
80757	136 EMERALD AVE	LISTED_HD	
80758	121 THIRD AVENUE	LISTED_HD	
80820	407 BROADWAY	LISTED_HD	
80821	162 LEAMING AVENUE	LISTED_HD	
80822	138 EMERALD AVE	LISTED_HD	
80823	126 THIRD AVE	LISTED_HD	

80824	144 EMERALD AVE	LISTED_HD	
80825	146 EMERALD AVE	LISTED_HD	
80826	309 PARK BLVD	LISTED_HD	
80827	515-517 BROADWAY	LISTED_HD	
80828	3 E MECHANIC ST	LISTED_HD	
80829	12 E MECHANIC ST	LISTED_HD	
80830	11 E. MECHANIC ST.	LISTED_HD	
80831	SIXTH AVENUE	LISTED_HD	
80832	109 SIXTH AVE	LISTED_HD	
80833	102 SECOND AVENUE	LISTED_HD	
80834	401 BROADWAY	LISTED_HD	
80835	149 EMERALD AVENUE	LISTED_HD	
80836	FIRST AVENUE	LISTED_HD	
80837	306 FOW AVE.	LISTED_HD	
80838	311 FOW AVE	LISTED_HD	
80839	301 FOW AVE	LISTED_HD	
80840	298 SOUTH BROADWAY	LISTED_HD	
80841	298 1/2 BROADWAY	LISTED_HD	
80842	296 SOUTH BROADWAY	LISTED_HD	
80843	109 W. GRANT ST.	LISTED_HD	
80844	107 W GRANT ST	LISTED_HD	
80845	113 ELDREDGE AVE	LISTED_HD	
80846	511 BROADWAY	LISTED_HD	
80847	603 PARK BLVD	LISTED_HD	
80848	201 PACIFIC AVE	LISTED_HD	
80849	107 GOLDBEATEN ALLEY	LISTED_HD	
80850	115 SECOND AVENUE	LISTED_HD	
80851	136-138 ELDREDGE AVE	LISTED_HD	
80852	142 ELDREDGE AVE	LISTED_HD	
80915	144 ELDREDGE AVE	LISTED_HD	
80916	154 LEAMING AVENUE	LISTED_HD	
80917	501-503 PACIFIC AVE.	LISTED_HD	
80918	106-108 SUNSET BLVD	LISTED_HD	
80919	409 FIRST AVENUE	LISTED_HD	
80920	6 E MECHANIC ST	LISTED_HD	
80921	523 BROADWAY	LISTED_HD	
80922	500-502 BROADWAY	LISTED_HD	
80923	158 LEAMING AVENUE	LISTED_HD	
80924	8 E MECHANIC ST	LISTED_HD	
80925	160 LEAMING AVE	LISTED_HD	
80926	10 E MECHANIC	LISTED_HD	
80927	521 BROADWAY	LISTED_HD	
80928	412 PARK BLVD	LISTED_HD	
80929	306 BROADWAY	LISTED_HD	
80930	147 EMERALD AVE	LISTED_HD	
80931	104 EMERALD AVENUE	LISTED_HD	
80932	302 BROADWAY	LISTED_HD	
80933	110 EMERALD AVE	LISTED_HD	
80934	116 EMERALD AVE	LISTED_HD	
80935	136 THIRD AVENUE	LISTED_HD	
80936	118 EMERALD AVE	LISTED_HD	
80937	122 EMERALD AVE	LISTED_HD	
80938	311 BROADWAY	LISTED_HD	
80939	126 EMERALD AVE	LISTED_HD	
80940	612 BROADWAY	LISTED_HD	
80941	115 BROADWAY	LISTED_HD	
80942	101 SUNSET BLVD	LISTED_HD	
80943	127 MYRTLE AVE	LISTED_HD	
80944	601 BROADWAY	LISTED_HD	
80945	506 BROADWAY	LISTED_HD	

80946	150 LEAMING AVE	LISTED_HD	
80947	2 E MECHANIC STREET	LISTED_HD	
80948	4 E MECHANIC ST	LISTED_HD	
81012	406 BROADWAY	LISTED_HD	
81013	503 BROADWAY	LISTED_HD	
81014	[Park] MYRTLE & PERRY AVE	LISTED_HD	
81015	304 SOUTH BROADWAY	LISTED_HD	
81016	306 SOUTH BROADWAY	LISTED_HD	
81017	102-104 SUNSET BLVD	LISTED_HD	
81018	479 W PERRY ST	LISTED_HD	
81019	435 W PERRY ST	LISTED_HD	
81020	433 W PERRY ST	LISTED_HD	
81021	431 W PERRY ST	LISTED_HD	
81022	429 W PERRY ST	LISTED_HD	
81023	427 W PERRY ST	LISTED_HD	
81024	310 SO. BROADWAY	LISTED_HD	
81025	392 MYRTLE AVE	LISTED_HD	
81026	425 W PERRY ST	LISTED_HD	
81027	423 W PERRY ST	LISTED_HD	
81028	421 W PERRY ST	LISTED_HD	
81029	419 W PERRY ST	LISTED_HD	
81030	405 W PERRY ST	LISTED_HD	
81031	394 MYRTLE AVE	LISTED_HD	
81032	308 SOUTH BROADWAY	LISTED_HD	
81033	116 FOURTH AVENUE	LISTED_HD	
81034	128 ELDREDGE AVE	LISTED_HD	
81035	185 E MECHANIC ST	LISTED_HD	
81036	414-416 BROADWAY	LISTED_HD	
81037	123 FOURTH AVENUE	LISTED_HD	
81038	PARK BLVD	LISTED_HD	
81039	507 BROADWAY	LISTED_HD	
81040	121 FOURTH AVENUE	LISTED_HD	
81041	17 E MECHANIC ST	LISTED_HD	
81042	119 FOURTH AVE	LISTED_HD	
81043	601 PARK BLVD	LISTED_HD	
81044	E MECHANIC ST	LISTED_HD	
Other/No District			
93209	609 Sunset Boulevard (demolished)	IDENTIFIED_INDV 1980-01-01	Historic Sites Survey, Cape May County, 1980: Lower Township (1980-01-01)
93350	601 SUNSET BLVD (demolished)	IDENTIFIED_INDV 1980-01-01	Historic Sites Survey, Cape May County, 1980: Lower Township (1980-01-01)
93905	Ezekiel Eldredge House (alternate name George Reeves House) 108 Stevens Street	IDENTIFIED_INDV 2006-11-01	A survey of the Heavy Timber Frame Buildings of Cape May - Phase II (2006-10-12)
93943	Whilldin-Eldredge House (alternate name Stillwell Stevens House) 213 STEVENS ST	IDENTIFIED_INDV 2006-11-01	A survey of the Heavy Timber Frame Buildings of Cape May - Phase II (2006-11-01)
Source: NJDEP HPO, February 21, 2017a; NJDEP, February 21, 2017b			





Figure 9b: West Cape May Historic Districts and Properties
The four homes highlighted on the map were featured in a 2006 survey of the Heavy Timber Frame Buildings of Cape May.



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

References: Historic Resources

Cape May County Website. 2017. History of Cape May County. <http://capemaycountynj.gov/982/History-of-Cape-May-County>. Accessed March 2017.

Fulginiti, R.M. 2017. From the Beginning: Fun Facts about Early Cape May Government. <http://capemaycountynj.gov/DocumentCenter/View/2332>. Accessed March 2017.

Newsworks. July 20, 2016. Cape May becoming a popular spot to find Native American artifacts. <http://www.newsworks.org/index.php/local/new-jersey/95503-cape-may-becoming-a-popular-spot-to-find-native-american-artifacts?l=me>

NJDEP Historic Preservation Office (HPO). November 20, 2017. New Jersey Department of Environmental Protection - Historic Preservation Office. New Jersey and National Registers of Historic Places: Cape May County. http://www.nj.gov/dep/hpo/1identify/nrsr_lists/Cape%20May.pdf. Accessed November 24, 2017.

NJDEP Historic Preservation Office (HPO). February 21, 2017a. Historic Properties of New Jersey, Edition 20170221 (Land_use_HPO_property). http://njogis-newjersey.opendata.arcgis.com/datasets/4d1c7b7379ce4de0816de81aeef293bb_7

NJDEP Historic Preservation Office (HPO). February 21, 2017b. Historic Property Features of New Jersey, Edition 20170221 (Land_use_HPO_property_feature). http://njogis-newjersey.opendata.arcgis.com/datasets/86fbaa4d2c8147e2b2494f1cb18a6e9e_6

NJDEP Historic Preservation Office (HPO). September 22, 2016a. New Jersey Department of Environmental Protection (NJDEP). Archaeological Site Grid of New Jersey, Edition 20160922 (Land use HPO arch grid). http://njogis-newjersey.opendata.arcgis.com/datasets/a6ade30b86fe465fa319c4a7b8aac175_8

NJDEP Historic Preservation Office (HPO). September 22, 2016b. Historic Districts of New Jersey, Edition 20160922 (Land use HPO district). http://njogis-newjersey.opendata.arcgis.com/datasets/53861c4495914d2381794948adef58e0_9

Skinner, A. and M. Schrabish. 1913. Bulletin 9, A Preliminary Report of the Archaeology Survey of the State of New Jersey. Made by the Department of Anthropology in the American Museum of Natural History, Under the Direction of the State Geological Survey. MacCrellish & Quigley, State Printers, Trenton, New Jersey.

State of New Jersey. 2017.

Stevens, L.T. 1897. The History of Cape May County New Jersey: From the Aboriginal Times to the Present Day. Press of the Star of the Cape Publishing Company, Cape May, New Jersey.

Wikipedia. Accessed March 20, 2017. Whilldin-Miller House. https://en.wikipedia.org/wiki/Whilldin-Miller_House

Internet Resources: Historic Resources

Ancient America: Learning Lenape: http://www.njskylands.com/hs_lenape_083.htm

Cape May County Library: Genealogy and Local History - <http://www.cmclibrary.org/services/genealogy-local-history>

Cape May County New Jersey: Historic Sites - <http://capemaycountynj.gov/374/Historic-Sites>

History of Cape May County: <http://capemaycountynj.gov/982/History-of-Cape-May-County>

New Jersey National Landmarks and Historic Sites: <http://www.nationalregisterofhistoricplaces.com/nj/cape+may/state.html>

New Jersey Comprehensive Statewide Historic Preservation Plan, 2013-2019:

http://www.nj.gov/dep/hpo/Index_HomePage_images_links/hpo_plan%202013_2019/hpoplan2014.pdf

The Museum of Cape May County - 504 North Route 9, Cape May Court House - <http://www.cmcmuseum.org/>

Revolutionary War Sites in Cape May County:

http://www.revolutionarywarnewjersey.com/new_jersey_revolutionary_war_sites/counties/cape_may_county_revolutionary_war_sites.htm

10: REGIONAL RELATIONSHIPS

A. Sustainable Jersey

According to the organization’s website, “Sustainable Jersey is a nonprofit organization that provides tools, training and financial incentives to support communities as they pursue sustainability programs. By supporting community efforts to reduce waste, cut greenhouse gas emissions, and improve environmental equity, Sustainable Jersey is empowering communities to build a better world for future generations.” (Sustainable Jersey, 2017a)

The voluntary Sustainable Jersey certification is a significant achievement for municipal governments in New Jersey. Municipalities are awarded points for completing and documenting actions that increase sustainability. Nearly 80% of New Jersey’s municipalities are listed as participating in the program, while 44% of these are currently certified at either the Bronze or Silver level.



West Cape May Borough received a *Sustainable Jersey Community Bronze Certification* on October 12, 2016 with 185 points (see **Table 10.1**) (Sustainable Jersey, 2017b).

There are currently two solar photovoltaic installations on public buildings in West Cape May, including a 7.56 mw system on the Borough Hall (installed in 2008) and a 66.89 mw system on the elementary school (installed in 2012) (NJDEP, January 20, 2017).

Table 10.1: Sustainability Actions Implemented in West Cape May for Bronze Certification

Category	Action	Points	Comment
Community Partnership & Outreach	Community Education and Outreach	10	
Community Partnership & Outreach	Create Green Team	10	Bronze Mandatory
Support Local Food	Farmers Market	10	
Support Local Food	Making Farmers Markets Accessible	5	
Innovation & Demonstration Projects	Innovative Community Project (Arbor Day activities)	10	
Innovation & Demonstration Projects	Wind Ordinance	10	
Solar Actions	On-site solar system	10	
Land Use & Transportation	Sustainable Land Use Pledge	20	Bronze Priority Silver Priority
Natural Resources	Environmental Commission	10	
Natural Resources	Environmental Commission Site Plan Review	10	
Natural Resources Protection Ordinances	Tree Protection Ordinance	10	
Tree and Woodlands Management	Community Forestry Plan and Tree Cover Goal	20	
Tree and Woodlands Management	Tree Planting Programs	10	
Green Purchasing Program	Recycled paper	10	
Access to Public Information	Digitizing Public Information	10	
Waste Management	Prescription Drug Safety and Disposal	10	Bronze Priority Silver Priority
Recycling	Recycling Depot	10	
	Total:	185	
Source: Sustainable Jersey, 2017b.			

B. Water Supply Planning

The goal of statewide water supply planning, mandated by the Water Supply Management Act (N.J.S.A. 58:1A-1), is to improve the management and protection of the State's water supplies to ensure that the State's water supplies could withstand foreseeable drought and that aquifers are not depleted.

The first New Jersey Water Supply Plan (NJSWSP) was adopted in 1982, and has been periodically updated and revised. A new plan was released for public comment in June 2017 and finalized in October. The goal of this 5 year (2017-2022) NJSWSP is "to form the foundation of a "living" resource able to be updated on a continuous basis as reliable new data becomes available and improved upon as new scientific methods are identified." (NJDEP Division of Water Supply and Geoscience, October 5, 2017)

Appendix A of the plan presents a characterization of confined and unconfined ground water and surface water supplies on a HUC11 basis. DEP analyzed data available from 1998 through 2015 to determine a period of time representing peak consumptive demand in New Jersey. West Cape May is within two HUC11s; 02040206230 (Cape May Tribs West) and 02040302080 (Cape May Bays & Tribs East). The method calculated the quantity of water available for depletive/consumptive loss to be 4.2 million gallons per day (mgd) in these two HUC11s. The full allocation of depletive/consumptive water loss is calculated to be 1.9 mgd in Cape May Tribs West and -1.9 mgd (net gain) in Cape May Bays & Tribs East. Based on this information, one of the stated management options is that NJDEP will continue to monitor the Cape May Tribs West watershed as they approach the planning threshold for depletive/consumptive uses associated with unconfined groundwater or unregulated (non-safe yield) surface water (NJDEP Division of Water Supply and Geoscience, October 5, 2017).

D. Water Quality Management Planning

In 2015, NJDEP released a new Continuing Planning Process (CPP) document, which was prepared pursuant to the federal Clean Water Act (CWA) and the New Jersey Water Pollution Control Act (WQPA), both of which require the NJDEP to formulate a continuing planning process (CPP) to achieve the water quality standards and maintain, improve, and protect water quality throughout the State. The CPP is intended to serve as an easily accessible planning tool, to be used not only as a listing of current NJDEP programs and rules relating to water quality, but as a resource for planning entities and members of the public on current policies and technical guidance on water quality issues, including:

- Establishing water quality standards and goals
- Assessing water quality and identify priority problems
- Water Quality Management Planning
- Identifying and controlling sources and causes of water quality impairment
- Intergovernmental Coordination (NJDEP Water Resources Management, December 18, 2015).

The *Water Quality Management Planning* rules at N.J.A.C 7:15 represent one component of the CPP. The current rules were adopted November 7, 2016, repealing and replacing the prior rules from 2008. The new rules streamline the wastewater planning process and eliminate the some of the requirements found previously in both wastewater planning and other permitting

programs. These rules focus on procedures for adopting new or amended areawide water quality management (WQM) plans, including Wastewater Management Plans (WMPs); Lists of water quality limited (impaired) waters; and total maximum daily loads (TMDL) for impaired waters. The CPP describes how these processes, along with other Department programs, integrate and unify water quality management planning processes, establish and assess attainment of water quality goals and standards, and implement control measures necessary to maintain, improve, and protect water quality throughout the State (NJDEP Water Resources Management, December 18, 2015; NJDEP, November 7, 2016).

A *Wastewater management planning agency* or *WMP agency* is defined in the rule as a governmental entity that has wastewater management planning responsibility (NJDEP, November 7, 2016). The Cape May County Board of Chosen Freeholders is the designated planning agency with approved plan amendment procedures for the Cape May County WQM planning area, which includes the Borough of West Cape May. One of the WQM agency's roles is to update the WMP at least once every 10 years (NJDEP Office of Water Resources Management Coordination, November 16, 2017).

There have been no recent changes to the West Cape May portion of the Cape May County WQM planning area (NJDEP Office of Water Resources Management Coordination, November 3, 2017). The last change was a 2006 revision in order to correct, clarify or update the boundaries using newer, more accurate mapping (NJDEP Division of Watershed Management, July 11, 2006; NJDEP, January 24, 2017) (see **Figure 2d**).

The wastewater from the Cape May Co MUA discharges into Atlantic Ocean via an outfall pipe in Cape May Point Borough (discharge id. number NJ0020371) (NJDEP, February 13, 2017) (see **Figure 2d**).

References: Regional Relationships

Sustainable Jersey

NJDEP. January 20, 2017. Public Solar Facilities of New Jersey, Edition 20170120 (Util_solar_public). Data obtained from AEG & NJBPU on 20160826. GIS data.

Sustainable Jersey. 2017a. Sustainable Jersey. <http://www.sustainablejersey.com/>. Accessed September 4, 2017.

Sustainable Jersey. 2017b. West Cape May Sustainable Jersey Community Certification Report. [HTML version of report](#) or [PDF version of report](#). Accessed October 26, 2017.

Water Supply Planning

NJDEP Division of Water Supply and Geoscience. October 5, 2017. New Jersey Water Supply Plan 2017-2022. <http://www.state.nj.us/dep/watersupply/wsp.html>

Water Quality Management Planning

NJDEP. February 13, 2017. NJPDES Surface Water Discharges in New Jersey, (1:12,000) Edition 20170213 (Strc_NJPDES_sw_pipe). GIS data. http://njogis-newjersey.opendata.arcgis.com/datasets/2ceba1ef852b4940afc3e0d94fb5d327_6

NJDEP. January 24, 2017. Statewide Sewer Service Area for New Jersey, Edition 20170124 (Util_wastewater_servicearea). GIS data. http://njogis-newjersey.opendata.arcgis.com/datasets/2ceba1ef852b4940afc3e0d94fb5d327_6

NJDEP. November 7, 2016. N.J.A.C. 7:15: Water Quality Management Planning. Date last amended: November 7, 2016. http://www.nj.gov/dep/rules/rules/njac7_15.pdf

NJDEP Division of Watershed Management. July 11, 2006. Public Notice: Revision to the Cape May County Water Quality Management Plan. <http://www.nj.gov/dep/wqmp/docs/wqmp/capemay/20060711a.pdf>

NJDEP Office of Water Resources Management Coordination. November 16, 2017. The Water Quality Management Rules Home Page. <http://www.nj.gov/dep/wqmp/wqmmps.html>. Accessed November 19, 2017.

NJDEP Office of Water Resources Management Coordination. November 3, 2017. Water Quality Management Planning Program. <http://www.nj.gov/dep/wqmp/wmpadopted.html#capemay>. Accessed November 19, 2017.

NJDEP Water Resources Management. December 18, 2015. New Jersey's Continuing Planning Process. <http://www.nj.gov/dep/wrm/docs/cpp.pdf>.

Internet Resources: Regional Relationships

Sustainability

NJDEP Office of Sustainability: <http://www.nj.gov/dep/aqes/sustainability.html>

Rethink Energy NJ: <http://rethinkenergynj.org/>

Sustainable Jersey: <http://www.sustainablejersey.com/>

USEPA Greener Living: <https://www.epa.gov/environmental-topics/greener-living>

Water Quality Management Planning

NJDEP Water Quality Management Planning: <http://www.nj.gov/dep/wrm/index.html>

NJDEP Water Supply Planning: <http://www.state.nj.us/dep/watersupply/wsp.html>

11: COMPOSITE MAP OF ENVIRONMENTALLY CRITICAL AREAS

Throughout this document, many environmental and natural features of the Borough of West Cape May have been documented, described and mapped. One of the greatest values of mapping with GIS is to easily combine features in new ways. To accomplish this, **Figure 11** combines some of the mapped layers from previous sections, displaying features that make an area environmentally critical together on one map.

A useful definition of an "environmentally critical area" is provided in the Stormwater Management regulations (N.J.A.C. 7:8):

" 'Environmentally critical area' means an area or feature which is of significant environmental value, including, but not limited to: stream corridors; natural heritage priority sites; habitats of endangered or threatened species; large areas of contiguous open space or upland forest; steep slopes; and well head protection and groundwater recharge areas. Habitats of endangered or threatened species are identified using the Department's Landscape Project as approved by the Department's Endangered and Nongame Species Program." (NJDEP, June 20, 2016)

Figure 11 combines the following:

- Floodplains
- Wetlands (based on 2012 Land Use data; Note: an LOI is necessary to determine actual boundary of wetlands)
- 150 foot wetlands buffers (based on 2012 Land Use data; an LOI is necessary to determine actual buffer for wetlands – wetlands are given a 150', 50' or 0' buffer)
- Riparian buffers: 50 foot buffers are given for most streams)
- Landscape Project version 3.3: Rank 4 habitats
- Open space

References: Environmentally Critical Areas

NJDEP. June 20, 2016. N.J.A.C. 7:8 Stormwater Management Rule. Date last amended: June 20, 2016. 39 pages. http://www.nj.gov/dep/rules/rules/njac7_8.pdf

GIS data:


Floodplains: See Section 6B; Figure 6c

Wetlands & wetlands buffers: See Section 6C; Figure 6d

Riparian buffers: See Section 6D; Figure 6b


Landscape Project v3.3 Rank 3 or 4: See Section 7F; Figure 7d

Open Space: See Section 8C; Figure 8a



Legend

- | | | | |
|--|--------------------------------|-------------------------------------|---|
| | West Cape May Borough | | Open Space |
| | Roads | | Preserved Farmland |
| | Water | | Landscape v.3.3 Rank 4 - State Endangered |
| | 50' Riparian Buffer | | Wetlands |
| | Potential vernal pool location | | 150' buffer may apply to some wetlands |
| | Vernal pool location | Flood Zone (FEMA, 10/4/2017) | |
| | Vernal habitat area | | AE (base elevations determined) |
| | Potential vernal habitat area | | 0.2 % annual chance flood hazard |



Data Sources: NJDEP, NJDOT
Note: Map accuracy is limited to the accuracy and scale of the original data sets.
Disclaimer: This map was developed using NJDEP GIS digital data, but this secondary product has not been verified by NJDEP and is not NJDEP authorized.

Figure 11: West Cape May Environmentally Critical Areas
"Environmentally critical area" means an area or feature which is of significant environmental value. This map combines some of the mapped layers from previous sections that make an area environmentally critical.

APPENDIX A: DATA USE AGREEMENTS

Contents

A.1. Terms of Agreement for use of NJDEP GIS data

A.2. Cautions and Restrictions on Use of Natural Heritage Data

A.1 Terms of Agreement for use of NJDEP GIS data

(Required by NJDEP Office of Information Management, Bureau of Geographic Information and Analysis.)

1. Digital data received from the NJDEP are to be used solely for internal purposes in the conduct of daily affairs.
2. The data are provided, as is, without warranty of any kind and the user is responsible for understanding the accuracy limitations of all digital data layers provided herein, as documented in the accompanying Data Dictionary and Readme files. Any reproduction or manipulation of the above data must ensure that the coordinate reference system remains intact.
3. Digital data received from the NJDEP may not be reproduced or redistributed for use by anyone without first obtaining written permission from the NJDEP. This clause is not intended to restrict distribution of printed mapped information produced from the digital data.
4. Any maps, publications, reports, or other documents produced as a result of this project that utilize NJDEP digital data will credit the NJDEP Geographic Information System (GIS) as the source of the data with the following credit/disclaimer:

This (map/publication/report) was developed using New Jersey Department of Environmental Protection Geographic Information System digital data, but this secondary product has not been verified by NJDEP and is not state-authorized.

5. Users shall require any independent contractor, hired to undertake work that will utilize digital data obtained from the NJDEP, to agree not to use, reproduce, or redistribute NJDEP GIS data for any purpose other than the specified contractual work. All copies of NJDEP GIS data utilized by an independent contractor will be required to be returned to the original user at the close of such contractual work. Users hereby agree to abide by the use and reproduction conditions specified above and agree to hold any independent contractor to the same terms. By using data provided herein, the user acknowledges that terms and conditions have been read and that the user is bound by these criteria.

A.2 Cautions and Restrictions on Use of Natural Heritage Data

(Required by NJDEP Division of Parks and Forestry, Natural Lands Management.)

CAUTIONS AND RESTRICTIONS ON NATURAL HERITAGE DATA

The quantity and quality of data collected by the Natural Heritage Program is dependent on the research and observations of many individuals and organizations. Not all of this information is the result of comprehensive or site-specific field surveys. Some natural areas in New Jersey have never been thoroughly surveyed. As a result, new locations for plant and animal species are continuously added to the database. Since data acquisition is a dynamic, ongoing process, the Natural Heritage Program cannot provide a definitive statement on the presence, absence, or condition of biological elements in any part of New Jersey. Information supplied by the Natural Heritage Program summarizes existing data known to the program at the time of the request regarding the biological elements or locations in question. They should never be regarded as final statements on the elements or areas being considered, nor should they be substituted for on-site surveys required for environmental assessments. The attached data is provided as one source of information to assist others in the preservation of natural diversity.

This office cannot provide a letter of interpretation or a statement addressing the classification of wetlands as defined by the Freshwater Wetlands Act. Requests for such determination should be sent to the DEP Division of Land Use Regulation, P.O. Box 439, Trenton, NJ 08625-0439.

The Landscape Project was developed by the Division of Fish & Wildlife, Endangered and Nongame Species Program in order to map critical habitat for rare animal species. Natural Heritage Database response letters will also list all species (if any) found during a search of the Landscape Project. However, this office cannot answer any inquiries about the Landscape Project. All questions should be directed to the DEP Division of Fish and Wildlife, Endangered and Nongame Species Program, P.O. Box 400, Trenton, NJ 08625-0400.

This cautions and restrictions notice must be included whenever information provided by the Natural Heritage Database is published.



NJ Department of Environmental Protection
Division of Parks and Forestry

Natural Lands Management

APPENDIX B: METADATA FOR GIS DATA LAYERS USED FOR THIS REPORT

Data Disclaimers in **Appendix A** apply to the use of these data layers and the maps created from them. The user is responsible for understanding the accuracy limitations of the digital data layers, as documented in the accompanying report and metadata summaries, and the metadata files which accompany the data.

APPENDIX B: GIS DATA LAYERS USED FOR THIS REPORT

Figure	Source of Data*	Data Title	Date	Scale	Online Linkage
Base Layers on many or most maps	NJDEP BGIS	Municipalities of New Jersey (Clipped to Coast), Edition 20121228 (Govt_admin_mun_coast_bnd)	12/28/2012	1:2,400	http://www.state.nj.us/dep/gis/stateshp.html#MUNCOAST
	NJDEP BGIS	NJDEP State Boundary of New Jersey	11/1/1998	1:24,000	http://www.state.nj.us/dep/gis/stateshp.html#STATE
	NJDEP BGIS	NJDEP County Boundaries for the State of New Jersey	1/23/2003	1:100,000	http://njogis-newjersey.opendata.arcgis.com/datasets/5f45e1ece6e14ef5866974a7b57d3b95_1
	NJDEP BGIS	National Hydrography Dataset (NHD) Waterbody 2002	11/1/2010	1:2,400	http://www.state.nj.us/dep/gis/digidownload/zip/statewide/nhdwaterbody2002shp.zip
	NJDEP BGIS	National Hydrography Dataset (NHD) Streams 2002	11/1/2010	1:2,400	http://www.state.nj.us/dep/gis/digidownload/zip/statewide/nhdstreams2002shp.zip
	Cape May County Planning Department	CMC_Parcels (20110923)	9/23/2011	1:400	https://njgin.state.nj.us/NJ_NJGINExplorer/DataDownloads.jsp
Figure 1a: Location of Municipality	NJDEP	Alternative Fueled Vehicle Fueling Stations for New Jersey, Edition 20160706 (Strc_alt_fuel)	7/6/2016	na	http://njogis-newjersey.opendata.arcgis.com/datasets/4b93a5c694f7413a9def60139a49ae1f_9
	NJDEP	Solar PV Grid Supply Installations of New Jersey, Edition 20170927 (Util_solar_supply)	1/20/2017	na	http://njogis-newjersey.opendata.arcgis.com/datasets/new-jersey-solar-pv-grid-supply-installations
	NJDOT	New Jersey Lighthouses	2/22/2017	na	http://njogis-newjersey.opendata.arcgis.com/datasets/new-jersey-lighthouses
Figure 1b: 1930 Aerial Photography	NJOIT OGIS	1930s Aerial Photography of New Jersey Web Map Service (WMS)	10/1/2009	na	WMS Server: http://geodata.state.nj.us/imagerywms/BlackWhite1930?
Figure 1c: 1995 Aerial Photography	USGS	1994/1995 Digital Orthophoto Quadrangles (DOQ)	6/21/1995	1:12,000	https://njgin.state.nj.us/NJ_NJGINExplorer/IW.jsp
Figure 1d: 2002 Aerial Photography	NJOIT OGIS	New Jersey 2002 High Resolution Orthophotography	7/31/2003	1:2,400	https://njgin.state.nj.us/NJ_NJGINExplorer/IW.jsp
Figure 1e: 2007 Aerial Photography	NJOIT OGIS	New Jersey 2007 - 2008 High Resolution Orthophotography, MrSID 5K Tiles	10/1/2008	1:19,200	https://njgin.state.nj.us/NJ_NJGINExplorer/IW.jsp
Figure 1f: 2012 Aerial Photography	NJOIT OGIS	New Jersey 2012 - 2013 High Resolution Orthophotography, NAD83 NJ State Plane Feet, MrSID Tiles	3/1/2013	1:2,400	https://njgin.state.nj.us/NJ_NJGINExplorer/IW.jsp
Figure 1g: 2015 Aerial Photography	NJOIT OGIS	New Jersey 2015 High Resolution Orthophotography, NAD83(2011) NJ State Plane Feet, MrSID Tiles	2/24/2016	1:2,400	https://njgin.state.nj.us/NJ_NJGINExplorer/IW.jsp
Figure 1h: Land Use Type (2012)	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice (Land_lu_2012_hu02040204_206)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
Figure 1h: Land Use Type (2012)	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague (Land_lu_2012_hu02040302_303)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
Figure 1i: Change in Land Use Type	NJDEP BGIS	NJDEP 1986 Land Use/Land Cover for Cape May County, New Jersey	11/1/1998	1:24,000	http://www.nj.gov/dep/gis/lulcshp.html#CAP
	NJDEP BGIS	NJDEP 1995/97 Land use/Land cover Update, Cape May Watershed Management Area, WMA-16	12/1/2000	1:12,000	http://www.nj.gov/dep/gis/lulc95shp.html

Figure	Source of Data*	Data Title	Date	Scale	Online Linkage
	NJDEP BGIS	NJDEP 2002 Land use/Land cover Update, Cape May Watershed Management Area, WMA-16, Edition 20080304	3/4/2008	1:2,400	http://www.nj.gov/dep/gis/lulc02cshp.html
	NJDEP BGIS	NJDEP 2007 Land use/Land Cover Update, Cape May Watershed Management Area, WMA16	7/12/2010	1:2,400	http://www.nj.gov/dep/gis/lulc07cshp.html
Figure 2a: Historical Coastline	NJDEP BGIS	NJDEP Coastline of New Jersey, 1836-1977 (created 1991)	1/1/1991	unknown	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/histshore.zip
	NJDEP BGIS	NJDEP Coastline of New Jersey (published in 1996 from 1986 aerials)	1/1/1996	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/coast.zip
	NJDEP BGIS	Coastline of New Jersey, Edition 20150501 (Land_coastline_2007)	5/1/2015	1:2,400	http://www.state.nj.us/dep/gis/stateshp.html#NJCOAST07
	NJDEP BGIS	Coastline of New Jersey, Edition 20150501 (Land_coastline_2012)	5/1/2015	1:2,400	http://www.state.nj.us/dep/gis/stateshp.html#NJCOAST12
	ESRI	Imagery Basemap from ESRI		1:70,000	http://goto.arcgisonline.com/maps/World_Imagery
Figure 2b: Drought Regions & Air Quality	NADP	National Atmospheric Deposition Network	6/27/1905	na	http://nadp.sws.uiuc.edu/data/
	NJDEP NJGS	DGS00-1: Drought Regions of New Jersey	5/1/2004	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs00-1.htm
	NJDEP BAM	NJDEP Ambient Air Quality Monitors	3/1/2006	1:1,600	http://njogis-newjersey.opendata.arcgis.com/datasets/new-jersey-air-quality-monitoring-stations
Figure 2c: Public Water	NJDEP DSRT BEA	NJDEP Public Community Water Purveyor Service Areas, 1998	7/12/2004	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/watpurv1998.zip
	NJGS	DGS97-1 NJDEP Public-Community Water-Supply Wells of New Jersey	7/7/2011	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs97-1.htm
Figure 2d: Public Sewer	NJDEP	Statewide Sewer Service Area for New Jersey, Edition 20170124 (Util_wastewater_servicearea)	1/24/2017	na	http://njogis-newjersey.opendata.arcgis.com/datasets/2ceba1ef852b4940afc3e0d94fb5d327_6
	NJDEP	NJPDES Surface Water Discharges in New Jersey, (1:12,000) Edition 20170213 (Strc_NJPDES_sw_pipe)	2/13/2017	1:12,000	http://njogis-newjersey.opendata.arcgis.com/datasets/2ceba1ef852b4940afc3e0d94fb5d327_6
Figure 3a: Physiographic Provinces	NJDEP BGIA	NJDEP Digital Elevation Grid for New Jersey (100 meter)	5/1/2002	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/nj100mlat.zip
	NJDEP BGIA	NJDEP Hillshade Grid for New Jersey (100 meters)	5/1/2002	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/nj100mhill.zip
	NJGS	DGS02-7: Physiographic Provinces of New Jersey	6/30/2002	1:100,000	http://www.state.nj.us/dep/njgs/geodata/dgs02-7.htm
	NJGS	DGS04-1: Earthquakes Epicentered In New Jersey, Edition 20161107	11/7/2016	n/a	http://www.state.nj.us/dep/njgs/geodata/dgs04-1.htm
Figure 3b: Bedrock Geology	NJGS	DGS04-6: Bedrock Geology for New Jersey 1:100,000 Scale	6/30/1999	1:100,000	http://www.state.nj.us/dep/njgs/geodata/dgs04-6.htm
Figure 3c: Surficial Geology	NJGS	DGS07-2: Surficial Geology of New Jersey	9/11/2013	1:100,000	http://www.state.nj.us/dep/njgs/geodata/dgs07-2.htm
	NJGS	Thickness of Surficial Deposits in New Jersey	9/11/2013	1:100,000	http://njgeology.org/geodata/dgs07-2.htm
	NJDEP NJG&WS	DGS04-7: Historic Fill For New Jersey as of January 2016	1/26/2016	1:100,000	http://www.state.nj.us/dep/njgs/geodata/dgs04-7.htm
Figure 3d: Shaded Elevation (Hillshade)	NJDEP BGIA	NJDEP Digital Elevation Grid for New Jersey (100 meter)	5/1/2002	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/nj100mlat.zip
	NJDEP BGIA	NJDEP Hillshade Grid for New Jersey (100 meters)	5/1/2002	1:24,000	http://www.state.nj.us/dep/gis/digdownload/zips/statewide/nj100mhill.zip
Figure 4a-4g: Soil Maps	USDA	Soil Survey Geographic (SSURGO) database for Cape May County, New Jersey	9/28/2016	1:24,000	http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx
Figure 5b: Aquifer, GW Monitoring wells	NJGS	DGS98-5 Aquifers of New Jersey	5/21/1998	1:100,000	http://www.state.nj.us/dep/njgs/geodata/dgs98-5.htm
	NJGS	DGS98-6 NJDEP Sole-Source Aquifers in New Jersey	4/5/2000	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs98-6.htm
	NJDEP	New Jersey Pollution Discharge Elimination System (NJPDES) Regulated Facility Locations, Edition 20070718 (Envr_NJEMS_NJPDES_all)	7/18/2007	unknown	http://njogis-newjersey.opendata.arcgis.com/datasets/njdep-non-point-wastewater-sites
	NJGS	DGS97-1 NJDEP Public-Community Water-Supply Wells of New Jersey	7/7/2011	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs97-1.htm ; new message (noticed it on 10/8/17) sensitive nature so no longer available
	NJGS	DGS04-5 Well Head Protection Areas For Public Non-Community Water Supply Wells In New Jersey.	2/19/2015	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs04-5.htm

Figure	Source of Data*	Data Title	Date	Scale	Online Linkage
	USGS	USGS ground water level monitoring	na	na	
Figure 5d: Ground Water Recharge & Aquifer Rank	NJGS	DGS02-3 Ground-Water Recharge for Cape May County, NJ	10/8/2004	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs02-3.htm
	NJGS	DGS07-1: Aquifer-Recharge Potential for Cape May County, NJ	1/4/2005	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs07-1.htm
Figure 6a: Watershed	NJDEP OEA CTD	NJDEP Head of Tide Points for Watercourses of New Jersey	1/1/1986	1:24,000	http://www.state.nj.us/dep/gis/digidownload/zips/statewide/hot.zip
	NJGS	DGS06-2 New Jersey Tidal Benchmark Network	3/1/2006	na	http://www.state.nj.us/dep/njgs/geodata/dgs06-2.htm
	NJGS	Canals and Water Raceways of New Jersey	10/1/2008	1:24,000	http://www.state.nj.us/dep/njgs/geodata/dgs08-1.htm#image
	NJDEP DWM	NJDEP Watershed Management Areas in New Jersey (Version 200901)	1/1/2009	1:24,000	http://www.nj.gov/dep/gis/stateshp.html#VMAS
	NJGS	14 Digit Hydrologic Unit Code Delineations for New Jersey (Version 20110225)	2/25/2011	1:24,000	http://www.nj.gov/dep/gis/stateshp.html#HUC14
Figure 6b: Subwatershed & SWQS	NJDEP WMS	NJDEP Surface Water Quality Standards of New Jersey (Version 201012)	12/1/2010	1:2,400	http://www.state.nj.us/dep/gis/stateshp.html#SWQS
Figure 6c: Floodplains	FEMA	National Flood Hazard Layer (NFHL) (Cape May, NJ)	10/4/2017	1:12,000	https://msc.fema.gov
Figure 6d: Wetlands	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice (Land_lu_2012_hu02040204_206)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague (Land_lu_2012_hu02040302_303)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
Figure 6e: Integrated List	NJDEP WMS	Integrated List of Waters for New Jersey, 2014 (Integrated List), Edition 20170927 (Envr_mon_water_IR_2014_use)	9/27/2017	1:24,000	http://njgis-newjersey.opendata.arcgis.com/datasets/2014-new-jersey-integrated-list-of-waters-integrated-list?geometry=-82.144,38.633,-61.237,41.573
Figure 6f: Percent Impervious Surface & Discharge	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice (Land_lu_2012_hu02040204_206)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague (Land_lu_2012_hu02040302_303)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP	NJPDES Surface Water Discharges in New Jersey, (1:12,000) Edition 20170614 (Strc_NJPDES_sw_pipe)	6/14/2017	1:12,000	http://njgis-newjersey.opendata.arcgis.com/datasets/55e8b1a25706431490a8f3becf861f91_5
Figure 6g: Monitoring	NJDEP BFBM	STORET Water Quality Monitoring Stations	8/1/2005	unknown	http://www.state.nj.us/dep/gis/digidownload/zips/statewide/storet.zip
Figure 7a-7c: Land Cover (2012)	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice (Land_lu_2012_hu02040204_206)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague (Land_lu_2012_hu02040302_303)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
Figure 7d: Wildfire Fuel Hazard	NJDEP NJFFS	2002 NJFFS Wildfire Fuel Hazard for Cape May County, New Jersey	4/17/2009	1:2,400	http://www.state.nj.us/dep/gis/njfn.html#CAP
Figure 7e: Important Bird Areas (Audubon)	State Audubon	Important Bird Areas	3/2/2016	1:50,000	ArcGIS Map Service or http://gis.audubon.org/arcgisweb/rest/services/NAS/ImportantBirdAreas_Poly/MapServer
Figure 7f: Landscape Project version 3.3	NJDEP DFW ENSP	NJDEP Species Based Habitat, Vernal Habitat, Version 3.3, 20170509 (Envr_hab_ls_v3_3_vernalhabitat)	5/9/2017	1:12,000	http://www.state.nj.us/dep/gis/landscape.html#geodatabase
	NJDEP DFW ENSP	NJDEP Species Based Habitat, Vernal Pools, Version 3.3, 20170509	5/9/2017	1:12,000	http://www.state.nj.us/dep/gis/landscape.html#geodatabase

Figure	Source of Data*	Data Title	Date	Scale	Online Linkage
		(Envr_hab_ls_v3_3_vernalpools)			
	NJDEP DFW ENSP	NJDEP Species Based Habitat, Delaware Bay Region, Version 3.3, 20170509 (Envr_hab_ls_v3_3_delbay)	5/9/2017	1:12,000	http://www.state.nj.us/dep/gis/landscape.html#geodatabase
	NJDEP DFW ENSP	NJDEP Species Based Habitat, Atlantic Coastal Region, Version 3.3, 20170509 (Envr_hab_ls_v3_3_coastal)	5/9/2017	1:12,000	http://www.state.nj.us/dep/gis/landscape.html#geodatabase
Figure 7g: Natural Heritage Priority Sites and Grid	NJDEP ONLM	NJDEP Natural Heritage Priority Sites	3/1/2007	1:24,000	http://www.state.nj.us/dep/gis/stateshp.html#PRIORITY
	NJDEP ONLM	NJDEP Natural Heritage Grid Map, Version 200911	3/21/2013	1:24,000	http://www.state.nj.us/dep/gis/stateshp.html#NHPGRID
Figure 8a: Preserved Open Space & Recreation	NJDA SADC	NJFPP - New Jersey Farmland Preservation Program	1/8/2013	1:24,000	http://www.nj.gov/agriculture/sadc/farmpreserve/resources/njfp.zip
	NJDEP	State Owned, Protected Open Space and Recreation Areas in New Jersey, Edition 20160527 (Land_owner_openspace_state)	5/27/2016	1:2,400	http://njgis-newjersey.opendata.arcgis.com/datasets/b75fe11ed90543c1b4ee87e66af63b8b_1
	KES	Open Space - Other (selected parcels from CMC_parcel)	3/16/2017	na	na
Figure 9a: Archeological & Historical	NJDEP NHR HPO	Historic Property Features of New Jersey, Edition 20160922 (Land_use_HPO_property_feature)	9/22/2016	NA	https://geodata.state.nj.us/arcgis/rest/services/Features/Land_Use_and_Land_Cover/MapServer/7
	NJDEP NHR HPO	Archaeological Site Grid of New Jersey, Edition 20160922 (Land_use_HPO_arch_grid)	9/22/2016	NA	https://geodata.state.nj.us/arcgis/rest/services/Features/Land_Use_and_Land_Cover/MapServer/8
Figure 9b: Historical Properties	NJDEP NHR HPO	Historic Districts of New Jersey, Edition 20160922 (Land_use_HPO_district)	9/22/2016	NA	https://geodata.state.nj.us/arcgis/rest/services/Features/Land_Use_and_Land_Cover/MapServer/9
	NJDEP NHR HPO	Historic Property Features of New Jersey, Edition 20111121 (Land_use_HPO_property_feature)	11/21/2017	NA	https://geodata.state.nj.us/arcgis/rest/services/Features/Land_Use_and_Land_Cover/MapServer/6
Figure 11: Composite Map of Environmentally Critical Areas	NJDEP BGIS	National Hydrography Dataset (NHD) Streams 2002	11/1/2010	1:2,400	http://www.state.nj.us/dep/gis/digidownload/zips/statewide/nhdstreams2002shp.zip
	NJDA SADC	NJFPP - New Jersey Farmland Preservation Program	1/8/2013	1:24,000	http://www.nj.gov/agriculture/sadc/farmpreserve/resources/njfp.zip
	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040204 - Delaware Bay, Subbasin 02040206 - Cohansey-Maurice (Land_lu_2012_hu02040204_206)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP BGIS	Land Use/Land Cover 2012 Update, Edition 20150217 Subbasin 02040302 - Great Egg Harbor, Subbasin 02040303 - Chincoteague (Land_lu_2012_hu02040302_303)	2/17/2015	1:2,400	http://www.nj.gov/dep/gis/listall.html
	NJDEP	State Owned, Protected Open Space and Recreation Areas in New Jersey, Edition 20160527 (Land_owner_openspace_state)	5/27/2016	1:2,400	http://njgis-newjersey.opendata.arcgis.com/datasets/b75fe11ed90543c1b4ee87e66af63b8b_1
	KES	Open Space - Other (selected parcels from CMC_parcel)	3/16/2017	1:400	na
	FEMA	National Flood Hazard Layer (NFHL) (Cape May, NJ)	10/4/2017	1:12,000	https://msc.fema.gov

***Sources of GIS Data:**

Data Sources	Full Name of Data Source
Cape May County PD	Cape May County Planning Department
DRBC	Delaware River Basin Commission
ESRI	ESRI Inc.
FEMA	Federal Emergency Management Agency
KES	Kratzer Environmental Services
NADP	National Atmospheric Deposition Program
NJDA SADC	New Jersey Department of Agriculture (NJDA), State Agriculture Development Committee (SADC)
NJDEP	NJ Department of Environmental Protection
NJDEP BAM	NJDEP Dept of Environmental Regulation, Bureau of Air Monitoring
NJDEP BFBM	NJDEP Bureau of Freshwater Biological Monitoring
NJDEP BGIA	NJDEP Bureau of Geographic Information and Analysis
NJDEP BGIS	NJDEP Bureau of Geographic Information Services
NJDEP DFW ENSP	NJDEP Division of Fish & Wildlife, Endangered Nongame Species Program
NJDEP DSRT BEA	NJDEP Division of Science, Research, and Technology, Bureau of Environmental Assessment
NJDEP DWM	NJDEP Division of Watershed Management
NJDEP GA	NJDEP Green Acres
NJDEP NHR HPO	NJDEP Natural and Historic Resources, Historic Preservation Office
NJDEP NJFFS	NJDEP New Jersey Forest Fire Service
NJDEP NJG&WS	NJDEP New Jersey Geological and Water Survey
NJDEP OEA CTD	NJDEP Office of Environmental Analysis (OEA), Coast survey Limited (CTD)
NJDEP ONLM	NJDEP Office of Natural Lands Management
NJDEP WMS	NJDEP Division of Water Monitoring and Standards
NJDOT	New Jersey Department of Transportation
NJGIN	New Jersey Geographic Information Network
NJGS	New Jersey Geological Service
NJOIT OGIS	New Jersey Office of Information Technology, Office of GIS
PA DCNR	Pennsylvania Department of Conservation and Natural Resources
State Audubon	NJ State Audubon
USDA/NRCS	United States Department of Agriculture, Natural Resources Conservation Service
USGS	United States Geological Survey

APPENDIX C: REGIONAL FAUNA

Contents:

C.1 List of Birds

C.2 List of Mammals

C.3 List of Reptiles

C.4 List of Amphibians

C.5 List of Fish

C.1 Cape May County Birds

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
<u>Ducks, Geese & Swans</u>					
RV	Greater White-fronted Goose	<i>Anser albifrons</i>	7	3/3/2014	✓
RV	Pink-footed goose	<i>Anser brachyrhynchus</i>	1	2/26/2017	
	Snow Goose	<i>Chen caerulescens</i>	15000	12/8/2008	✓
RV	Ross's Goose	<i>Chen rossii</i>	2	12/18/2014	✓
	Brant	<i>Branta bernicla</i>	3200	1/2/2014	✓
RV	Cackling Goose	<i>Branta hutchinsii</i>	7	12/6/2014	✓
	Canada Goose	<i>Branta canadensis</i>	20325	12/9/2010	✓
RV	Barnacle goose	<i>Branta leucopsis</i>	1	11/24/2013	
int	Mute Swan	<i>Cygnus olor</i>	135	1/10/2003	✓
RV	Trumpeter Swan	<i>Cygnus buccinator</i>	1	1/5/2006	
	Tundra Swan	<i>Cygnus columbianus</i>	102	11/18/2014	✓
	Fulvous whistling duck	<i>Dendrocygna bicolor</i>	2	10/5/1974	
RV	Black-bellied whistling duck	<i>Dendrocygna autumnalis</i>	12	7/14/2016	✓
	Wood Duck	<i>Aix sponsa</i>	428	10/21/2013	✓
	Gadwall	<i>Anas strepera</i>	450	11/7/2014	✓
	Eurasian Wigeon	<i>Anas penelope</i>	5	10/31/2014	✓
	American Wigeon	<i>Anas americana</i>	425	10/31/1998	✓
	American Black Duck	<i>Anas rubripes</i>	1100	2/10/1929	✓
	Mallard	<i>Anas platyrhynchos</i>	880	2/6/2014	✓
	Blue-winged Teal	<i>Anas discors</i>	120	9/15/2008	✓
	Northern Shoveler	<i>Anas clypeata</i>	200	11/10/2013	✓
	Northern Pintail	<i>Anas acuta</i>	1000	2/12/2012	✓
	Green-winged Teal	<i>Anas crecca</i>	2510	3/7/2008	✓
	Canvasback	<i>Aythya valisineria</i>	12	1/11/2015	
	Redhead	<i>Aythya americana</i>	60	12/14/2013	✓
	Ring-necked Duck	<i>Aythya collaris</i>	600	2/19/2017	✓
	Greater Scaup	<i>Aythya marila</i>	1075	1/1/2014	✓
	Lesser Scaup	<i>Aythya affinis</i>	300	11/2/2008	✓
	King Eider	<i>Somateria spectabilis</i>	3	12/2/2014	
	Common Eider	<i>Somateria mollissima</i>	227	11/27/2009	
	Harlequin Duck	<i>Histrionicus histrionicus</i>	8	3/14/2004	
	Surf Scoter	<i>Melanitta perspicillata</i>	66587	10/26/2007	✓
	White-winged Scoter	<i>Melanitta fusca</i>	424	11/29/2010	
	Black Scoter	<i>Melanitta americana</i>	74998	10/26/2007	✓
	Long-tailed Duck	<i>Clangula hyemalis</i>	1000	3/5/2000	✓
	Bufflehead	<i>Bucephala albeola</i>	1063	11/17/2007	✓
	Common Goldeneye	<i>Bucephala clangula</i>	51	1/7/2012	
RV	Barrow's Goldeneye	<i>Bucephala islandica</i>	1	11/28/2007	
	Hooded Merganser	<i>Lophodytes cucullatus</i>	405	12/17/2006	✓
	Common Merganser	<i>Mergus merganser</i>	59	3/7/2008	
	Red-breasted Merganser	<i>Mergus serrator</i>	523	11/19/2007	✓
	Ruddy Duck	<i>Oxyura jamaicensis</i>	2254	1/28/2016	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
<u>Grouse, Quail & Allies</u>					
	Northern Bobwhite	<i>Colinus virginianus</i>	30	10/21/2008	✓
	Ring-necked Pheasant	<i>Phasianus colchicus</i>	1	4/1/2016	✓
	Ruffed Grouse	<i>Bonasa umbellus</i>	1	11/3/2014	✓
	Wild Turkey	<i>Meleagris gallopavo</i>	61	1/12/2012	✓
<u>Grebes</u>					
Ebr, SCnb	Pied-billed Grebe	<i>Podilymbus podiceps</i>	20	11/23/1998	✓
	Horned Grebe	<i>Podiceps auritus</i>	70	1/11/2015	
	Red-necked Grebe	<i>Podiceps grisegena</i>	13	3/2/2015	
RV	Eared Grebe	<i>Podiceps nigricollis</i>	1	9/30/2012	
RV	Western Grebe	<i>Aechmophorus occidentalis</i>	1	12/1/2013	
<u>Pigeons & Doves</u>					
int	Rock Pigeon	<i>Columba livia</i>	389	12/18/2016	✓
RV	Band-tailed Pigeon	<i>Patagioenas leucocephala</i>	1	1/18/2007	✓
RV	Eurasian Collared-Dove	<i>Streptopelia decaocto</i>	5	9/6/2012	✓
RV	White-winged Dove	<i>Zenaida asiatica</i>	1	6/11/2016	✓
	Mourning Dove	<i>Zenaida macroura</i>	302	12/16/2012	✓
RV	Common Ground-Dove	<i>Columbina passerina</i>	1	10/29/2010	
<u>Cuckoos</u>					
	Yellow-billed Cuckoo	<i>Coccyzus americanus</i>	10	9/1/1983	✓
SCbr	Black-billed Cuckoo	<i>Coccyzus erythrophthalmus</i>	3	8/13/2011	✓
<u>Nightjars</u>					
SC	Common Nighthawk	<i>Chordeiles minor</i>	1202	9/6/2012	✓
RV	Lesser Nighthawk	<i>Chordeiles acutipennis</i>	1	11/25/2007	
	Chuck-will's-widow	<i>Antrostomus carolinensis</i>	12	5/21/2015	✓
SCbr, Unb	Eastern Whip-poor-will	<i>Antrostomus vociferous</i>	18	5/11/2012	✓
<u>Swifts & Hummingbirds</u>					
	Chimney Swift	<i>Chaetura pelagica</i>	517	10/7/2012	✓
RV	Black Swift	<i>Cypseloides niger</i>	1	8/28/2011	
	Ruby-throated Hummingbird	<i>Archilochus colubris</i>	35	8/17/2014	✓
RV	Black-chinned Hummingbird	<i>Archilochus alexandri</i>	1	11/17/2016	
RV	Broad-tailed Hummingbird	<i>Selasphorus platycercus</i>	1	4/3/2012	
RV	Rufous Hummingbird	<i>Selasphorus rufus</i>	2	12/26/2008	✓
RV	Allen's Hummingbird	<i>Selasphorus sasin</i>	1	12/7/2004	
RV	Calliope Hummingbird	<i>Selasphorus calliope</i>	1	11/20/2015	
<u>Cranes & Rails</u>					
	Corn Crane	<i>Crex crex</i>	1	11/11/1905	
	Yellow rail	<i>Coturnicops</i>	1	10/26/2015	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
		<i>noveboracensis</i>			
Ebr, Tnb	Black rail	<i>Laterallus jamaicensis</i>	3	5/5/2008	✓
	Clapper Rail	<i>Rallus longirostris</i>	113	6/20/2009	✓
	King Rail	<i>Rallus elegans</i>	3	5/8/2013	✓
	Virginia Rail	<i>Rallus limicola</i>	17	12/15/2013	✓
	Sora	<i>Porzana carolina</i>	11	9/29/1972	✓
	Purple gallinule	<i>Poryphyryla martinica</i>	1	10/26/2016	✓
	Common Gallinule	<i>Gallinula galeata</i>	10	9/22/1973	✓
	American Coot	<i>Fulica americana</i>	266	11/25/2011	✓
	Sandhill Crane	<i>Grus canadensis</i>	14	11/26/2016	✓
<u>Plovers, Sandpipers & Allies</u>					
RV	Northern Lapwing	<i>Vanellus vanellus</i>	1	1/3/1997	
	Black-bellied Plover	<i>Pluvialis squatarola</i>	1000	10/4/1986	✓
	American Golden-Plover	<i>Pluvialis dominica</i>	21	10/23/2007	✓
RV	Wilson's Plover	<i>Charadrius wilsonia</i>	1	6/6/2016	
	Semipalmated Plover	<i>Charadrius semipalmatus</i>	1795	8/13/2004	✓
E	Piping Plover	<i>Charadrius melodus</i>	30	9/6/2011	✓
	Killdeer	<i>Charadrius vociferus</i>	415	10/23/2014	✓
RV	Lesser Sand-Plover	<i>Charadrius mongolus</i>	1	7/13/1990	
SC	American Oystercatcher	<i>Haematopus palliatus</i>	500	9/26/2006	✓
	Black-necked Stilt	<i>Himantopus mexicanus</i>	6	5/13/1988	
	American Avocet	<i>Recurvirostra americana</i>	4	7/15/2016	
SCbr	Spotted Sandpiper	<i>Actitis macularius</i>	25	7/27/2013	✓
	Solitary Sandpiper	<i>Tringa solitaria</i>	31	5/2/2016	✓
	Greater Yellowlegs	<i>Tringa melanoleuca</i>	660	11/1/2011	✓
	Willet	<i>Tringa semipalmata</i>	472	4/22/2014	✓
	Lesser Yellowlegs	<i>Tringa flavipes</i>	854	9/12/2009	✓
E	Upland Sandpiper	<i>Bartramia longicauda</i>	4	8/23/1987	✓
SCnb	Whimbrel	<i>Numenius phaeopus</i>	1144	5/5/2012	✓
RV	Long-billed Curlew	<i>Numenius americanus</i>	1	7/18/2011	
	Hudsonian Godwit	<i>Limosa haemastica</i>	29	10/10/2015	
	Marbled Godwit	<i>Limosa fedoa</i>	20	10/8/2006	✓
	Ruddy Turnstone	<i>Arenaria interpres</i>	3300	5/27/2014	✓
Enb	Red Knot	<i>Calidris canutus</i>	95530	5/21/1982	✓
SCnb	Sanderling	<i>Calidris alba</i>	4629	8/27/2004	✓
SCnb	Semipalmated Sandpiper	<i>Calidris pusilla</i>	5000	5/29/2014	✓
	Western Sandpiper	<i>Calidris mauri</i>	800	8/5/2007	✓
	Least Sandpiper	<i>Calidris minutilla</i>	1200	7/26/2013	✓
	White-rumped Sandpiper	<i>Calidris fuscicollis</i>	295	10/4/2010	✓
	Baird's Sandpiper	<i>Calidris bairdii</i>	3	9/8/2008	
	Pectoral Sandpiper	<i>Calidris melanotos</i>	101	9/5/2012	✓
	Purple Sandpiper	<i>Calidris maritima</i>	84	12/14/2008	
	Dunlin	<i>Calidris alpina</i>	11000	2/26/2012	✓
	Stilt Sandpiper	<i>Calidris himantopus</i>	120	8/13/2014	✓
	Curlew sandpiper	<i>Calidris ferruginea</i>	2	5/16/2005	
RV	Little Stint	<i>Calidris minuta</i>	1	7/10/2003	

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
RV	Red-necked Stint	<i>Calidris ruficollis</i>	1	8/30/2008	
RV	Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	1	10/8/2002	
	Ruff	<i>Calidris pugnax</i>	1	5/7/2016	
	Buff-breasted Sandpiper	<i>Tryngites subruficollis</i>	22	9/8/2007	✓
	Short-billed Dowitcher	<i>Limnodromus griseus</i>	2300	5/7/2009	✓
	Long-billed Dowitcher	<i>Limnodromus scolopaceus</i>	30	10/16/2011	✓
	Wilson's Snipe	<i>Gallinago delicata</i>	75	10/4/2010	✓
	American Woodcock	<i>Scolopax minor</i>	140	1/13/2011	✓
	Wilson's Phalarope	<i>Phalaropus tricolor</i>	3	9/19/1993	
	Red-necked Phalarope	<i>Phalaropus lobatus</i>	262	9/2/2006	
	Red Phalarope	<i>Phalaropus fulicarius</i>	235	11/8/2016	
	Black-legged Kittiwake	<i>Rissa tridactyla</i>	500	12/4/1982	
RV	Ivory Gull	<i>Pagophila eburnea</i>	1	12/9/2009	
RV	Sabine's Gull	<i>Xema sabini</i>	1	10/11/2009	
	Bonaparte's Gull	<i>Chroicocephalus philadelphia</i>	1900	12/20/2015	✓
	Black-headed Gull	<i>Chroicocephalus ridibundus</i>	5	3/11/2014	
	Little Gull	<i>Hydrocoloeus minutus</i>	3	2/10/2002	✓
	Laughing Gull	<i>Leucophaeus atricilla</i>	20000	5/13/2016	✓
RV	Franklin's Gull	<i>Leucophaeus pipixcan</i>	114	11/13/2015	✓
	Ring-billed Gull	<i>Larus delawarensis</i>	8000	11/24/2013	✓
RV	California Gull	<i>Larus californicus</i>	1	11/11/2011	
	Herring Gull	<i>Larus argentatus</i>	8000	12/14/2008	✓
RV	Thayer's Gull	<i>Larus thayeri</i>	1	2/19/2000	✓
	Iceland Gull	<i>Larus glaucooides</i>	2	2/4/2017	✓
	Lesser Black-backed Gull	<i>Larus fuscus</i>	32	7/20/2015	✓
	Glaucous Gull	<i>Larus hyperboreus</i>	2	12/21/2010	
	Great Black-backed Gull	<i>Larus marinus</i>	1200	12/14/2008	✓
RV	Black-tailed Gull	<i>Larus crassirostris</i>	1	12/10/1998	
	Bridled Tern	<i>Onychoprion anaethetus</i>	23	9/3/2000	
RV	Sooty Tern	<i>Onychoprion fuscatus</i>	25	9/16/1999	
E	Least Tern	<i>Sternula antillarum</i>	380	7/28/2016	✓
SC	Gull-billed Tern	<i>Gelochelidon nilotica</i>	15	7/5/2016	✓
SCbr	Caspian Tern	<i>Hydroprogne caspia</i>	250	9/13/2016	✓
	Black Tern	<i>Chlidonias niger</i>	125	9/3/2016	✓
RV	White-winged Tern	<i>Chlidonias leucopterus</i>	1	6/10/1989	
RV	Whiskered Tern	<i>Chlidonias hybrida</i>	1	9/20/2014	
E	Roseate Tern	<i>Sterna dougallii</i>	16	9/19/2003	
SCbr	Common Tern	<i>Sterna hirundo</i>	3000	5/18/2007	✓
RV	Arctic Tern	<i>Sterna paradisaea</i>	7	5/7/1993	
	Forster's Tern	<i>Sterna forsteri</i>	3788	11/4/2014	✓
	Royal Tern	<i>Thalasseus maximus</i>	1027	10/17/2006	✓
	Sandwich Tern	<i>Thalasseus sandvicensis</i>	34	9/19/2003	✓
E	Black Skimmer	<i>Rynchops niger</i>	2002	7/11/2008	✓
RV	Great Skua	<i>Stercorarius skua</i>	1	2/6/2016	

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
RV	South Polar Skua	<i>Stercorarius maccormicki</i>	2	5/25/2015	
	Pomarine Jaeger	<i>Stercorarius pomarinus</i>	91	10/30/2012	
	Parasitic Jaeger	<i>Stercorarius parasiticus</i>	172	11/11/2013	✓
	Long-tailed Jaeger	<i>Stercorarius longicaudus</i>	3	9/2/2006	
	Dovekie	<i>Alle alle</i>	397	2/12/2013	
	Common Murre	<i>Uria aalge</i>	27	1/16/2011	
	Thick-billed murre	<i>Uria lomvia</i>	1	11/7/2008	
	Razorbill	<i>Alca torda</i>	131	1/11/2015	
RV	Black Guillemot	<i>Cepphus grylle</i>	1	12/22/2011	
	Atlantic Puffin	<i>Fratercula arctica</i>	17	2/6/2016	
<u>Loons</u>					
	Red-throated Loon	<i>Gavia stellata</i>	20000	11/14/1985	✓
	Common Loon	<i>Gavia immer</i>	376	10/24/2016	✓
RV	Pacific Loon	<i>Gavia arctica</i>	1	10/24/2015	
<u>Subs</u>					
	Northern Fulmar	<i>Fulmarus glacialis</i>	10	3/14/2004	
	Cory's Shearwater	<i>Calonectris diomedea</i>	89	8/22/2008	
	Great Shearwater	<i>Puffinus gravis</i>	182	6/28/2015	
	Sooty Shearwater	<i>Puffinus griseus</i>	224	5/25/2002	
	Manx Shearwater	<i>Puffinus puffinus</i>	30	12/20/2016	
	Audubon's Shearwater	<i>Puffinus lherminieri</i>	12	9/18/2016	
RV	Black-capped Petrel	<i>Pterodroma hasitata</i>	8	7/13/1996	
	Wilson's Storm-Petrel	<i>Oceanites oceanicus</i>	419	5/20/2011	
	Leach's Storm-Petrel	<i>Oceanodroma leucorhoa</i>	16	5/20/2011	
RV	Band-rumped Storm-Petrel	<i>Oceanodroma castro</i>	11	7/13/1996	
RV	White-faced Storm-Petrel	<i>Pelagodroma marina</i>	1	9/18/2016	
RV	Black-browed Albatross	<i>Thalassarche melanophris</i>	1	5/12/1985	
RV	Yellow-nosed Albatross	<i>Thalassarche chlororhynchos</i>	1	5/23/2000	
<u>Storks</u>					
RV	Wood Stork	<i>Mycteria americana</i>	2	9/17/1983	✓
<u>Frigatebirds, Boobies, Cormorants, Darters & Allies</u>					
RV	Magnificent Frigatebird	<i>Fregata magnificens</i>	1	9/7/2012	✓
RV	Masked Booby	<i>Sula dactylatra</i>	1	10/15/2015	
RV	Brown Booby	<i>Sula leucogaster</i>	1	9/27/2016	
	Northern Gannet	<i>Morus bassanus</i>	16946	11/11/2008	✓
	Neotropic Cormorant	<i>Phalacrocorax brasilianus</i>	1	10/14/2015	
	Double-crested Cormorant	<i>Phalacrocorax auritus</i>	35500	10/17/2009	✓
	Great Cormorant	<i>Phalacrocorax carbo</i>	30	10/22/2012	✓
RV	Anhinga	<i>Anhinga anhinga</i>	1	5/1/2011	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
<u>Tropicbirds</u>					
RV	White-tailed Tropicbird	<i>Phaethon lepturus</i>	3	8/28/2011	
<u>Pelicans, Herons, Ibises & Allies</u>					
	American White Pelican	<i>Pelecanus erythrorhynchos</i>	11	11/29/2009	✓
	Brown Pelican	<i>Pelecanus occidentalis</i>	94	9/29/2010	
Ebr, SCnb	American Bittern	<i>Botaurus lentiginosus</i>	80	10/5/2008	✓
SC	Least Bittern	<i>Ixobrychus exilis</i>	6	6/18/2015	✓
SCbr	Great Blue Heron	<i>Ardea herodias</i>	3000	10/2/1994	✓
	Great Egret	<i>Ardea alba</i>	375	8/24/2014	✓
SCbr	Snowy Egret	<i>Egretta thula</i>	891	8/27/2014	✓
SC	Little Blue Heron	<i>Egretta caerulea</i>	100	4/4/1970	✓
SC	Tricolored Heron	<i>Egretta tricolor</i>	92	8/19/2014	✓
Tbr, SCnb	Cattle Egret	<i>Bubulcus ibis</i>	50	5/11/1986	✓
	Green Heron	<i>Butorides virescens</i>	310	9/13/2015	✓
Tbr, SCnb	Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>	300	10/5/2008	✓
T	Yellow-crowned Night-Heron	<i>Nyctanassa violacea</i>	58	7/5/2015	✓
RV	White Ibis	<i>Eudocimus albus</i>	4	9/1/2015	✓
SCbr	Glossy Ibis	<i>Plegadis falcinellus</i>	400	4/28/2016	✓
RV	White-faced Ibis	<i>Plegadis chihi</i>	3	5/14/2005	✓
<u>New World Vultures</u>					
	Black Vulture	<i>Coragyps atratus</i>	100	11/6/2010	✓
	Turkey Vulture	<i>Cathartes aura</i>	657	11/19/2014	✓
<u>Hawks, Kites, Eagles & Allies</u>					
Tbr	Osprey	<i>Pandion haliaetus</i>	800	10/2/1986	✓
	Swallow-tailed Kite	<i>Elanoides forficatus</i>	3	5/3/2013	✓
	Mississippi Kite	<i>Ictinia mississippiensis</i>	17	6/13/2016	✓
Ebr, Tnb	Bald Eagle ²⁸	<i>Haliaeetus leucocephalus</i>	36	9/14/2013	✓
Ebr, SCnb	Northern Harrier	<i>Circus cyaneus</i>	266	10/2/1986	✓
SC	Sharp-shinned Hawk	<i>Accipiter striatus</i>	5025	10/2/1986	✓
SCbr	Cooper's Hawk	<i>Accipiter cooperii</i>	587	10/11/2009	✓
Ebr, SCnb	Northern Goshawk	<i>Accipiter gentilis</i>	11	10/23/1993	✓
Ebr, SCnb	Red-shouldered Hawk	<i>Buteo lineatus</i>	233	11/4/2012	✓
SCbr	Broad-winged Hawk	<i>Buteo platypterus</i>	2986	10/2/1986	✓
RV	Swainson's Hawk	<i>Buteo swainsoni</i>	2	10/31/1998	✓
RV	Zone-tailed Hawk	<i>Buteo albonotatus</i>	1	9/23/2015	✓
	Red-tailed Hawk	<i>Buteo jamaicensis</i>	428	11/4/2012	✓
	Rough-legged Hawk	<i>Buteo lagopus</i>	4	3/8/2015	✓
	Golden Eagle	<i>Aquila chrysaetos</i>	14	11/4/2012	✓

²⁸ There are no recent bald eagle nests in West Cape May (Michael O'Brien, personal communication, May 15, 2018).

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
<u>Owls</u>					
	Barn Owl	<i>Tyto alba</i>	5	10/10/2010	✓
	Eastern Screech-Owl	<i>Megascops asio</i>	7	12/14/2014	✓
	Great Horned Owl	<i>Bubo virginianus</i>	9	12/18/2011	✓
	Snowy Owl	<i>Bubo scandiacus</i>	3	12/13/2013	✓
T	Barred Owl ²⁹	<i>Strix varia</i>	8	5/21/2015	✓
T	Long-eared Owl	<i>Asio otus</i>	13	10/29/1996	✓
Ebr, SCnb	Short-eared Owl	<i>Asio flammeus</i>	8	1/14/2017	✓
	Northern Saw-whet Owl	<i>Aegolius acadicus</i>	53	11/5/2012	✓
<u>Kingfishers & Allies</u>					
	Belted Kingfisher	<i>Megaceryle alcyon</i>	15	10/13/2013	✓
<u>Woodpeckers</u>					
	Red-headed Woodpecker	<i>Melanerpes erythrocephalus</i>	15	5/6/2015	✓
	Red-bellied Woodpecker	<i>Melanerpes carolinus</i>	50	10/5/2012	✓
	Yellow-bellied Sapsucker	<i>Sphyrapicus varius</i>	40	10/2/2010	✓
	Downy Woodpecker	<i>Picoides pubescens</i>	14	12/16/2012	✓
	Hairy Woodpecker	<i>Picoides villosus</i>	8	4/10/2011	✓
	Northern Flicker	<i>Colaptes auratus</i>	1350	9/30/2007	✓
	Pileated Woodpecker	<i>Dryocopus pileatus</i>	1	6/4/2016	✓
<u>Caracaras & Falcons</u>					
RV	Crested Caracara	<i>Caracara cheriway</i>	1	5/5/2014	✓
T	American Kestrel	<i>Falco sparverius</i>	2467	10/2/1986	✓
	Merlin	<i>Falco columbarius</i>	800	9/30/1999	✓
Ebr, SCnb	Peregrine Falcon	<i>Falco peregrines</i>	241	10/5/2005	✓
<u>Perching Birds</u>					
	Olive-sided Flycatcher	<i>Contopus cooperi</i>	3	9/10/2010	
	Eastern Wood-Pewee	<i>Contopus virens</i>	34	5/23/2009	✓
	Yellow-bellied Flycatcher	<i>Empidonax flaviventris</i>	4	9/5/2010	✓
	Acadian Flycatcher	<i>Empidonax virescens</i>	23	5/12/2016	✓
	Alder Flycatcher	<i>Empidonax alnorum</i>	4	9/15/2010	✓
	Willow Flycatcher	<i>Empidonax traillii</i>	8	5/25/2014	✓
SCbr	Least Flycatcher	<i>Empidonax minimus</i>	10	9/9/2009	✓
	Eastern Phoebe	<i>Sayornis phoebe</i>	150	10/19/2005	✓
RV	Say's Phoebe	<i>Sayornis saya</i>	1	9/3/2013	
RV	Vermilion Flycatcher	<i>Pyrocephalus rubinus</i>	1	10/14/2014	✓
RV	Ash-throated Flycatcher	<i>Myiarchus cinerascens</i>	1	12/20/2015	✓
	Great Crested Flycatcher	<i>Myiarchus crinitus</i>	26	8/28/2007	✓
	Western Kingbird	<i>Tyrannus verticalis</i>	3	10/21/2007	✓

²⁹ Barred owl nesting has been documented in West Cape May (Michael O'Brien, personal communication, May 15, 2018).

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
	Eastern Kingbird	<i>Tyrannus tyrannus</i>	908	8/30/2011	✓
RV	Scissor-tailed Flycatcher	<i>Tyrannus forficatus</i>	1	6/17/2015	✓
RV	Fork-tailed Flycatcher	<i>Tyrannus savana</i>	1	10/26/2014	✓
RV	Gray Kingbird	<i>Tyrannus dominicensis</i>	1	10/5/2011	✓
RV	Sulphur-bellied Flycatcher	<i>Myiodynastes luteiventris</i>	1	10/13/2006	
	Northern Shrike	<i>Lanius excubitor</i>	1	12/28/2012	✓
Enb	Loggerhead shrike	<i>Lanius ludovicianus</i>	2	9/11/1957	✓
	White-eyed Vireo	<i>Vireo griseus</i>	35	5/9/2009	✓
RV	Bell's Vireo	<i>Vireo bellii</i>	3	10/13/2015	✓
	Yellow-throated Vireo	<i>Vireo flavifrons</i>	7	5/2/2016	✓
SCbr	Blue-headed Vireo	<i>Vireo solitarius</i>	9	4/24/2010	✓
	Warbling Vireo	<i>Vireo gilvus</i>	4	9/8/2008	✓
	Philadelphia Vireo	<i>Vireo philadelphicus</i>	6	9/18/2008	✓
	Red-eyed Vireo	<i>Vireo olivaceus</i>	110	9/26/2011	✓
	Blue Jay	<i>Cyanocitta cristata</i>	1500	10/8/2010	✓
	American Crow	<i>Corvus brachyrhynchos</i>	1394	12/17/2006	✓
	Fish Crow	<i>Corvus ossifragus</i>	875	9/7/2011	✓
	Common Raven	<i>Corvus corax</i>	1	6/16/2016	✓
Tbr, SCnb	Horned Lark	<i>Eremophila alpestris</i>	65	12/28/1947	✓
	Purple Martin	<i>Progne subis</i>	3401	8/21/2015	✓
RV	Brown-chested Martin	<i>Progne tapera</i>	1	11/18/1997	✓
	Tree Swallow	<i>Tachycineta bicolor</i>	100000	9/26/2015	✓
RV	Violet-green Swallow	<i>Tachycineta thalassina</i>	1	9/19/2011	✓
	Northern Rough-winged Swallow	<i>Stelgidopteryx serripennis</i>	175	10/17/2014	✓
	Bank Swallow	<i>Riparia riparia</i>	100	5/13/2007	✓
SCbr	Cliff Swallow	<i>Petrochelidon pyrrhonota</i>	1500	9/12/1993	✓
	Cave Swallow	<i>Petrochelidon fulva</i>	662	11/14/2015	✓
	Barn Swallow	<i>Hirundo rustica</i>	4500	9/3/2015	✓
	Carolina Chickadee	<i>Poecile carolinensis</i>	75	1/2/1950	✓
	Black-capped Chickadee	<i>Poecile atricapillus</i>	1	2/11/2017	
	Tufted Titmouse	<i>Baeolophus bicolor</i>	140	4/6/2010	✓
	Red-breasted Nuthatch	<i>Sitta canadensis</i>	427	10/5/2012	✓
	White-breasted Nuthatch	<i>Sitta carolinensis</i>	23	10/21/2012	✓
RV	Brown-headed Nuthatch	<i>Sitta pusilla</i>	1	6/30/2016	
	Brown Creeper	<i>Certhia americana</i>	36	10/15/2015	✓
RV	Rock Wren	<i>Salpinctes obsoletus</i>	1	3/8/1993	
	House Wren	<i>Troglodytes aedon</i>	28	5/10/2013	✓
SCbr	Winter Wren	<i>Troglodytes hiemalis</i>	13	12/18/2011	✓
E	Sedge Wren	<i>Cistothorus platensis</i>	5	8/8/1988	✓
	Marsh Wren	<i>Cistothorus palustris</i>	35	6/24/2011	✓
	Carolina Wren	<i>Thryothorus ludovicianus</i>	58	4/24/2008	✓
	Blue-gray Gnatcatcher	<i>Poliophtila caerulea</i>	313	8/16/2012	✓
	Golden-crowned Kinglet	<i>Regulus satrapa</i>	232	10/15/2015	✓
	Ruby-crowned Kinglet	<i>Regulus calendula</i>	650	10/25/2014	✓
RV	Northern Wheatear	<i>Oenanthe oenanthe</i>	1	10/9/2012	
	Eastern Bluebird	<i>Sialia sialis</i>	1000	11/18/2008	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
RV	Mountain Bluebird	<i>Sialia currucoides</i>	1	11/12/1988	
RV	Townsend's Solitaire	<i>Myadestes townsendi</i>	1	10/31/2015	✓
SCbr	Veery	<i>Catharus fuscescens</i>	150	9/8/2009	✓
SCnb	Gray-cheeked Thrush	<i>Catharus minimus</i>	300	10/5/2008	✓
	Bicknell's Thrush	<i>Catharus bicknelli</i>	12	10/5/2008	✓
	Swainson's Thrush	<i>Catharus ustulatus</i>	475	10/5/2008	✓
	Hermit Thrush	<i>Catharus guttatus</i>	1000	10/30/2010	✓
SCbr	Wood Thrush	<i>Hylocichla mustelina</i>	50	5/3/2012	✓
	American Robin	<i>Turdus migratorius</i>	250000	11/3/2013	✓
RV	Varied Thrush	<i>Ixoreus naevius</i>	1	1/30/2016	
	Gray Catbird	<i>Dumetella carolinensis</i>	1053	10/5/2015	✓
	Northern Mockingbird	<i>Mimus polyglottos</i>	55	5/9/2009	✓
SCbr	Brown Thrasher	<i>Toxostoma rufum</i>	35	10/10/2008	✓
int	European Starling	<i>Sturnus vulgaris</i>	5000	11/25/2013	✓
	American Pipit	<i>Anthus rubescens</i>	410	11/14/2012	✓
RV	Bohemian Waxwing	<i>Bombycilla garrulus</i>	1	2/10/2004	
	Cedar Waxwing	<i>Bombycilla cedrorum</i>	5000	11/2/2012	✓
	Lapland Longspur	<i>Calcarius lapponicus</i>	5	2/1/2014	✓
RV	Smith's Longspur	<i>Calcarius pictus</i>	1	3/20/2014	
	Snow Bunting	<i>Plectrophenax nivalis</i>	140	12/14/2007	✓
	Ovenbird	<i>Seiurus aurocapilla</i>	186	4/24/2011	✓
SCbr	Worm-eating Warbler	<i>Helmitheros vermivorum</i>	51	5/1/2012	✓
	Louisiana Waterthrush	<i>Parkesia motacilla</i>	6	4/9/2005	✓
	Northern Waterthrush	<i>Parkesia noveboracensis</i>	200	9/4/2014	✓
Ebr,SCnb	Golden-winged Warbler	<i>Vermivora chrysoptera</i>	2	9/7/2014	✓
	Blue-winged Warbler	<i>Vermivora cyanoptera</i>	15	5/2/2007	✓
	Black-and-white Warbler	<i>Mniotilta varia</i>	676	9/15/2015	✓
	Prothonotary Warbler	<i>Protonotaria citrea</i>	6	5/4/2012	✓
RV	Swainson's Warbler	<i>Limnithlypis swainsonii</i>	1	5/5/2016	
	Tennessee Warbler	<i>Oreothlypis peregrina</i>	100	9/17/1997	✓
	Orange-crowned Warbler	<i>Oreothlypis celata</i>	6	12/21/2016	✓
SCbr	Nashville Warbler	<i>Oreothlypis ruficapilla</i>	30	9/14/1997	✓
	Connecticut Warbler	<i>Oporornis agilis</i>	8	9/13/2007	✓
RV	MacGillivray's Warbler	<i>Geothlypis tolmiei</i>	1	1/10/1998	✓
	Mourning Warbler	<i>Geothlypis philadelphia</i>	4	8/17/2012	✓
SC	Kentucky Warbler	<i>Geothlypis formosa</i>	4	5/13/2010	✓
	Common Yellowthroat	<i>Geothlypis trichas</i>	1200	10/5/2006	✓
SCbr	Hooded Warbler	<i>Setophaga citrina</i>	12	4/29/2007	✓
	American Redstart	<i>Setophaga ruticilla</i>	4065	9/15/2015	✓
	Cape May Warbler	<i>Setophaga tigrina</i>	150	9/30/1996	✓
SCbr	Cerulean Warbler	<i>Setophaga cerulea</i>	2	5/14/2016	✓
SCbr	Northern Parula	<i>Setophaga americana</i>	800	10/6/2006	✓
	Magnolia Warbler	<i>Setophaga magnolia</i>	73	9/29/2007	✓
	Bay-breasted Warbler	<i>Setophaga castanea</i>	17	9/24/2012	✓
SCbr	Blackburnian Warbler	<i>Setophaga fusca</i>	22	9/8/2008	✓
	Yellow Warbler	<i>Setophaga petechia</i>	365	8/14/2014	✓
	Chestnut-sided Warbler	<i>Setophaga pensylvanica</i>	30	8/27/2010	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
	Blackpoll Warbler	<i>Setophaga striata</i>	1068	9/29/2012	✓
SCbr	Black-throated Blue Warbler	<i>Setophaga caeruleascens</i>	500	10/6/2006	✓
	Palm Warbler	<i>Setophaga palmarum</i>	1500	10/10/2014	✓
	Pine Warbler	<i>Setophaga pinus</i>	128	4/6/2010	✓
	Yellow-rumped Warbler	<i>Setophaga coronata</i>	63640	10/29/2010	✓
	Yellow-throated Warbler	<i>Setophaga dominica</i>	32	5/12/2016	✓
	Prairie Warbler	<i>Setophaga discolor</i>	20	5/6/2011	✓
RV	Black-throated Gray Warbler	<i>Setophaga nigrescens</i>	1	8/29/2014	
RV	Townsend's Warbler	<i>Setophaga townsendi</i>	1	3/7/2013	✓
SCbr	Black-throated Green Warbler	<i>Setophaga virens</i>	90	9/29/2007	✓
SCbr	Canada Warbler	<i>Cardellina canadensis</i>	9	5/12/2012	✓
	Wilson's Warbler	<i>Cardellina pusilla</i>	5	5/23/2015	✓
SCbr	Yellow-breasted Chat	<i>Icteria virens</i>	12	5/21/2015	✓
RV	Spotted Towhee	<i>Pipilo maculatus</i>	1	4/5/1995	
	Eastern Towhee	<i>Pipilo erythrophthalmus</i>	50	4/29/2007	✓
	American Tree Sparrow	<i>Spizella arborea</i>	100	12/26/1938	✓
	Chipping Sparrow	<i>Spizella passerina</i>	1422	10/21/2007	✓
	Clay-colored Sparrow	<i>Spizella pallida</i>	6	10/21/2007	✓
	Field Sparrow	<i>Spizella pusilla</i>	100	10/22/2005	✓
Ebr, SCnb	Vesper Sparrow	<i>Poocetes gramineus</i>	35	12/26/1938	✓
	Lark Sparrow	<i>Chondestes grammacus</i>	3	9/11/1999	✓
RV	Lark Bunting	<i>Calamospiza malanocorys</i>	1	6/2/2008	
Tbr	Savannah Sparrow	<i>Passerculus sandwichensis</i>	600	10/5/2006	✓
Tbr, SCnb	Grasshopper Sparrow	<i>Ammodramus savannarum</i>	6	10/5/2008	✓
E	Henslow's sparrow	<i>Ammodramus henslowii</i>	10	7/28/1929	✓
RV	Le Conte's Sparrow	<i>Ammodramus leconteii</i>	1	1/10/2016	
	Nelson's Sparrow	<i>Ammodramus nelsoni</i>	20	10/21/2016	✓
SCbr	Saltmarsh Sparrow	<i>Ammodramus caudacutus</i>	37	7/13/2011	✓
	Seaside Sparrow	<i>Ammodramus maritimus</i>	60	7/13/2011	✓
	Fox Sparrow	<i>Passerella iliaca</i>	143	2/13/2010	✓
	Song Sparrow	<i>Melospiza melodia</i>	2000	10/30/2010	✓
	Lincoln's Sparrow	<i>Melospiza lincolni</i>	9	10/12/2013	✓
	Swamp Sparrow	<i>Melospiza georgiana</i>	10000	10/30/2010	✓
	White-throated Sparrow	<i>Zonotrichia albicollis</i>	1500	10/30/2010	✓
RV	Harris's Sparrow	<i>Zonotrichia querula</i>	1	11/4/1998	
	White-crowned Sparrow	<i>Zonotrichia leucophrys</i>	60	11/19/2008	✓
RV	Golden-crowned Sparrow	<i>Zonotrichia atricapilla</i>	1	2/21/1998	
	Dark-eyed Junco	<i>Junco hyemalis</i>	3397	10/29/2010	✓
	Summer Tanager	<i>Piranga rubra</i>	7	5/10/2016	✓
	Scarlet Tanager	<i>Piranga olivacea</i>	50	9/30/1996	✓
RV	Western Tanager	<i>Piranga ludoviciana</i>	2	12/25/2012	✓
	Northern Cardinal	<i>Cardinalis cardinalis</i>	87	4/24/2008	✓
	Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>	50	9/24/1983	✓
RV	Black-headed Grosbeak	<i>Pheucticus melanocephalus</i>	1	3/6/1999	✓

State Status	Common name	Scientific name	Cape May High Count*	Cape May High Count Date*	Recorded in West Cape May**
	Blue Grosbeak	<i>Passerina caerulea</i>	60	5/12/2007	✓
	Indigo Bunting	<i>Passerina cyanea</i>	150	9/30/1996	✓
RV	Painted Bunting	<i>Passerina ciris</i>	1	9/15/2015	✓
	Dickcissel	<i>Spiza americana</i>	12	10/10/2015	✓
Tbr, SCnb	Bobolink ³⁰	<i>Dolichonyx oryzivorus</i>	7500	9/4/2014	✓
	Red-winged Blackbird	<i>Agelaius phoeniceus</i>	70000	11/19/2013	✓
SCbr	Eastern Meadowlark	<i>Sturnella magna</i>	145	10/22/2007	✓
	Yellow-headed Blackbird	<i>Xanthocephalus xanthocephalus</i>	2	9/10/2011	✓
	Rusty Blackbird	<i>Euphagus carolinus</i>	250	11/19/2013	✓
RV	Brewer's Blackbird	<i>Euphagus cyanocephalus</i>	8	2/12/1990	✓
	Common Grackle	<i>Quiscalus quiscula</i>	17000	12/2/2016	✓
	Boat-tailed Grackle	<i>Quiscalus major</i>	700	10/15/2014	✓
int	Brown-headed Cowbird	<i>Molothrus ater</i>	5000	10/28/2007	✓
	Orchard Oriole	<i>Icterus spurius</i>	30	8/7/2015	✓
RV	Bullock's Oriole	<i>Icterus bullockii</i>	1	5/11/2016	✓
	Baltimore Oriole	<i>Icterus galbula</i>	420	9/4/1997	✓
	Purple Finch	<i>Haemorhous purpureus</i>	1000	11/3/2012	✓
int	House Finch	<i>Haemorhous mexicanus</i>	3000	10/31/2006	✓
	Red Crossbill	<i>Loxia curvirostra</i>	166	11/16/2012	✓
	White-winged Crossbill	<i>Loxia leucoptera</i>	360	11/3/2012	✓
	Common Redpoll	<i>Acanthis flammea</i>	30	12/29/2003	✓
RV	European Goldfinch	<i>Carduelis carduelis</i>	1	5/9/2007	
	Pine Siskin	<i>Spinus pinus</i>	4740	11/2/2012	
	American Goldfinch	<i>Spinus tristis</i>	9000	11/17/2008	✓
	Evening Grosbeak	<i>Coccothraustes vespertinus</i>	45	11/8/2012	✓
int	House Sparrow	<i>Passer domesticus</i>	394	12/18/2016	✓
RV	Eurasian Tree Sparrow	<i>Passer montanus</i>	1	4/9/2014	

*Data from Sullivan, B.L., C.L. Wood, M.J. Iliff, R.E. Bonney, D. Fink, and S. Kelling. 2009. eBird: a citizen-based bird observation network in the biological sciences. *Biological Conservation* 142: 2282-2292. <http://ebird.org/ebird/nj/places>. Website accessed March 2017.

**Michael O'Brien, personal communication, May 15, 2018.

³⁰ Bobolink nesting has not been documented in West Cape May, but migrants use the borough extensively (Michael O'Brien, personal communication, May 15, 2018).

C.2 List of New Jersey Land Mammals

Mammals have not been inventoried specifically in West Cape May Borough or Cape May County.

Status	Common Name	Scientific Name
	Opossum	<i>Didelphis marsupialis</i>
	Masked shrew	<i>Sorex cinereus</i>
	Tuckahoe masked shrew	<i>Sorex cinereus nigriculus</i>
	Water shrew	<i>Sorex palustris</i>
	Smokey shrew	<i>Sorex fumeus</i>
	Long-tailed shrew	<i>Sorex dispar</i>
	Short-tailed shrew	<i>Blarina brevicauda</i>
	Least shrew	<i>Cryptotis parva</i>
	Pygmy shrew	<i>Sorex hoyii</i>
	Hairy-tailed mole	<i>Parascalops breweri</i>
	Eastern mole	<i>Scalopus aquaticus</i>
	Star-nosed mole	<i>Condylura cristata</i>
	Little brown bat	<i>Myotis lucifugus</i>
E	Indiana bat	<i>Myotis sodalis</i>
	Keen myotis	<i>Myotis septentrionalis</i>
	Small-footed myotis	<i>Myotis leibii</i>
	Silver-haired bat	<i>Lasionycteris noctivagans</i>
	Eastern pipistrel	<i>Pipistrellus subflavus</i>
	Big brown bat	<i>Eptesicus fuscus</i>
	Red bat	<i>Lasiurus borealis</i>
peripheral	Northern yellow bat	<i>Lasiurus intermedius</i>
	Hoary bat	<i>Lasiurus cinereus</i>
	Eastern cottontail	<i>Sylvilagus floridanus</i>
	New England cottontail	<i>Sylvilagus transitionalis</i>
int	European hare	<i>Lepus capensis</i>
int	Black-tailed jackrabbit	<i>Lepus californicus</i>
int	White-tailed jackrabbit	<i>Lepus townsendii</i>
	Eastern chipmunk	<i>Tamias striatus</i>
	Woodchuck	<i>Marmota monax</i>
	Gray squirrel	<i>Sciurus carolinensis</i>
	Red squirrel	<i>Tamiasciurus hudsonicus</i>
	Southern flying squirrel	<i>Glaucomys volans</i>
	Northern flying squirrel	<i>Glaucomys sabrinus</i>
	Beaver	<i>Castor canadensis</i>
int	Nutria	<i>Myocastor coypus</i>
	Marsh rice rat	<i>Oryzomys palustris</i>
	White-footed mouse	<i>Peromyscus leucopus</i>
E	Eastern woodrat	<i>Neotoma floridana</i>
	Red-backed mouse	<i>Clethrionomys gapperi</i>
	Meadow vole	<i>Microtus pennsylvanicus</i>

Status	Common Name	Scientific Name
	Woodland vole	<i>Microtus pinetorum</i>
	Muskrat	<i>Ondatra zibethicus</i>
	Southern bog lemming	<i>Synaptomys cooperi</i>
int	Black rat	<i>Rattus rattus</i>
int	Brown rat	<i>Rattus norvegicus</i>
int	House mouse	<i>Mus musculus</i>
	Woodland jumping mouse	<i>Napaeozapus insignis</i>
	Meadow jumping mouse	<i>Zapus hudsonius</i>
	Porcupine	<i>Erethizon dorsatum</i>
	Eastern coyote	<i>Canis latrans, var</i>
	Red fox	<i>Vulpes vulpes</i>
	Gray fox	<i>Urocyon cinereoargenteus</i>
	Black bear	<i>Ursus americanus</i>
	Raccoon	<i>Procyon lotor</i>
	Ermine	<i>Mustela erminea</i>
	Long-tailed weasel	<i>Mustela frenata</i>
	Mink	<i>Mustela vison</i>
	Striped skunk	<i>Mephitis mephitis</i>
	River otter	<i>Lutra canadensis</i>
E	Bobcat	<i>Felis rufus</i>
	White-tailed deer	<i>Odocoileus virginianus</i>
Source: NJDEP Division of Fish and Wildlife. 2004b. <u>Checklist of NJ Mammals.</u> http://www.state.nj.us/dep/fgw/chkmamls.htm		

C.3 Cape May County Reptiles

New Jersey reptile species that may occur in Cape May County.

Status	Common Name	Scientific Name
<u>Lizards</u>		
	Common five-lined skink	<i>Plestiodon fasciatus</i>
	Eastern fence lizard	<i>Sceloporus undulatus</i>
<u>Snakes</u>		
	Eastern wormsnake	<i>Carphophis a. amoenus</i>
	Northern black racer	<i>Coluber c. constrictor</i>
	Northern ring-necked snake	<i>Diadophis punctatus edwardsi</i>
	Southern ring-necked snake	<i>Diadophis p. punctatus</i>
	Eastern hog-nosed snake	<i>Heterodon platirhinos</i>
SC	Eastern kingsnake	<i>Lampropeltis getula</i>
	Eastern milksnake	<i>Lampropeltis t. triangulum</i>
	Coastal plain milksnake	<i>Lampropeltis t. triangulum x L. t. elapsoides</i>
	Northern watersnake	<i>Nerodia s. sipedon</i>
	Rough greensnake	<i>Opheodrys aestivus</i>
E	Red cornsnake	<i>Pantherophis guttatus</i>
	Eastern ratsnake	<i>Pantherophis alleghaniensis</i>
	Northern brownsnake	<i>Storeria d. dekayi</i>
	Northern red-bellied snake	<i>Storeria o. occipitomaculata</i>
	Eastern ribbonsnake	<i>Thamnophis s. sauritus</i>
	Eastern gartersnake	<i>Thamnophis s. sirtalis</i>
	Eastern smooth earthsnake	<i>Virginia v. valeriae</i>
<u>Turtles</u>		
	Snapping turtle	<i>Chelydra serpentina</i>
	Eastern painted turtle	<i>Chrysemys p. picta</i>
SC	Spotted turtle	<i>Clemmys guttata</i>
E	Bog turtle	<i>Glyptemys muhlenbergii</i>
	Southeastern mud turtle	<i>Kinosternon s. subrubrum</i>
	Northern diamond-backed terrapin	<i>Malaclemys t. terrapin</i>
	Northern red-bellied cooter	<i>Pseudemys rubiventris</i>
	Eastern musk turtle	<i>Sternotherus odoratus</i>
SC	Woodland box turtle	<i>Terrapene c. carolina</i>
int	Red-eared slider	<i>Trachemys scripta elegans</i>
Source: NJDEP Division of Fish and Wildlife, June 6 2014.		

C.4 Cape May County Amphibians

New Jersey amphibian species that may occur in Cape May County.

Status	Common Name	Scientific Name
<u>Anurans</u>		
	Eastern cricket frog	<i>Acris crepitans</i>
SC	Fowler's toad	<i>Anaxyrus fowleri</i>
E	Cope's gray treefrog	<i>Hyla chrysoscelis</i>
	Barking treefrog	<i>Hyla gratiosa</i>
	Northern gray treefrog	<i>Hyla versicolor</i>
	Bullfrog	<i>Lithobates catesbeianus</i>
	Green frog	<i>Lithobates clamitans</i>
	Pickerel frog	<i>Lithobates palustris</i>
	Southern leopard frog	<i>Lithobates sphenoccephalus</i>
	Wood frog	<i>Lithobates sylvaticus</i>
SC	Carpenter frog	<i>Lithobates virgatipes</i>
	Spring peeper	<i>Pseudacris crucifer</i>
	New Jersey chorus frog	<i>Pseudacris kalmi</i>
	Eastern spadefoot toad	<i>Scaphiopus h. holbrooki</i>
<u>Salamanders</u>		
SC	Marbled salamander	<i>Ambystoma opacum</i>
E	Eastern tiger salamander	<i>Ambystoma tigrinum</i>
	Four-toed salamander	<i>Hemidactylium scutatum</i>
	Red-spotted newt	<i>Notophthalmus v. viridescens</i>
	Red-backed salamander	<i>Plethodon cinereus</i>
T	Eastern mud salamander	<i>Pseudotriton m. montanus</i>
	Northern red salamander	<i>Pseudotriton r. ruber</i>
Sources: NJDEP Division of Fish and Wildlife, June 6 2014; NJENSP November 2014.		

C.5 Freshwater Fish of New Jersey

State Status	Common Name	Scientific Name
<u>Northern Lampreys</u>		
	American Brook Lamprey	<i>Lampetra appendix</i>
	Sea Lamprey	<i>Petromyzon marinus</i>
<u>Sturgeons</u>		
E	Atlantic Sturgeon	<i>Acipenser oxyrhynchus</i>
E	Shortnose Sturgeon	<i>Acipenser brevirostrum</i>
<u>Gars</u>		
X	Longnose Gar	<i>Lepisosteus osseus</i>
<u>Bowfins</u>		
	Bowfin	<i>Amia calva</i>
<u>Freshwater Eels</u>		
	American Eel	<i>Anguilla rostrata</i>
<u>Swamp Eels</u>		
int*	Swamp Eel	<i>Monopterus albus</i>
<u>Herrings, Shads, Sardines, Menhadens</u>		
	Blueback Herring	<i>Alosa aestivalis</i>
	Hickory Shad	<i>Alosa mediocris</i>
	Alewife	<i>Alosa pseudoharengus</i>
	American Shad	<i>Alosa sapidissima</i>
	Gizzard Shad	<i>Dorosoma cepedianum</i>
<u>Salmonids</u>		
int	Rainbow Trout	<i>Oncorhynchus mykiss</i>
int	Brown Trout	<i>Salmo trutta</i>
	Brook Trout	<i>Salvelinus fontinalis</i>
int	Lake Trout	<i>Salvelinus namaycush</i>
<u>Smelts</u>		
	Rainbow Smelt	<i>Osmerus mordax</i>
<u>Mudminnows</u>		
	Eastern Mudminnow	<i>Umbra pygmaea</i>
<u>Pikes</u>		
	Redfin Pickerel	<i>Esox americanus</i>

State Status	Common Name	Scientific Name
int	Northern Pike	<i>Esox lucius</i>
	Chain Pickerel	<i>Esox niger</i>
int	Muskellunge	<i>Esox masquinongy</i>
<u>Carp, Minnows</u>		
int	Goldfish	<i>Carassius auratus</i>
int*	Grass Carp	<i>Ctenopharyngodon idella</i>
	Satinfin Shiner	<i>Cyprinella analostana</i>
	Spotfin Shiner	<i>Cyprinella spiloptera</i>
int	Common Carp	<i>Cyprinus carpio</i>
	Cutlip Minnow	<i>Exoglossum maxillingua</i>
	Eastern Silvery Minnow	<i>Hybognathus regius</i>
int*	Bighead Carp	<i>Hypophthalmichthys nobilis</i>
	Common Shiner	<i>Luxilus cornutus</i>
int	Allegheny Pearl Dace	<i>Margariscus margarita</i>
	Golden Shiner	<i>Notemigonus crysoleucas</i>
	Comely Shiner	<i>Notropis amoenus</i>
	Bridle Shiner	<i>Notropis bifrenatus</i>
	Ironcolor Shiner	<i>Notropis chalybaeus</i>
	Spottail Shiner	<i>Notropis hudsonius</i>
	Swallowtail Shiner	<i>Notropis procne</i>
int	Fathead Minnow	<i>Pimephales promelas</i>
int	Bluntnose Minnow	<i>Pimephales notatus</i>
	Blacknose Dace	<i>Rhinichthys atratulus</i>
	Longnose Dace	<i>Rhinichthys cataractae</i>
	Creek Chub	<i>Semotilus atromaculatus</i>
	Fallfish	<i>Semotilus corporalis</i>
<u>Suckers</u>		
	Quillback	<i>Carpiodes cyprinus</i>
	White Sucker	<i>Catostomus commersoni</i>
	Creek Chubsucker	<i>Erimyzon oblongus</i>
	Northern Hog Sucker	<i>Hypentelium nigricans</i>
<u>Freshwater Catfishes</u>		
	White Catfish	<i>Ameiurus catus</i>
int	Black Bullhead	<i>Ameiurus melas</i>
	Yellow Bullhead	<i>Ameiurus natalis</i>
	Brown Bullhead	<i>Ameiurus nebulosus</i>
int	Channel Catfish	<i>Ictalurus punctatus</i>
	Tadpole Madtom	<i>Noturus gyrinus</i>
	Margined Madtom	<i>Noturus insignis</i>
int*	Flathead Catfish	<i>Pylodictis olivarius</i>

State Status	Common Name	Scientific Name
	<u>Pirate Perches</u>	
	Pirate Perch	<i>Aphredoderus sayanus</i>
	<u>Killifishes</u>	
	Banded Killifish	<i>Fundulus diaphanus</i>
	Mummichog	<i>Fundulus heteroclitus</i>
	<u>Poeciliids</u>	
	Eastern Mosquitofish	<i>Gambusia holbrooki</i>
int	Mosquitofish	<i>Gambusia affinis</i>
	<u>Gasterosteidae:</u>	
	Fourspine Stickleback	<i>Apletes quadracus</i>
int*	Brook Stickleback	<i>Culaea inconstans</i>
	Threespine Stickleback	<i>Gasterosteus aculeatus</i>
	Ninespine Stickleback	<i>Pungitius pungitius</i>
	<u>Moronidae:</u>	
	White Perch	<i>Morone americana</i>
	Striped Bass	<i>Morone saxatilis</i>
	<u>Sticklebacks, Tubesnouts</u>	
	Mud Sunfish	<i>Acantharchus pomotis</i>
int	Rock Bass	<i>Ambloplites rupestris</i>
	Blackbanded Sunfish	<i>Enneacanthus chaetodon</i>
	Bluespotted Sunfish	<i>Enneacanthus gloriosus</i>
	Banded Sunfish	<i>Enneacanthus obesus</i>
int*	Green Sunfish	<i>Lepomis cyanellus</i>
	Pumpkinseed	<i>Lepomis gibbosus</i>
int	Bluegill	<i>Lepomis macrochirus</i>
	Redbreast Sunfish	<i>Lepomis auritus</i>
int*	Warmouth	<i>Lepomis gulosus</i>
int	Smallmouth Bass	<i>Micropterus dolomieu</i>
int	Largemouth Bass	<i>Micropterus salmoides</i>
int	White Crappie	<i>Pomoxis annularis</i>
int	Black Crappie	<i>Pomoxis nigromaculatus</i>
	<u>Perches</u>	
	Swamp Darter	<i>Etheostoma fusiforme</i>
	Tessellated Darter	<i>Etheostoma olmstedii</i>
	Yellow Perch	<i>Perca flavescens</i>
	Shield Darter	<i>Percina peltata</i>

State Status	Common Name	Scientific Name
int	Walleye	<i>Sander vitreus</i>
<u>Sculpins</u>		
	Slimy Sculpin	<i>Cottus cognatus</i>
<u>Loaches</u>		
int*	Oriental Weatherfish	<i>Misgurnus anguillicaudatus</i>
<u>Soles</u>		
	Hogchoker	<i>Trinectes maculatus</i>
<p>* indicates species that pose a serious threat to freshwater resources, and must be destroyed when encountered. An anticipated addition to this category, the silver carp (<i>Hypophthalmichthys molitrix</i>) has not yet been documented in NJ.</p> <p>Source: NJDEP Division of Fish and Wildlife. 2016. <u>Checklist of NJ Fish</u>. http://www.njfishandwildlife.com/chkfish.htm</p>		

APPENDIX D: ENDANGERED SPECIES

Contents:

D.1 Rare Plant Reporting Form

Source: http://www.nj.gov/dep/parksandforests/natural/heritage/natherrareplantspeciesreportform1_2008.doc

Note: Use the following address, not the one on the reporting form:

The New Jersey Natural Heritage Program
DEP - Office of Natural Lands Management
Mail Code 501-04
P.O. Box 420
Trenton, New Jersey 08625-0420

D.2 Rare Wildlife Reporting Form

Source: <http://www.state.nj.us/dep/fgw/ensp/pdf/rptform.pdf>

D.3. Habitat Requirements for Rare Wildlife Species Recorded in the Vicinity of West Cape May Borough

D.4. Rare Bird Species Documented in Cape May County

D.5 List of Rare Plant Species of Cape May County

Source: <http://www.state.nj.us/dep/parksandforests/natural/heritage/textfiles/hunterdon.pdf>

D.6 Cape May Migratory Bird Refuge Natural Heritage Priority Site



Natural Heritage Rare Plant Species Reporting Form

This form is used to report a personal field sighting of a rare plant species tracked by the Natural Heritage Database. It may also be used to summarize locational information from a published or unpublished report. Plant species tracked include those appearing on the State Endangered Plant Species List or the Plant Species of Concern List (<http://www.nj.gov/dep/parksandforests/natural/heritage/spplant.html>). The Office of Natural Lands Management can provide copies of the lists upon request. In order for this form to be processed, the sections preceded by an asterisk (*) must be completed.

Send completed form to: DEP, Division of Parks and Forestry, Office of Natural Lands Management, Natural Heritage Program, P.O. Box 404, Trenton, NJ 08625-0404.

Today's Date: (date this form is being completed) _____

Common Name: _____ **Scientific Name:** _____

***Location Map:** *A mapped location of the occurrence must accompany this form.* The ideal format is to locate the site on a photocopied section of a U.S. Geological Survey 7.5 minute topographical map, and to also sketch a second map showing finer details. Be sure to provide the name of the USGS map.

GPS Coordinates (If available please provide the following):

Datum Used:	<input type="checkbox"/> NAD 1983	<input type="checkbox"/> NAD 1927	<input type="checkbox"/> WGS84	<input type="checkbox"/> Other _____
Lat/Long (if applicable):	_____ N (Latitude)		_____ W (Longitude)	
UTM (if applicable)	18 N/S: _____	_____ Northing	_____ Easting	
Accuracy Level:	+/- _____ <input type="checkbox"/> feet or <input type="checkbox"/> meters			

***Directions to Site:** Directions to the element occurrence using a readily locatable and relatively permanent landmark on or near the site (such as a road intersection, a prominent hill or cliff) as the starting point. Use clear, complete sentences so that someone who is unfamiliar with the area will be able to relocate the element occurrence using your written directions (e.g., "About 50 ft. N. of small stream draining Brindel Lake, 0.5 mi. SE of Brindeltown and 0.2 mi. WSW of jct. of Range Rd. and Rt. 539, Fort Dix").

***Date(s) of the Observation(s):** _____

Identification: How was the species identification made? Name the identification manuals used or the experts consulted. Were there identification problems?

***Number of Individuals Observed:**

- 1-10 11-50 101-1,000 1,001-10,000 >10,000

If possible, provide the exact number of individuals and an estimated percentage of flowering/fruitlet individuals. For rhizomatous plants such as grasses and sedges, what was counted as individual – separate culms or entire clumps or patches?

Life Stages Present: Check life stages observed or provide an estimate of the numbers of individuals for each life stage.

- vegetative in bud flower seed dispersing seedling dormant

Associated Species/Additional Biological Data: List any associated species and/or additional rare species observed at this site. What else was observed? Provide information on the general condition or vigor of the individuals and viability of the population(s).

--

Habitat Data: Describe the specific area where the occurrence is located. List natural community types, dominant vegetation and information on the physical environment such as substrate type, hydrology, moisture regime, slope and aspect. Also, describe the surrounding landscape.

--

Threats: Describe any current or potential threats to this occurrence. If invasive species are present, please list.

--

Ownership: If known, please provide landowner(s) name, address, phone #.

--

Information Source:

***Name, Address and Phone # (of person filing report):**

Name:	
Address:	
Phone Number:	

*Does this information come directly from a field visit or a published or unpublished report?

Citation: For information taken from a published or unpublished report, please provide a copy of the cover page and the pertinent portions of the report. If a copy can not be provided, list below the author, date, title, publisher, and page numbers.

--

Voucher: Was the observation vouchered with a photograph? a video/digital format? a specimen?
If possible, attach a copy of the photograph or tape. If specimen voucher, please provide the name of the repository:

--

Confirmation: Would you accompany a biologist to the site if needed? yes no.

Additional Comments: (use extra sheets if needed)

RARE WILDLIFE SIGHTING REPORT FORM

REPORT FORM MUST BE ACCOMPANIED BY AN AERIAL PHOTOGRAPH, SATELLITE IMAGE, OR TOPOGRAPHIC MAP WITH THE LOCATION PRECISELY MARKED. PLEASE PRINT LEGIBLY.

***The inclusion of a map is mandatory, please see other side for further information on obtaining a map.**

General Information

Today's Date _____

Common Name _____

Scientific Name (If known) _____

Where did the sighting take place?

Municipality/ Township _____

County _____

Topographic quad (if known) _____

Coordinates in state plane feet (if known) _____

Directions to location with landmarks, which will enable the future relocation of the site where the species was sighted:

Land Owner (name, address and phone number, if known) _____

Describe habitat at the point of sighting and habitat in the general area of the sighting location.

Would you accompany a biologist to the site if needed? Yes No

Can you describe any immediate or future plans to develop or disturb the site? Yes No

If so, please describe. _____

Locational Accuracy

1. Is your depiction of the sighting location on the topographic map or aerial photo within 6m (20ft) of the animals actual location on the ground? Yes No (if no, answer question 2 below)

2. Your mapping is accurate to within ___ meters ___ feet ___ miles of the actual location.

What was observed?

How was the species identification made? (ex. Sighting, Call, Road Kill, etc.) _____

Date and time of this sighting (ex. August 20, 2004, 10:30am) _____

How frequently has this species been sighted at this location and over how long a period of time? _____

Number of individuals sighted: Adult ___ Immature ___ Larva ___ Unknown/Other ___

Describe sighting and activity observed (ex. Nesting, Perched, Flying, Sunning, etc.) _____

Describe physical features that identify the sighted animal as the species you are reporting. _____

Were photos taken? Yes No Was video recorded? Yes No Was audio recorded? Yes No
(PHOTOS/VIDEO/AUDIO ARE STRONGLY ENCOURAGED IN ORDER TO VERIFY THE ACCURACY OF A SIGHTING. Items should be identified with the date taken, location, and observer signature. Items will not be returned.)

List manuals used or experts consulted to verify identification. _____

Provide a brief background on wildlife knowledge and/or experience, or additional information that would add to the validity of the sighting. _____

Can this be verified by someone else or can anyone vouch for your identification skills? Yes No _____

Describe any additional information that may be useful in regards to the condition of the animal or location. _____

Your Contact information

Name _____
Street _____
City _____ State _____ ZIP _____
Daytime Phone () _____ - _____ E-mail _____

Preferred method of contact _____

Signature _____



Return to:
Endangered and Nongame Species Program
NJ Division of Fish and Wildlife
PO Box 400
Trenton, NJ 08625-0400
(609) 292-9400



Instructions

1. Complete this form for first-hand field observations only.
2. **DO NOT COMPLETE THIS FORM** if the source of your information is a report, letter, conversation, or other document. Send us the documentation instead.
3. Attach a copy of a map. (*see below)
4. Only report one species at each location per form and map.

***Mapping**

A map is necessary to help our biologists determine if suitable habitat is present at the location. Once the suitability of the area is determined the map provided aids in the delineation of land to be protected. Ideally the most accurate form of map is an aerial photo, which can be obtained from <http://www.state.nj.us/dep/gis/newmapping.htm>, if you are comfortable with your ability to identify the location of the sighting accurately on them. In addition, satellite-derived images are available at <http://www.maps.google.com>. These images can be printed and clearly marked with a pen. An alternative to an aerial photo or satellite image is a topographic map. You may also print copies of topographic maps from the internet at <http://www.topozone.com>. Please use 1:24,000 scale topographic maps only. Please provide either an image or a topographic map, but **NOT** both. Thank you.

Refer to the DFW website for further information: <http://www.njfishandwildlife.com/ensp/rptform.htm>

D.3 Habitat Requirements for Rare Wildlife Species Recorded in the Immediate Vicinity of West Cape May Borough

Habitat notes are generally direct quotes, as excerpted from the cited sources.

Cope's Gray Treefrog

(State Endangered)

Cope's gray treefrogs require both aquatic and terrestrial habitats, relying on small freshwater ponds, old fields, and hardwood forests throughout their annual cycle. Accepting both natural and manmade basins, they breed in vernal ponds, gravel pits, retention basins, floodplain corridors, bogs, weedy lakes, marshes and farm ponds. Breeding pools must contain water long enough to ensure metamorphosis but often dry up by the end of summer, thereby prohibiting the establishment of predatory fish (Beans and Niles, 2003).

Eastern Tiger Salamander

(State Endangered)

Tiger salamanders require both upland and wetland habitats that contain suitable breeding ponds, forests, and soil types appropriate for burrowing (loamy sand and sandy loam). Ponds may have gravel, mud, sand or clay substrates and pH levels between 3.5 and 7.9, but they must contain clean, unpolluted water and be fish free. Terrestrial habitats utilized by the species include old fields and deciduous or mixed woods (Beans and Niles, 2003).

Bald Eagle

(Endangered breeding, Threatened non-breeding)

Bald Eagles live near rivers, lakes, and marshes where they can find fish, their staple food. Bald Eagles will also feed on waterfowl, turtles, rabbits, snakes, and other small animals and carrion. Bald Eagles require a good food base, perching areas, and nesting sites. Their habitat includes estuaries, large lakes, reservoirs, rivers, and some seacoasts. In winter, the birds congregate near open water in tall trees for spotting prey and night roosts for sheltering. (USFWS, 2015).

Barred Owl

(State Threatened)

Barred owls require mature, wet woods that contain large trees with cavities suitable for nesting. The lack of large nesting cavities is often the primary limiting factor for the birds. Their habitats typically have an open understory through which the owls can fly and hunt. Large, unbroken tracts that are remote from human habitation are preferred by the species (Beans and Niles, 2003).

Black Skimmer

(State Endangered)

Black skimmers nests on open, sandy beaches, inlets, sandbars, offshore islands and dredge disposal islands that are sparsely vegetated and contain shell fragments. Shallow tidal creeks, inlets and ponds are utilized for foraging. Similar coastal and estuarine habitats are used throughout the year (Beans and Niles, 2003).

Black-crowned Night Heron

(Threatened breeding, Special Concern non-breeding)

Forests, scrub/shrubland, marshes and ponds serve as nesting, roosting and foraging habitats for black-crowned night herons. The birds nest colonially, sometimes in mixed-species colonies, in

wooded swamps, coastal dune forests, vegetated dredge spoil islands, scrub thickets or marshes. The herons forage in marshes, along the edges of ponds and creeks, and in saline habitats including shallow tide pools, tidal channels and mudflats (Beans and Niles, 2003).

Bobolink

(Threatened breeding, Special Concern non-breeding)

Bobolinks inhabit low-intensity agricultural habitats, such as hayfields and pastures, during the breeding season. In addition, lush fallow fields and meadows of grasses, forbs, and wildflowers are occupied. Bobolink nests are often placed in areas of greatest vegetative height and density. Similar habitats are occupied by bobolinks throughout their annual cycle. During migration, bobolinks inhabit fallow and agricultural fields, as well as coastal and freshwater marshes. On their South American wintering grounds, they occur in grasslands, marshes, rice fields, and farm fields. (NJENSP, undated-a).

Brown Thrasher

(Special Concern breeding, Stable non-breeding)

In eastern North America, brown thrashers nest in thickets, hedgerows, forest edges, and overgrown clearings in deciduous forest. They're often found in woodlands with cottonwood, willow, dogwood, American plum, saltcedar, hawthorn, pitch pine, or scrub oak. On rare occasions they breed in backyards and gardens, although they are more likely to breed in suburban settings in the western part of their range. (Cornell Lab of Ornithology, 2015-a; Ehrlich et. al., 1988).

Cattle Egret

(Threatened breeding, Special Concern non-breeding)

Cattle Egrets breed in coastal barrier islands, marshes, reservoirs, lakes, quarries, swamps, riverside woodlands, and upland forests. They usually nest in colonies already established by native herons and egrets, and forage in fields with grazing livestock. During spring and fall migration they stop along marine shorelines as well as in farm fields (Cornell Lab of Ornithology, 2015-b).

Common Tern

(Special Concern breeding)

Common terns utilize a variety of coastal habitats, including sand and shell beaches, grassy uplands, or rocky island shores. In some areas, the birds have resorted to nesting on spoil banks. Availability of suitable nesting sites for this colonial species is a limiting factor to its success in the eastern United States (Harrison, 1975).

Glossy Ibis

(Special Concern breeding)

The glossy ibis breeds in colonies along the coast, often associating with other species such as herons. A variety of habitat conditions may be utilized including fresh, brackish or saltwater settings. Rookeries may be found in swamps, marshes or on islands (Harrison, 1975).

Least Tern

(State Endangered)

Least terns nest in colonies along barrier island beaches or mainland beach strands. They prefer bare or sparsely vegetated sandy areas just beyond the reach of normal spring tides. Sandy dredge disposal sites or sand piles near mining operations may also be utilized. The birds typically forage in bays, lagoons, estuaries, rivers and lakes along the coast (Beans and Niles, 2003).

Little Blue Heron

(Special Concern)

Little Blue Herons nest and forage in many kinds of wetlands, including swamps, marshes, ponds, streams, lagoons, tidal flats, canals, ditches, fish hatcheries, and flooded fields. They nest mostly in shrubs and small trees in standing water or upland sites on islands, including artificial islands created from dredged material. Rarely, they seek prey in upland pasture sites. They usually forage in water 2–6 inches deep, often gravitating toward densely vegetated foraging sites. In wintertime, Little Blue Herons make especially frequent use of salt ponds, mudflats, and savannas (Cornell Lab of Ornithology, 2015-c).

Osprey

(Threatened breeding, Stable non-breeding)

As a piscivorous species, the osprey is strictly associated with bodies of water that support adequate fish populations. Consequently, ospreys inhabit coastal rivers, marshes, bays and inlets as well as inland rivers, lakes and reservoirs. Ospreys nest on live or dead trees, artificial nesting platforms, light poles, channel markers, abandoned duck blinds, or other artificial structures that are in close proximity to fishing areas and offer an unobstructed view of the surrounding landscape. Territories typically contain poles, snags, or structures near the nest on which the ospreys perch. (Beans and Niles, 2003).

Red-shouldered Hawk

(Endangered breeding, Special Concern non-breeding)

Mature wet woods such as hardwood swamps and riparian forests typify red-shouldered hawk breeding habitat. Nesting territories, which occur in deciduous, coniferous, or mixed woodlands, are typically located within remote and extensive old growth forests containing standing water. Red-shouldered hawks select large deciduous and, to a lesser extent, coniferous trees for nesting. Forest characteristics include a closed canopy of tall trees, an open subcanopy, and variable amounts of understory cover. An-area sensitive species, the red-shouldered hawk typically nests away from residences, roads, and development. During the nonbreeding season, red-shouldered hawks are less restrictive in their habitat use. They inhabit the traditional wetland forests occupied during the breeding season as well as uplands, fragmented woods, smaller forests, open areas, and edges. (NJENSP, undated-b).

Snowy Egret

(Special Concern breeding)

Snowy Egrets nest in colonies on thick vegetation in isolated places such as barrier islands, dredge-spoil islands, salt marsh islands, swamps, and marshes. They often change location from year to year. During the breeding season Snowy Egrets feed in estuaries, saltmarshes, tidal channels, and shallow bays. They winter in freshwater swamps, grassy ponds, and temporary pools, and forage on beaches, shallow reefs, and wet fields (Cornell Lab of Ornithology, 2015-d).

Tricolored Heron

(Special Concern)

Tricolored herons, also known as Louisiana herons, may breed in both inland freshwater and saltwater habitats, although it prefers the latter. Like other herons, it is a colonial nester and may be found in mixed-species colonies located on islands, in marshes, or adjacent to ponds, lakes and rivers (Ehrlich et. al, 1988; Harrison 1975).

Precious Underwing

(not currently listed but tracked by Endangered and Nongame Species Program)

The precious underwing is a moth that inhabits mesic to wet hardwood forests, including riparian and seepage swamps. Best habitats are usually headwaters swamps with an abundant supply of a preferred food plant in the understory. Larval host plants are members of the rose family including red chokeberry (*Aronia arbutifolia*), beach plum (*Prunus maritima*), serviceberry (*Amelanchier sp.*), and apple (*Malus sp.*) (Natureserve, 2015; BugGuide, 2017).

Southeastern Tiger Beetle

(not currently listed but tracked by Endangered and Nongame Species Program)

The southeastern tiger beetle may be found on open, pristine white sandy beaches along the Atlantic coast during the summer months. Adults run actively on sand to hunt, while larvae hide in burrows with only their jaws protruding to catch unwary prey. Therefore, larvae need a safe place for development above the high tide line, and both stages require a large enough expanse of naturally shifting ocean beach so that sufficient prey is available. (SCDNR, 2015; BugGuide, 2017).

Piping Plover

(Federally Threatened, State Endangered)

Piping plovers inhabit ocean-front beaches and barrier islands, typically nesting on the stretch of beach between dunes and the high-tide line. Nests are often located in sparsely vegetated areas with shell fragments. During the non-breeding season, the plovers inhabit coastal beaches, barrier islands, inlets, sand flats, mudflats and dredged material islands. Intertidal beaches, washover areas, exposed mud and sand flats, wracklines and shorelines are preferred foraging sites (Beans and Niles, 2003).

American Oystercatcher

(Special Concern)

During the breeding season, American Oystercatchers may be found in coastal habitats including sand or shell beaches, dunes, salt marshes, marsh islands, mudflats, and dredge spoil islands made of sand or gravel. During migration and winter, look for them feeding in mud or sand flats exposed by the tide, or on shellfish beds. Oystercatchers tend to roost on beaches, dunes, or marsh islands near their foraging sites, and rarely venture far inland (Cornell Lab of Ornithology, 2015-e).

Bronze Copper

(State Endangered)

The bronze copper is a butterfly that inhabits moist or wet areas including brackish and freshwater marshes, bogs, fens, seepages, wet sedge meadows, riparian zones, wet grasslands and drainage ditches. Various kinds of dock (*Rumex spp.*) are the preferred larval host plants, and adults nectar on an assortment of wildflowers (Beans and Niles, 2003).

Citations:

Beans, B.E. and L. Niles. 2003. Endangered and Threatened Wildlife of New Jersey. Rutgers University Press, New Brunswick, New Jersey.

BugGuide. 2017. Identification, Images and Information for Insects, Spiders and their kin for the United States and Canada. A shared content website hosted by the Iowa State University Department of Entomology. <http://bugguide.net/node/view/15740>. Accessed March 2017.

Cornell Lab of Ornithology. 2015-a. All About Birds: Brown Thrasher.
https://www.allaboutbirds.org/guide/Brown_Thrasher/lifehistory

Cornell Lab of Ornithology. 2015-b. All About Birds: Cattle Egret.
https://www.allaboutbirds.org/guide/Cattle_Egret/lifehistory

Cornell Lab of Ornithology. 2015-c. All About Birds: Little Blue Heron.
https://www.allaboutbirds.org/guide/Little_Blue_Heron/lifehistory

Cornell Lab of Ornithology. 2015-d. All About Birds: Snowy Egret.
https://www.allaboutbirds.org/guide/Snowy_Egret/lifehistory

Cornell Lab of Ornithology. 2015-e. All About Birds: American Oystercatcher.
https://www.allaboutbirds.org/guide/American_Oystercatcher/lifehistory

Ehrlich, P.R., D.S. Dobkin and D. Wheye. 1988. The Birder's Handbook: A Field Guide to the Natural History of North American Birds. Simon & Schuster Inc., New York, New York.

Harrison, H.H. 1975. A Field Guide to Birds' Nests , United States East of the Mississippi River. Houghton Mifflin Company, New York, New York.

Natureserve. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. <http://explorer.natureserve.org>. Accessed March 2017.

NJENSP (Endangered and Nongame Species Program). Undated -a. Bobolink Fact Sheet. <http://www.nj.gov/dep/fgw/ensp/pdf/end-thrtened/bobolink.pdf> . Accessed February 2017.

NJENSP (Endangered and Nongame Species Program). Undated -b. Red-shouldered Hawk Fact Sheet. <http://www.nj.gov/dep/fgw/ensp/pdf/end-thrtened/redshldhwk.pdf>. Accessed February 2017.

SCDNR (South Carolina Department of Natural Resources). 2015. White Beach Tiger Beetle (*Cicindela dorsalis media*). Species summary by J.C. Ciegler and J. Holmes.
<http://www.dnr.sc.gov/cwcs/pdf/tigerbeetle.pdf>

USFWS (U.S. Fish and Wildlife Service). 2015. Bald and Golden Eagle Information. U.S. Fish & Wildlife Service, Migratory Bird Program. <https://www.fws.gov/birds/management/managed-species/bald-and-golden-eagle-information.php> Site updated September 25, 2015.

D.4 Rare Bird Species Documented in Cape May County

This is a subset of the full county list provided in Appendix C.1.

ENDANGERED		
E	Piping Plover	<i>Charadrius melodus</i>
E	Upland Sandpiper	<i>Bartramia longicauda</i>
E	Least Tern	<i>Sternula antillarum</i>
E	Roseate Tern	<i>Sterna dougallii</i>
E	Black Skimmer	<i>Rynchops niger</i>
E	Sedge Wren	<i>Cistothorus platensis</i>
E	Henslow's sparrow	<i>Ammodramus henslowii</i>
Ebr, SCnb	Pied-billed Grebe	<i>Podilymbus podiceps</i>
Ebr, SCnb	American Bittern	<i>Botaurus lentiginosus</i>
Ebr, SCnb	Northern Harrier	<i>Circus cyaneus</i>
Ebr, SCnb	Northern Goshawk	<i>Accipiter gentilis</i>
Ebr, SCnb	Red-shouldered Hawk	<i>Buteo lineatus</i>
Ebr, SCnb	Short-eared Owl	<i>Asio flammeus</i>
Ebr, SCnb	Peregrine Falcon	<i>Falco peregrines</i>
Ebr, SCnb	Vesper Sparrow	<i>Poocetes gramineus</i>
Ebr, Tnb	Black rail	<i>Laterallus jamaicensis</i>
Ebr, Tnb	Bald Eagle	<i>Haliaeetus leucocephalus</i>
Ebr,SCnb	Golden-winged Warbler	<i>Vermivora chrysoptera</i>
Enb	Red Knot	<i>Calidris canutus</i>
Enb	Loggerhead shrike	<i>Lanius ludovicianus</i>
THREATENED		
T	Yellow-crowned Night-Heron	<i>Nyctanassa violacea</i>
T	Barred Owl	<i>Strix varia</i>
T	Long-eared Owl	<i>Asio otus</i>
T	American Kestrel	<i>Falco sparverius</i>
Tbr	Osprey	<i>Pandion haliaetus</i>
Tbr	Savannah Sparrow	<i>Passerculus sandwichensis</i>
Tbr, SCnb	Cattle Egret	<i>Bubulcus ibis</i>
Tbr, SCnb	Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>
Tbr, SCnb	Horned Lark	<i>Eremophila alpestris</i>
Tbr, SCnb	Grasshopper Sparrow	<i>Ammodramus savannarum</i>
Tbr, SCnb	Bobolink	<i>Dolichonyx oryzivorus</i>
SPECIAL CONCERN		
SC	Common Nighthawk	<i>Chordeiles minor</i>
SC	American Oystercatcher	<i>Haematopus palliatus</i>
SC	Gull-billed Tern	<i>Gelochelidon nilotica</i>
SC	Least Bittern	<i>Ixobrychus exilis</i>
SC	Little Blue Heron	<i>Egretta caerulea</i>
SC	Tricolored Heron	<i>Egretta tricolor</i>

SC	Sharp-shinned Hawk	<i>Accipiter striatus</i>
SC	Kentucky Warbler	<i>Geothlypis formosa</i>
SCbr	Black-billed Cuckoo	<i>Coccyzus erythrophthalmus</i>
SCbr	Spotted Sandpiper	<i>Actitis macularius</i>
SCbr	Caspian Tern	<i>Hydroprogne caspia</i>
SCbr	Common Tern	<i>Sterna hirundo</i>
SCbr	Great Blue Heron	<i>Ardea herodias</i>
SCbr	Snowy Egret	<i>Egretta thula</i>
SCbr	Glossy Ibis	<i>Plegadis falcinellus</i>
SCbr	Cooper's Hawk	<i>Accipiter cooperii</i>
SCbr	Broad-winged Hawk	<i>Buteo platypterus</i>
SCbr	Least Flycatcher	<i>Empidonax minimus</i>
SCbr	Blue-headed Vireo	<i>Vireo solitarius</i>
SCbr	Cliff Swallow	<i>Petrochelidon pyrrhonota</i>
SCbr	Winter Wren	<i>Troglodytes hiemalis</i>
SCbr	Veery	<i>Catharus fuscescens</i>
SCbr	Wood Thrush	<i>Hylocichla mustelina</i>
SCbr	Brown Thrasher	<i>Toxostoma rufum</i>
SCbr	Worm-eating Warbler	<i>Helmitheros vermivorum</i>
SCbr	Nashville Warbler	<i>Oreothlypis ruficapilla</i>
SCbr	Hooded Warbler	<i>Setophaga citrina</i>
SCbr	Cerulean Warbler	<i>Setophaga cerulea</i>
SCbr	Northern Parula	<i>Setophaga americana</i>
SCbr	Blackburnian Warbler	<i>Setophaga fusca</i>
SCbr	Black-throated Blue Warbler	<i>Setophaga caerulescens</i>
SCbr	Black-throated Green Warbler	<i>Setophaga virens</i>
SCbr	Canada Warbler	<i>Cardellina canadensis</i>
SCbr	Yellow-breasted Chat	<i>Icteria virens</i>
SCbr	Saltmarsh Sparrow	<i>Ammodramus caudacutus</i>
SCbr	Eastern Meadowlark	<i>Sturnella magna</i>
SCbr, Unb	Eastern Whip-poor-will	<i>Antrostomus vociferous</i>
SCnb	Whimbrel	<i>Numenius phaeopus</i>
SCnb	Sanderling	<i>Calidris alba</i>
SCnb	Semipalmated Sandpiper	<i>Calidris pusilla</i>
SCnb	Gray-cheeked Thrush	<i>Catharus minimus</i>

D.5 Rare Plant Species of Cape May County.

Species shown in **red** are known from West Cape May Borough, and those in **green** from the immediate vicinity.

Ferns & Allies

<i>Isoetes melanopoda</i>	black-base quillwort	mostly in shallow, temporary ponds or nearly terrestrial
<i>Ophioglossum pusillum</i>	northern adder's tongue	grassy marsh edges and ditches
<i>Ophioglossum vulgatum</i>	southern adder's-tongue	low woods and floodplains
<i>Schizaea pusilla</i>	curly grass fern	in sandy, acid soils of white cedar bog edges and hummocks, just above the water line

Graminoids

<i>Carex jorii</i>	cypress-swamp sedge	wet woods and swamps, mainly on the coastal plain
<i>Carex mitchelliana</i>	Mitchell's sedge	swamp forests and wet thickets and meadows
<i>Carex silicea</i>	seabeach sedge	sand or sandy soil near the coast
<i>Cyperus lancastricensis</i>	Lancaster flat sedge	woods and fields
<i>Cyperus polystachyos</i>	coast flat sedge	wet soil
<i>Cyperus refractus</i>	reflexed flat sedge	woods and fields
<i>Cyperus retrofractus</i>	rough flat sedge	pastures, roadsides and other disturbed places
<i>Eleocharis brittonii</i>	Britton's spike-rush	acid swamps near the coast
<i>Eleocharis halophila</i>	salt-marsh spike-rush	mainly of coastal salt marshes
<i>Eleocharis melanocarpa</i>	black-fruit spike-rush	wet sand and pine barrens, mostly near the coast
<i>Eleocharis quadrangulata</i>	angled spike-rush	shallow water
<i>Eleocharis tortilis</i>	twisted spike-rush	swamps and bogs, chiefly near the coast
<i>Eriophorum tenellum</i>	rough cotton-grass	bogs and swamps, often under conifers
<i>Fimbristylis castanea</i>	marsh fimbry	brackish coastal marshes, seldom in inland alkaline sites
<i>Fuirena squarrosa</i>	hairy umbrella-sedge	moist or wet, often sandy places
<i>Rhynchospora cephalantha</i>	large-headed beaked-rush	wet acid soil on the coastal plain
<i>Rhynchospora filifolia</i>	thread-leaf beaked-rush	pine-barren bogs on the coastal plain
<i>Rhynchospora glomerata</i>	clustered beaked-rush	bogs and wet sand
<i>Rhynchospora inundata</i>	slender horned-rush	swamps and pond margins
<i>Rhynchospora nitens</i>	short-beaked bald-rush	wet sandy soil and bogs
<i>Rhynchospora pallida</i>	pale beaked-rush	acid bogs along the coast
<i>Rhynchospora rariflora</i>	rare-flower beaked-rush	bogs and wet soil on the coastal plain
<i>Rhynchospora recognita</i>	coarse grass-like beaked-rush	low, wet to moist ground in swamps and bogs, sandy depressions
<i>Rhynchospora scirpoides</i>	long-beak bald-rush	wet sandy soil
<i>Schoenoplectus novae-angliae</i>	New England bulrush	brackish transitional zones of tidal river systems
<i>Scirpus longii</i>	Long's woolgrass	marshes near the coast
<i>Scleria verticillata</i>	whorled nut-rush	wet sandy soil
<i>Juncus articulatus</i>	jointed rush	bogs, wet meadows and shores
<i>Juncus brachycarpus</i>	short-fruit rush	damp or wet soil
<i>Juncus caesariensis</i>	New Jersey rush	sphagnum bogs in the pine barrens
<i>Juncus coriaceus</i>	awl-leaf rush	wet ground, swamps or brackish marshes
<i>Juncus diffusissimus</i>	slim-pod rush	wet soil and muddy shores

<i>Juncus elliotii</i>	Elliott's rush	damp or wet, sandy or peaty soil, especially in pine barrens
<i>Juncus torreyi</i>	Torrey's rush	wet sunny places or shallow standing water, especially on prairies
<i>Andropogon glomeratus</i> <i>var hirsutior</i>	hairy beardgrass	wet, open soil
<i>Andropogon gyrans</i>	Elliott's beardgrass	fields and open woods
<i>Andropogon ternarius</i> <i>var ternarius</i>	silvery beardgrass	open woods and dry fields
<i>Aristida virgata</i>	wand-like three-awn grass	dry sandy soil of the coastal plain
<i>Coelorachis rugosa</i>	wrinkled jointgrass	wet pine barrens
<i>Gymnopogon brevifolius</i>	short-leaf skeleton grass	sandy or peaty dry ground of bog margins, pinelands
<i>Leptochloa fascicularis</i> <i>var maritima</i>	long-awn sprangletop	wet soil of watersides, fresh or usually brackish water of marshes
<i>Muhlenbergia torreyana</i>	Pine Barren smoke grass	wet, open soil of bogs, sandy pinelands
<i>Panicum aciculare</i>	bristling panic grass	dry to moist sandy soil of the coastal pinelands
<i>Panicum dichotomum</i> <i>var roanokense</i>	bluish panic grass	marshes, swampy woods, esp in pinelands
<i>Panicum hemitomon</i>	maiden-cane	moist soil or shallow water of swamp edges, watersides, wet fields, ponds
<i>Panicum leucothrix</i>	rough panic grass	wet, sandy soil of coastal pinelands, bogs
<i>Panicum longiligulatum</i>	coastal-plain panic grass	moist or dry sites, opens woods, dunes, shores and prairies
<i>Panicum tenue</i>	white-edge panic grass	damp sandy soil of the coastal plain
<i>Panicum wrightianum</i>	Wright's panic grass	moist or wet, sandy or peaty soil of watersides, bogs, beaches
<i>Paspalum dissectum</i>	mudbank crown grass	usually in wet, muddy soil or in shallow water of watersides, also in dried pond bottoms
<i>Paspalum floridanum</i>	Florida crown grass	sandy, usually moist open ground, brackish marshes
<i>Puccinellia fasciculata</i>	saltmarsh alkali grass	wet soil or water of salt marsh borders, beaches
<i>Sacciolepis striata</i>	American cupscale	shallow water or muddy ground of swamps, ditches
<i>Setaria magna</i>	giant fox-tail	low, wet disturbed soil in fresh to brackish conditions, found in marshes and swamps
<i>Sphenopholis pensylvanica</i>	swamp oats	swamps and wet woods
<i>Sporobolus clandestinus</i>	rough rush-grass	dry, sandy to rocky ground of woods edges, meadows
<i>Tridens flavus</i> <i>var chapmanii</i>	Chapman's redtop	dry open woods and shaded edges, disturbed ground, fields
<i>Vulpia ellioatea</i>	squirrel-tail six-weeks grass	dry sandy soils

Terrestrial Herbs

<i>Ruellia caroliniensis</i>	Carolina petunia	moist or dry woods
<i>Sesuvium maritimum</i>	seabeach purslane	sea beaches
<i>Sagittaria australis</i>	southern arrowhead	mostly in circumneutral water of lakes, ponds or swamps
<i>Sagittaria teres</i>	slender arrowhead	sandy soil in shallow acid water
<i>Amaranthus pumilus</i>	seabeach amaranth	sea-beaches
<i>Centella erecta</i>	erect coinleaf	marshes and wet soil along the coast
<i>Eryngium aquaticum</i> <i>var aquaticum</i>	marsh rattlesnake-master	bogs and marshes near the coast
<i>Hydrocotyle prolifera</i>	Canby's marsh-pennywort	wet or moist ground of bogs, swampy woods, watersides


<i>Hydrocotyle verticillata</i> <i>var verticillata</i>	whorled marsh-pennywort	wet or moist ground of bogs, swampy woods, watersides
<i>Asclepias lanceolata</i>	smooth orange milkweed	swamps, bogs and brackish marshes on the coastal plain
<i>Asclepias rubra</i>	red milkweed	swamps, bogs and wet woods on or near the coastal plain
<i>Asclepias variegata</i>	white milkweed	upland woods and thickets
<i>Asclepias verticillata</i>	whorled milkweed	fields, roadsides, upland woods and prairies
<i>Aster concolor</i>	eastern silvery aster	dry sandy places, often among pines
<i>Aster ericoides</i> <i>var ericoides</i>	white heath aster	dry open places
<i>Bidens mitis</i>	small-fruit beggar-ticks	swamps and other wet places
<i>Boltonia asteroides</i> <i>var glastifolia</i>	southern boltonia	moist or wet places
<i>Cirsium virginianum</i>	Virginia thistle	savannas, bogs and wet pinelands
<i>Eupatorium album</i> <i>var vaseyi</i>	Vasey's boneset	dry open woods, especially in sandy pinelands
<i>Eupatorium aromaticum</i> <i>var aromaticum</i>	smaller white snakeroot	dry woods, especially in sandy soil
<i>Eupatorium capillifolium</i>	dog-fennel thoroughwort	open places, often in old fields and pastures
<i>Eupatorium coelestinum</i>	mist-flower	woods, streambanks, meadows and fields
<i>Eupatorium resinosum</i>	Pine Barren boneset	pocosins, bogs and other wet places, often in pine barrens
<i>Gnaphalium helleri</i> <i>var micradenium</i>	small everlasting	dry, commonly sandy soil, often in woods
<i>Hieracium marianum</i>	Maryland hawkweed	mostly in dry open woods
<i>Krigia dandelion</i>	potato dwarf-dandelion	low prairies, fields and other moist, chiefly open places
<i>Pluchea foetida</i>	stinking fleabane	permanently wet soil, often in meadows or swampy woods
<i>Prenanthes autumnalis</i>	Pine Barren rattlesnake-root	sandy, usually moist places, often among pines
<i>Senecio tomentosus</i>	wooly ragwort	dry open places and pine woods, especially in sandy soil
<i>Smallanthus uvedalius</i>	bear's-foot	woods and meadows
<i>Solidago elliotii</i>	Elliott's goldenrod	fresh or brackish swamps near the coast
<i>Solidago uliginosa</i> <i>var uliginosa</i>	bog goldenrod	bogs, acid swamps, meadows and moist to dryish thickets
<i>Arabis drummondii</i>	Drummond's rockcress	various habitats
<i>Polyprenum procumbens</i>	juniper-leaf	dry sandy soil and waste places
<i>Lobelia boykinii</i>	Boykin's lobelia	ponds and swamps on the coastal plain
<i>Lobelia canbyi</i>	Canby's lobelia	swamps on the coastal plain
<i>Honckenya peploides</i> <i>var robusta</i>	seabeach sandwort	sea beaches and sand dunes
<i>Chenopodium berlandieri</i> <i>var macrocalycium</i>	large-calyx goosefoot	mainly coastal or coastal plain, often on sea beaches
<i>Suaeda rolandii</i>	Roland's seablite	saline or alkaline soil
<i>Hypericum adpressum</i>	Barton's St. John's-wort	marshes, shores and wet meadows
<i>Triadenum walteri</i>	Walter's St. John's-wort	swamps and marshes on the coastal plain
<i>Cuscuta indecora</i>	collared dodder	on a wide variety of hosts
<i>Cuscuta polygonorum</i>	smartweed dodder	on Polygonum and other hosts
<i>Croton willdenowii</i>	elliptical rushfoil	in sand, sandy soil or rocky barrens
<i>Euphorbia purpurea</i>	Darlington's glade spurge	dry or moist woods

<i>Aeschynomene virginica</i>	sensitive joint-vetch	sandy or muddy riverbanks and fresh to brackish tidal shores
<i>Centrosema virginianum</i>	spurred butterfly-pea	dry sandy woods and barrens
<i>Clitoria mariana</i>	butterfly-pea	dry upland woods and barrens
<i>Desmodium strictum</i>	pineland tick-trefoil	pine barrens on the coastal plain
<i>Desmodium viridiflorum</i>	velvety tick-trefoil	dry woods
<i>Galactia volubilis</i>	downy milk-pea	dry upland woods and barrens
<i>Lespedeza stuevei</i>	Stueve's downy bush-clover	dry upland woods and barrens
<i>Lupinus perennis</i> var <i>perennis</i>	sundial lupine	dry open woods and clearings
<i>Phaseolus polystachios</i> var <i>polystachios</i>	wild kidney bean	moist woods and thickets
<i>Stylosanthes biflora</i>	pencil-flower	dry or rocky woods and barrens
<i>Gentiana autumnalis</i>	Pine Barren gentian	wet, open sandy ground of pinelands and edges, roadsides
<i>Sabatia dodecandra</i> var <i>dodecandra</i>	large marsh-pink	salt or brackish marshes near the coast
<i>Sisyrinchium fuscatum</i>	sand plain blue-eyed grass	sandy areas, mainly near the coast
<i>Lycopus rubellus</i>	stalked water-hoarhound	low wet or moist ground of marshes, alluvial woods and edges
<i>Trichostema setaceum</i>	narrow-leaf bluecurls	dry, open sandy or rocky ground of thin woods fields or sterile disturbed soils
<i>Helonias bullata</i>	swamp-pink	swamps and bogs
<i>Linum intercursum</i>	sandplain flax	dry or moist, sandy, open ground of thin woods and shaded edges
<i>Ammannia latifolia</i>	Koehn's toothcup	wet ground on the coastal plain
<i>Rotala ramosior</i>	tooth-cup	wet, open muddy or sandy ground of watersides, depressions, swamps
<i>Rhexia aristosa</i>	awned meadow-beauty	local in sandy or peaty acid bogs on the coastal plain
<i>Rhexia mariana</i> var <i>ventricosa</i>	showy meadow-beauty	moist, open places, often in disturbed habitats
<i>Epilobium angustifolium</i> ssp <i>circumvagum</i>	narrow-leaf fireweed	many habitats, esp. moist soils rich in humus, often abundant after fires
<i>Ludwigia hirtella</i>	hairy primrose-willow	swamps on the coastal plain
<i>Ludwigia linearis</i>	narrow-leaf primrose-willow	wet soil, muddy shores or shallow waters
<i>Oenothera humifusa</i>	sea-beach evening-primrose	dry sand of upper sea beaches, dunes and hollows
<i>Oenothera oakesiana</i>	Oakes' evening-primrose	disturbed open places
<i>Arethusa bulbosa</i>	dragon mouth	sphagnum bogs and swampy meadows
<i>Cleistes divaricata</i>	spreading pogonia	swamps and wet woods, mostly on the coastal plain
<i>Listera australis</i>	southern twayblade	shaded bogs and wet woods, mainly on the coastal plain
<i>Platanthera ciliaris</i>	yellow fringed orchid	bogs, fields and woods
<i>Platanthera cristata</i>	crested yellow orchid	low moist meadows and damp pine woods, especially along the coastal plain
<i>Platanthera flava</i> var <i>flava</i>	southern rein orchid	boggy or swampy ground and floodplains
<i>Platanthera integra</i>	yellow fringeless orchid	open acid bogs and pine barrens on the coastal plain
<i>Platanthera nivea</i>	snowy orchid	open acid bogs on the coastal plain
<i>Platanthera peramoena</i>	purple fringeless orchid	open, swampy or vernal wet places, often in acid soil
<i>Spiranthes laciniata</i>	lace-lip ladies'-tresses	open wet places on the coastal plain

<i>Spiranthes lucida</i>	shining ladies'-tresses	damp woods, marshes and wet shores; calciphile
<i>Spiranthes odorata</i>	fragrant ladies'-tresses	open moist, often sandy places; mainly on the coastal plain, often in estuaries
<i>Spiranthes tuberosa</i>	little ladies'-tresses	acid, usually rather dry soil
<i>Plantago maritima</i> var <i>juncooides</i>	seaside plantain	salt or brackish conditions, in water or wet soil of marshes, swamps, beaches, tidal streams, headlands
<i>Plantago pusilla</i>	dwarf plantain	dry, sandy open ground of woods edges, fields, dunes
<i>Phlox maculata</i> var <i>maculata</i>	spotted phlox	moist or wet, low ground of streamsides, meadows, floodplains
<i>Polygala mariana</i>	Maryland milkwort	dry, usually sandy ground but sometimes in clay or peat
<i>Polygala ramosa</i>	low Pine Barren milkwort	water of pineland swamps, wet fields
<i>Polygonum densiflorum</i>	dense-flower knotweed	shallow water or wet ground of shaded swamps, pool edges, woods edges
<i>Polygonum glaucum</i>	sea-beach knotweed	dry open sand of beaches, dune hollows, saline pond edges, fill dirt
<i>Polygonum hydropiperoides</i> var <i>opelousanum</i>	Opelousas water-pepper	shallow water and wet ground of alluvial woods and edges, swamps, beaches, watersides
<i>Polygonum setaceum</i> var <i>setaceum</i>	bristly smartweed	shallow water and wet ground of alluvial woods and edges, swamps, beaches, watersides
<i>Ranunculus cymbalaria</i>	seaside buttercup	in mud, especially in brackish or alkaline places
<i>Diodia virginiana</i>	larger buttonweed	moist, open low ground of watersides and ditches
<i>Galium hispidulum</i>	coast bedstraw	sandy soil, esp pineland on the coastal plain
<i>Gratiola pilosa</i>	hairy hedge-hyssop	sandy, usually moist ground of salt marsh edges, pinelands, open woods & shaded edges
<i>Gratiola virginiana</i>	round-fruit hedge-hyssop	wet woods
<i>Schwalbea americana</i>	chaffseed	moist to dry, sandy ground of pine-oak woods and shaded edges, marshes
<i>Valerianella radiata</i>	beaked cornsalad	moist ground of streamsides, fields, meadows, woods and edges
<i>Phyla lanceolata</i>	fogfruit	moist low ground and muddy flats, both coastal and inland
<i>Verbena simplex</i>	narrow-leaf vervain	dry, open, sandy or rocky non-acid ground of thin woods, fields, roadsides
<i>Viola brittoniana</i> var <i>brittoniana</i>	Britton's coast violet	low, wet or moist, sandy or peaty ground, oft in brackish conditions, in woods edges & meadows
<i>Viola brittoniana</i> var <i>pectinata</i>	cut-leaf coast violet	low, wet or moist, sandy or peaty ground, oft in brackish conditions, in woods edges & meadows
<i>Xyris jupicai</i>	Richard's yellow-eyed grass	wet, sunny disturbed sites, often weedy
Aquatic Herbs		
<i>Sclerolepis uniflora</i>	bog buttons	shallow water on the coastal plain
<i>Ceratophyllum echinatum</i>	spiny coontail	quiet water
<i>Elatine americana</i>	American waterwort	muddy tidal shores
<i>Elatine minima</i>	small waterwort	on mud
<i>Eriocaulon parkeri</i>	Parker's pipewort	tidal flats and muddy shores, oft submerged in fresh to slightly brackish water
<i>Myriophyllum pinnatum</i>	cutleaf water-milfoil	submersed in quiet water or rooting on muddy shores
<i>Lemna perpusilla</i>	minute duckweed	floating on quiet waters of ponds, streams, springs, lakes
<i>Utricularia gibba</i>	humped bladderwort	shallow water

<i>Utricularia minor</i>	lesser bladderwort	shallow water
<i>Utricularia purpurea</i>	purple bladderwort	quiet shallow water of ponds, slow streams, ditches
<i>Utricularia radiata</i>	small swollen bladderwort	quiet shallow water of ponds, ditches
<i>Utricularia resupinata</i>	reversed bladderwort	very shallow water or wet mud of watersides
<i>Nuphar microphyllum</i> (<i>lutea</i> ssp. <i>Pumila</i>)	small yellow pond-lily	ponds
<i>Potamogeton oakesianus</i>	Oakes' pondweed	in quiet shallow waters of ponds; soils usually acid, sandy or rocky
<i>Hottonia inflata</i>	featherfoil	quiet shallow water or wet soil of swamps, slow streams, ditches
Shrubs		
<i>Rhododendron atlanticum</i>	dwarf azalea	sandy soil on the coastal plain
<i>Malus angustifolia</i> var <i>puberula</i>	spiny wild crabapple	woods and thickets
<i>Prunus angustifolia</i> var <i>angustifolia</i>	Chickasaw plum	dry, open sandy or sterile ground of woods and shaded edges, dune sands
<i>Rubus gnarus</i>	Pollock's mill blackberry	state historic from one occurrence
<i>Rubus longii</i>	Long's blackberry	dry sandy soil, chiefly on the coastal plain
<i>Rubus novocaesarius</i>	New Jersey dewberry	worldwide occurrence restricted to a single occurrence in Cape May County
Trees		
<i>Quercus michauxii</i>	basket oak	low or wet soil, especially alluvial flood plains, on the coastal plain
<i>Quercus nigra</i>	water oak	usually in damp or wet soil
<i>Fraxinus profunda</i>	pumpkin ash	swamps and wet woods
<i>Pinus serotina</i>	pond pine	swamps and wet soil
<i>Pinus taeda</i>	loblolly pine	moist, sandy soil
<i>Populus heterophylla</i>	swamp cottonwood	wet low woods and swamps

D.6 Cape May Migratory Bird Refuge Natural Heritage Priority Site



APPENDIX E: INVASIVE SPECIES

Species Tracked by the New Jersey Invasive Species Strike Team in Cape May County.

Common Name	Scientific Name	Taxa
<u>ANIMALS</u>		
mute swan		bird
<u>PLANTS</u>		
Black locust	<i>Robinia pseudoacacia</i>	tree
Callery pear (Bradford pear)	<i>Pyrus calleryana</i>	tree
Mimosa	<i>Albizia julibrissin</i>	tree
Norway maple	<i>Acer platanoides</i>	tree
Paper-mulberry	<i>Broussonetia papyrifera</i>	tree
Tree-of-heaven	<i>Ailanthus altissima</i>	tree
Amur honeysuckle	<i>Lonicera mackii</i>	shrub
Autumn olive	<i>Elaeagnus umbellata</i>	shrub
Butterflybush	<i>Buddleja davidii</i>	shrub
European privet	<i>Ligustrum vulgare</i>	shrub
Multiflora rose	<i>Rosa multiflora</i>	shrub
Privet	<i>Ligustrum sp.</i>	shrub
Seaside rose	<i>Rosa rugosa</i>	shrub
Trifoliolate orange	<i>Citrus trifoliata</i>	shrub
Chinese wisteria	<i>Wisteria sinensis</i>	vine
English ivy	<i>Hedera helix</i>	vine
Japanese honeysuckle	<i>Lonicera japonica</i>	vine
Kudzu	<i>Pueraria montana var. lobata</i>	vine
Mile-a-minute vine	<i>Persicaria perfoliata</i>	vine
Oriental bittersweet	<i>Celastrus orbiculatus</i>	vine
Porcelain-berry	<i>Ampelopsis brevipedunculata</i>	vine
Sweet autumn virginbower	<i>Clematis terniflora</i>	vine
Common reed	<i>Phragmites australis</i>	grass
Weeping lovegrass	<i>Eragrostis curvula</i>	grass
Chinese bush clover	<i>Lespedeza cuneata</i>	herb
Garlic mustard	<i>Alliaria petiolata</i>	herb
Japanese knotweed	<i>Fallopia japonica</i>	herb
Lesser celandine, fig buttercup	<i>Ficaria verna</i>	herb
Mugwort	<i>Artemesia vulgaris</i>	herb
Purple loosestrife	<i>Lythrum salicaria</i>	herb
Yellow iris	<i>Iris pseudacorus</i>	herb
Spotted knapweed	<i>Centaurea stoebe ssp. micranthos</i>	plant
Parrotfeather	<i>Myriophyllum aquaticum</i>	aquatic
Data from NJISST (undated). http://www.njisst.org/		

